



Turf Products

Fertilizer 22-0-3

Net Wt. 50 Lbs. (22.68 kg)

- Contains Fortify®-N

Guaranteed Analysis

Total Nitrogen (N)	22.00%
22.00% Urea Nitrogen*	
Soluble Potash (K ₂ O)	3.00%
Magnesium (Mg)	2.00%
0.00% Water Soluble Magnesium	
Iron (Fe)	2.00%

Derived from urea, potassium chloride, magnesium oxide and ferric oxide.

Chlorine (Cl) Max.	2.54%
--------------------	-------

*11.00% urea nitrogen stabilized with N-(n-butyl) thiophosphoric triamide and dicyandi-
amide

F002091

Nitrogen Notice: For those states with Nitrogen application restrictions, including but not limited to, MD, NH, NJ, VA, do not apply more than 3.4 lbs. of this product per 1,000 sq. ft.

DIRECTIONS FOR USE:

Apply only as directed. Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

Application Rates:

LIGHT:	3.4 lbs/1,000 sq. ft.
LOW:	4.5 lbs/1,000 sq. ft.
MEDIUM:	5.6 lbs/1,000 sq. ft.
HIGH:	6.8 lbs/1,000 sq. ft.

FLORIDA APPLICATIONS:

We recommend that you follow the Florida Green Industries BMP's at:
<http://www.flrules.org/Gateway/reference.asp?No=Ref-04706>

We recommend that you follow the Florida Golf Course BMP's at:
<http://www.flrules.org/Gateway/reference.asp?No=Ref-04702>

For all applications:

- Apply to dry foliage.
- Water in after application with a minimum of 0.2 inches to help move particles into the turf canopy.
- Do not collect clippings for at least one mowing following application to avoid particle pick-up.

Cool Season Grasses:

- Apply at the LOW or MEDIUM RATE every 6-8 weeks during the periods of active growth.
- Apply at the LIGHT RATE when temperatures exceed 85°F.

Warm Season Grasses:

- Apply at the LOW to HIGH RATE every 4-6 weeks during the period of active growth.

STORAGE AND USE:

Store in a clean, dry place.
Reseal opened bag by folding top down and securing.
Use only in clean, dry spreaders.
Do not leave unused product in spreaders.

Notice: This product contains the micronutrient iron for extra greening and to help prevent iron deficiency. Iron may stain concrete and other surfaces. Misapplied product to these surfaces should be removed promptly by sweeping or blowing. Do not wash off with water.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS.

LABEL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Product of the U.S.A.
Manufactured by:
The Andersons, Inc.
P.O. Box 119 Maumee, OH 43537
www.andersonsplantnutrient.com

©2020 The Andersons, Inc.
The Andersons logo and Fortify are registered trademarks of The Andersons, Inc. 101620

FRONT

Suggested Spreader Settings						
These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product. 50 lbs. feeds 14,700 sq. ft. (0.34 acre) with 0.75 lbs. N/1,000 sq. ft. at the LIGHT RATE 50 lbs. feeds 11,100 sq. ft. (0.25 acre) with 1.00 lbs. N/1,000 sq. ft. at the LOW RATE 50 lbs. feeds 8,900 sq. ft. (0.20 acre) with 1.23 lbs. N/1,000 sq. ft. at the MEDIUM RATE 50 lbs. feeds 7,300 sq. ft. (0.17 acre) with 1.50 lbs. N/1,000 sq. ft. at the HIGH RATE						
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS			
			LIGHT RATE 3.4 lbs/1,000 sq. ft.	LOW RATE 4.5 lbs/1,000 sq. ft.	MEDIUM RATE 5.6 lbs/1,000 sq. ft.	HIGH RATE 6.8 lbs/1,000 sq. ft.
Andersons Model 2000 / AccuPro 2000 (cone 4); Andersons 2000 SR / SR-2000 (cone 4)	3 mph	10 feet	L 1/2	M 1/2	O	O 1/2
Andersons LCO-1000	3 mph	10 feet	L 1/2	M 1/2	O	O 1/2
Scotts R-8, R-8A (cone 4)	3 mph	10 feet	K 1/4	L 1/2	M 3/4	O 1/4
Lesco 020093 (pattern 0.25)	3 mph	7 feet	E	E 1/2	F	F 1/2
Lesco 705698 (pattern 0.25)	3 mph	7 feet	12	13	14	15
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm) Pattern II-A	4.5 mph	38 feet	5 1/2	6 1/4	7	7 3/4
Vicon (03 Series)	4.5 mph	30 feet	28	33	37	41



Turf Products Fertilizer 22-0-3



WARNING

CAUSES SKIN, RESPIRATORY AND SERIOUS EYE IRRITATION

PRECAUTIONARY STATEMENTS

Wear protective gloves and clothing. Remove contaminated clothing and wash before reuse. Do not breathe dusts. Use only outdoors or in well-ventilated area. Call poison control or seek medical attention if you feel unwell. Wash thoroughly after handling. Do not eat, drink or smoke when using this product. Store container tightly closed. Dispose of contents/container in accordance with National/Regional/Local regulations.

FIRST AID

If swallowed:

Rinse mouth. DO NOT INDUCE VOMITING unless directed by a medical professional. Seek medical attention if unwell.

If on skin:

Wash affected areas with soap and water. Seek medical attention if irritation persists. Wash contaminated clothing before reuse.

If in eyes:

Immediately flush with water for at least 15-20 minutes, remove contact lenses if present and easy to do - continue rinsing. Seek medical attention if irritation persists.

If inhaled:

Move victim to fresh air and keep comfortable. Call POISON CENTER or doctor if unwell.

FPO

0 31865 31968 6

www.AndersonsPlantNutrient.com

APT22UM5.5

APT22UM5.5B