



GARDENLINE®

WEED and FEED with Trimec Herbicide .53% (15-5-10) FOR SENSITIVE GRASSES

Guaranteed Fertilizer Analysis:

Total Nitrogen (N)	15%
Available Phosphate (P ₂ O ₅)	5%
Soluble Potash (K ₂ O)	10%
Derived from: ammonium sulfate, urea, ammonium phosphate, and potassium chloride.	

ACTIVE INGREDIENTS:

Dimethylamine Salt of (+)-(R)-2-(2-methyl-4 chlorophenoxy) propionic acid....	0.29%
Dimethylamine Salt of 2, 4-dichlorophenoxyacetic acid.....	0.17%
Dimethylamine Salt of Dicamba (3, 6-dichloro-o-anisic acid)	0.07%

INERT INGREDIENT	99.47%
Total	100.00%

THIS PRODUCT CONTAINS: 4.80 lb (+)-(R)-2-(2-methyl-4 chlorophenoxy), propionic acid equivalent per ton or 0.24%. 2.80 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.14%. 1.20 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.06% Isomer Specific by AOAC Methods. TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Produced by:
Bonus Crop Fertilizer, Inc.
PO Box 1062, Greenville, Texas 75403-1062

NET WEIGHT: 40 LBS
EPA EST # 49991-TX-2

18.14 KGS

EPA REG NO. 2217-603-49991

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants, shoes plus socks and chemical-resistant gloves (made of any waterproof material). User should wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll free 1-877-800-5556 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

USE PRECAUTIONS:

Do not exceed specified dosages for any area. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

Other Use Precautions:

Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.

Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS:

This product controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 15-5-10 fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days. Do not use on dichondra. Do not overapply. Under conditions of adequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on St. Augustinegrass. Do not use this product on 'Floritam' St. Augustinegrass.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

HOW MUCH TO APPLY: For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, zoysiagrass, St. Augustinegrass, Bahiagrass and centipedegrass: Apply 3.2 to 4.0 pounds of product per 1,000square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq. ft. per application (0.24 lb 2,4-D ae, 0.42 lb MCPP-p ae and 0.10 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq. ft. (0.48 lb 2,4-D ae, 0.84 lb MCPP-p ae and 0.20 lb dicamba ae per acre). A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage. Do not exceed the application rate for any area.

Bag Size, Pounds	Application Rate, Pounds/1000 Sq. Ft.	Treated Area for This Bag Size, Square Feet
40	3.2	12,500
	3.6	11,000
	4.0	10,000

CALIBRATION INSTRUCTIONS:

- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

Spreader Settings	
Lawn Beauty	5
Jackson	7-1/2
Sears	4
Scotts	6

- A lawn will typically utilize 1 to 4 pounds of nitrogen per 1,000 square feet each growing season. When applied as directed, this product supplies (0.72) pounds of nitrogen per 1,000 square feet with each application. Use this product in conjunction with an additional fertilizer program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your local agricultural extension agent for the proper amount of nitrogen to be applied in your area.
- Information regarding the contents and levels of metal in this product is available on the Internet at: <http://www.aapfco.org/metals.htm>

BROADLEAF WEEDS			
Bedstraw	Curly dock	Pepperweed	Spiny Cocklebur
Black medic	Dandelion	Pigweed	Spurweed
Buckhorn	Ground ivy	Plantain	Thistle
Bullnettle	Healall	Poison ivy	Virginia creeper
Burdock	Henbit	Poison oak	Wild carrot
Chickweed, common	Knotweed	Purslane, common	Wild garlic
Chicory	Lambsquarters	Ragweed	Wild lettuce
Clover, red	Lespedeza	Sheep sorrel	Wild onion
Clover, sweet	Mallow	Shepherdspurse	Yarrow
	Morning glory	Speedwell	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return the product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

GARDENLINE® is a registered trademark of Bonus Crop Fertilizer, Inc.

EPA Reg. No. 2217-603-49991

0735129

Account	EPA Company N	EPA Product N	EPA Distributor N	Product Name	Account Expiration Date
0795288	053883	00172	049991	BCF TURF FERTILIZER WITH PRODIAMINE 0.37% (6-3-0/FE)	
0797002	008378	00068	049991	BCF TURF FERTILIZER WITH PENDIMETHALIN 1.07% (16-0-16) 50% XCU/3FE	
0756072	053883	00180	049991	BCF TURF FERTILIZER PLUS OXADIAZON 1.0% (20-0-6)/UFLEXX	03-31-2020
0782452	053883	00172	049991	BCF TURF FERTILIZER WITH PRODIAMINE 0.37% (20-3-10)	03-31-2020
0797000	053883	00170	049991	BCF TURF FERTILIZER WITH PRODIAMINE 0.29% (5-0-20)	03-31-2020
0756053	062719	00588	049991	BCF TURF FERTILIZER WITH LOCKUP PLUS (16-0-5)/UFLEXX	03-31-2020
0734226	008378	00067	049991	BCF TURF FERTILIZER WITH PENDIMETHALIN 0.86% (20-0-10)	03-31-2020
0737031	053883	00172	049991	BCF TURF FERTILIZER WITH PRODIAMINE 0.37% (14-14-14 LANDSCAPE)	03-31-2020
0702019	070506	00052	049991	TURF FERTILIZER WITH SURFLAN 0.75% (15-5-10)	03-31-2020
0735271	062719	00485	049991	BCF TURF FERTILIZER WITH DIMENSION 0.17% FG (19-0-5)/30% UFLEXX	03-31-2020
0717292	060063	00028	049991	TURF FERTILIZER WITH PRODIAMINE 20/75 (.43%)(0-0-7)	03-31-2020
0717291	062719	00494	049991	BCF TURF FERTILIZER WITH DIMENSION 0.22% FG (25-0-10)	03-31-2020
0736522	000100	01456	049991	BCF TURF FERTILIZER PLUS 0.29% BARRICADE HERBICIDE (15-5-10)	03-31-2020
0732243	000100	01456	049991	TURF FERTILIZER PLUS 29% BARRICADE HERBICIDE (0-0-7)	03-31-2020
0734272	008378	00067	049991	BCF TURF FERTILIZER WITH PENDIMETHALIN 0.86% (22-0-0)	03-31-2020
0703778	060063	00028	049991	BCF TURF FERTILIZER WITH PRODIAMINE 20/75 (.29%) 16-0-16 (UFLEXX AND IRON)	03-31-2020
0736481	000100	01340	049991	BCF TURF FERTILIZER PLUS 0.43% BARRICADE HERBICIDE WITH 30% SCU (19-0-7)	03-31-2020
0734997	002217	00603	049991	BCF WEED AND FEED WITH TRIMEC HERBICIDE .53% (16-3-4)	03-31-2020
0703127	000100	01339	049991	TURF FERTILIZER PLUS 0.38% BARRICADE HERBICIDE (0-0-7)	03-31-2020
0735129	002217	00603	049991	GARDENLINE WEED AND FEED WITH TRIMEC HERBICIDE .53% (15-5-10)	03-31-2020
0735304	000100	01340	049991	BCF TURF FERTILIZER PLUS 0.43% BARRICADE HERBICIDE (18-0-18)	03-31-2020
0734233	008378	00067	049991	BCF PROFESSIONAL WEED AND FEED WITH PENDIMETHALIN 0.86% WITH 50% SCU (18-0-18)	03-31-2020
0745536	053883	00246	049991	BCF TURF FERTILIZER WITH PRODIAMINE 0.29% (5-0-20)	03-31-2020
0730637	009404	00082	049991	WEED & FEED WITH 1.15% ATRAZINE FOR ST. AUGUSTINE LAWNS WITH 30% XCU/ 1 FE (25-0-5)	03-31-2020
0735804	053883	00170	049991	TURF FERTILIZER WITH PRODIAMINE 0.29% (19-0-5)	03-31-2020
0734662	053883	00362	049991	TURF FERTILIZER WITH 40% XCU/0.2% IMIDACLOPRID (22-0-6)	03-31-2020
0690261	062719	00494	049991	TURF FERTILIZER WITH DIMENSION 0.22% FG (19-0-10)	03-31-2020
0725166	008378	00067	049991	BCF TURF FERTILIZER WITH PENDIMETHALIN 0.86% (15-5-10)	03-31-2020
0679683	060063	00028	049991	TURF FERTILIZER WITH PRODIAMINE 20/75 (.37%)(0-0-7)	03-31-2020
0682304	060063	00028	049991	TURF FERTILIZER WITH PRODIAMINE 20/75 (.43%)(19-0-7)	03-31-2020
0643706	060063	00028	049991	TURF FERTILIZER WITH BARRICADE 20/75 (22-0-6)	03-31-2020
0573524	062719	00494	049991	BCF TURF FERTILIZER WITH DIMENSION 0.22% FG (0-0-7)	03-31-2020
0597753	000228	00435	049991	BIENTHRIN 0.069% PLUS FERTILIZER (19-0-6)	03-31-2020
0607841	060063	00028	049991	TURF FERTILIZER WITH BARRICADE 20/75 19-0-5	03-31-2020
0603846	008378	00067	049991	BCF FERTILIZER WITH PENDIMETHALIN 0.86% (0-0-7)	03-31-2020
0596687	008378	00067	049991	BCF FERTILIZER WITH PENDIMETHALIN 0.86% (18-0-18)	03-31-2020
0590177	009404	00082	049991	WEED & FEED WITH 1.15% ATRAZINE FOR ST AUGUSTINE LAWNS (25-0-5)	03-31-2020
0529991	062719	00494	049991	TURF FERTILIZER WITH DIMENSION .22% FG 16-8-8-40% XCU	03-31-2020
0691240	009404	00082	049991	WEED & FEED WITH 1.15% ATRAZINE FOR ST. AUGUSTINE LAWNS 0-0-7	03-31-2018