WEED BEATER PLUS®

BONIDE

Crabgrass & Broadleaf Weed Killer

Ready to Use

Kills Crabgrass **PLUS** broadleaf weeds

Kills the weeds not the lawn

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	. 0.313%
Quinclorac	. 0.104%
Dicamba, dimethylamine salt	. 0.029%
OTHER INGREDIENTS:	
TOTAL:	

EPA Est. No. 4-NY-1 EPA Reg. No. 2217-906-4

Keep Out Of Reach Of Children CAUTION (See attached booklet for complete Precautionary Statement and Directions for Use, including Storage

Kills to the root

Ready to Use

- √ Controls over 200 Broadleaf weeds as listed
- √ Starts working on contact
- Controls: crabgrass, dandelion, black medic, chickweed, clover, creeping charlie, ground ivy, lambsquarters, morningglory, oxalis, poison ivy, poison oak, speedwell, spurge, wild carrot, yarrow (See inside booklet for a complete list of weeds controlled)

THIS PRODUCT CONTAINS:

- 2,4-dichlorophenoxyacetic acid equivalent, 0.260%
- 3,7-dichloro-8-quinolinecarboxylic acid, 0.104%
- 3,6-dichloro-o-anisic acid equivalent, 0.024%
- Isomer Specific by AOAC Methods.

EDA E-t N- 4 NV 4

EPA Est. No. 4-NY-1 EPA Reg. No. 2217-906-4

©Bonide Products, LLC All Rights Reserved

Distributed by: Bonide Products, LLC 6301 Sutliff Road, Oriskany, NY 13424 R۱



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

- Po not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Apply when daytime temperatures are between 50°and 90°F (do not apply above 85°F for bermudagrass).
 Do not treat when air temperatures exceed 90°F as damage to turfgrass may occur.
- Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

- This product is for spot treatment only.
 Do not use clippings from treated area as common and treatment only.
- Do not use clippings from treated area as compost or mulch in vegetable gardens, flower beds or around other desirable plants.

PRODUCT FACTS:

What It Does: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed, clover as well as crabgrass and foxtail. (See inside for complete list) Where to Use: Lawns: Kills the weeds not the lawn when used as directed.

When to Use: Apply when daytime temperatures are between 50° and 90°F, (do not apply above 85°F for bermudagrass).

SHAKE WELL BEFORE USING.

HOW IT WORKS

This product contains three proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns listed on this label will not be harmed when used as directed.

O USE

This product is a ready-to-use product intended for the treatment of individual weeds, patches of weeds or sparse weeds scattered throughout home lawns. This product can be applied to the following established turfgrass species:

Cool season turfgrass	Warm season turfgrass
Kentucky bluegrass	Bermudagrass, common and hybrid
Ryegrass, perennial and annual	Zoysiagrass
Tall fescue	Buffalograss
Mixtures of cool season grasses containing fine fescues (red, hard and chewings) and bentgrass	

DO NOT apply this product to the following:

Warm season turfgrass	Other areas
St. Augustinegrass	Lawns with desirable clovers or legumes
	Dichondra
Centipedegrass	Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs)

Warm season turfgrass	Other areas
Carpetgrass	Turfgrass species that are not listed on the label.
	Gardens including vegetables, fruit tress, vines and berries.

WEEDS CONTROLLED GRASS WEEDS:

crabgrass (large and smooth) foxtail (green, yellow and giant) kikuyugrass and broadleaf signalgrass

BROADLEAF WEEDS

		DIVOADELA	IVLLDO	
	Annual yellow sweetclover Aster	Cocklebur Common mullein	Narrowleaf plantain Narrowleaf vetch	Spiny cocklebur Spiny sowthistle
	Austrian Fieldcress	Creeping Jenny	Nettle	Spotted catsear
	Bedstraw	(Creeping Charlie)	Orange hawkweed	Spotted spurge
	Beggarticks.	Cudweed	Oriental cocklebur	Spurweed
	Betony, Florida	Curly dock	Oxalis	Stinging nettle
	Bindweed, field	Daisy, English	Parsley-piert	Strawberry, India mock
	Bird vetch	Daisy, fleabane	Parsnip	Tall nettle
	Bitter wintercress Bittercress, hairy	Daisy oxeye Dandelion	Pearlwort	Tall vervain
	Bitterweed	Dichondra	Pennycress Pennywort	Tansy ragwort Tansy mustard
	Black-eyed Susan	Dogbane	Peppergrass	Tanweed
	Black medic	Dogfennel	Pepperweed	Thistle
	Black mustard	Dollarweed	Piaweed	Trailing crownvetch
	Blackseed plantain	False dandelion		Tumble mustard
	Blessed thistle	Falseflax	Plains coreopsis	Tumble pigweed
	Bloodflower milkweed	False sunflower	(tickseed)	Velvetleaf
ı	Blue lettuce	Fiddleneck	Plantain	Venice mallow

Blue vervain	Florida pusley	Poison ivy	Virginia buttonweed
Bracted plantain	Frenchweed	Poison oák	Virginia creeper
Brassbuttons	Galinsoga	Pokeweed	Virginia pepperweed
Bristly oxtongue	Goathead	Poorjoe	Wavyleaf bullthistle
Broadleaf dock	Goldenrod	Prairie sunflower	Western clematis
Broadleaf plantain	Ground ivv	Prickly lettuce	Western salsify
Broomweed	Gumweed	Prickly sida	White mustard
Buckhom	Hairy fleabane	Prostrate knotweed	Wild aster
Buckhom plantain	Hawkweed	Prostrate pigweed	Wild buckwheat
Bulbous buttercup	Healall	Prostrate spurge	Wild carrot
Bull thistle	Heartleaf drymary	Prostrate vervăin	Wild four-o'clock
Bullnettle	Heathaster '	Puncturevine	Wild garlic
Burclover	Hedge bindweed	Purslane, common	Wild garlic Wild geranium Wild lettuce
Burdock	Hedge mustard	Ragweed	Wild lettuce
Burning nettle	Hemp	Red sorrel	Wild marigold
Burweed	Henbit	Redroot pigweed	Wild mustard
Buttercup	Hoary cress	Redstem filaree	Wild onion
Buttonweed	Hoary plantain	Rough cinquefoil	Wild parsnip
Canada thistle	Hoarý vervain	Rough fleabane	Wild radish
Carolina geranium	Horsénettle	Russian pigweed	Wild rape
Carpetweed	Jimsonweed	Russian thistle	Wild strawberry
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild sweet potato
Catnip	Knotweed	Scotch thistle	Wild vetch
Catsear	Kochia	Sheep sorrel	Wild violet
Chickweed, common	Lambsquarters	Shepherdspurse	Woodsorrel
Chickweed, mouseear	Lespedeza	Slender plantain	Woolly croton
Chicory	Mallow	Smallflower galinsoga	Woolly morningglory
Cinquéfoil	Matchweed	Smooth dock	Woolly plantain
Clover, crimson	Mexicanweed	Smooth Pigweed	Wormseed
Clover, hop	Milk vetch	Sorrel	Yarrow
Clover, red	Morningglory	Sowthistle	Yellow rocket
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellowflower
Clover, sweet	Mugwort	Speedwell	pepperweed
Clover, white	Musk thistle	Spiny amaranth	popposition.
Cockle	Mustard	- - - - - - - - - -	

Doicon isas

Virginia buttonwood

Elorida puelov

Pluo vongin

WHEN TO USE

Apply this product when weeds are small and actively growing.

CRABGRASS (large and smooth):

• **Spring:** Apply this product from crabgrass emergence until the crabgrass is 3-4 inches in height. This period begins the first time that you can see crabgrass in your lawn. Re-treat at 21 days later for dense populations.

FOXTAIL (GREEN, GIANT AND YELLOW):

• **Spring and early summer:** Apply this product to foxtail during the early growth stages or when foxtail is 1-4 inches in height. Re-treat at 21 days later for dense populations.

BROADLEAF WEEDS:

- Spring: Apply when dandelion, plantain, buttercup, speedwell, knotweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.
- Fall: Apply when henbit, chickweed, white clover,

ground ivy, lawn burweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

HOW TO USE - TRIGGER SPRAYERS

- 1. Shake well.
- Spray the weeds only. Spray to lightly cover broadleaf and grass weeds.

Excessive spraying can injure some lawns. Do not apply when daytime temperatures are above 90°F as lawn injury can occur (do not apply above 85°F for bermudagrass).

USE PRECAUTIONS

For newly seeded areas:

 Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass, fine fescue mixtures and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.

For newly sodded sprigged, or plugged areas:

• The application of this product to newly sodded,

sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Seeding:

• Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the application of this product.

Irrigation:

- Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application.
- Avoid applications of this product when turfgrasses are under stress since injury may result.

Mowing:

 Delay mowing 2 days before and until 2 days after the application of this product.

STORAGE AND DISPOSAL PESTICIDE STORAGE: Keep from freezing. Store in original

container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your

local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.		
	FIRST AID	
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 	
If	Call a poison control center or doctor immediately	
arrellarred.	for trootmant advisa	

for treatment advice. swallowed: Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment advice.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash clothing before reuse. Wear long-sleeved shirt, long pants, shoes and socks.

ENVIRONMENTAL HAZARDS This pesticide is toxic to fish and aquatic invertebrates. Drift and

runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. This product has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

IMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES. INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR

PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL. CONSEQUENTIAL. OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.