

# Specimen Label

2,4-D CHOLINE SALT	GROUP	4	HERBICIDE
FLUROXYPYR-MEPTYL	GROUP	4	HERBICIDE
HALAUXIFEN-METHYL	GROUP	4	HERBICIDE



**GameOn™**  
with Arylex™ active

**SPECIALTY HERBICIDE**

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

**For postemergence control of annual and perennial broadleaf weeds in established turfgrass and ornamental grasses in golf courses (including fairways, aprons and roughs, tee boxes), industrial sites, cemeteries, commercial sod farms, and unimproved turfgrass areas. Unimproved turfgrass areas are defined as sites that are not regularly maintained, may only be mowed and treated 1-3 times per year.**

### Not for residential use

#### Active Ingredient:

2,4-D Choline salt:	
ethanaminium, 2-hydroxy-N,N,N-trimethyl-, 2-(2,4-dichlorophenoxy)acetic acid	32.00%
Fluroxypyr meptyl:	
[(4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy] Acetic acid, 1-methylheptyl ester	4.30%
halauxifen-methyl: 2-pyridinecarboxylic acid, 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-, methyl ester	0.21%
Other Ingredients	63.49%
Total	100.0%

Active Ingredient: Contains 2.87 lb of 2,4-D Choline (ethanaminium, 2-hydroxy-N,N,N-trimethyl-, 2-(2,4-dichlorophenoxy)acetic acid) per gallon. Acid Equivalent for 2,4-D Choline is 1.95 lb per gallon.

Active Ingredient: Contains 0.39 lb of fluroxypyr meptyl [(4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy] Acetic acid, 1-methylheptyl ester per gallon.

Acid Equivalent for fluroxypyr meptyl is 0.27 pounds per gallon.

Active Ingredient: Contains 0.02 lb of halauxifen (2-pyridinecarboxylic acid, 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)) per gallon. Acid Equivalent for halauxifen-methyl acid is 0.02 pounds per gallon.

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

EPA Reg. No. 62719-724

### Keep Out of Reach of Children

## WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove**

**and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.**

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of Barrier Laminate, ≥14 mils Butyl Rubber, ≥ 14 mils Nitrile Rubber or ≥ 14 mils Viton.
- Shoes plus socks
- Protective eyewear (goggles, face shield, or safety glasses)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

#### Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

### Environmental Hazards

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This product is toxic to fish and aquatic invertebrates. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants in water adjacent to treated areas. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

**Groundwater Advisory:** This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**Surface Water Advisory:** This product has a potential for reaching surface water via runoff after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of halauxifen from runoff water. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation in turfgrass setting of greater than 1 inch per hour is expected to occur within 48 hours.

### Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), restricted-entry interval, and notification to workers (as applicable). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of Barrier Laminate, ≥14 mils Butyl Rubber, ≥ 14 mils Nitrile Rubber or ≥ 14 mils Viton.
- Shoes plus socks
- Protective eyewear (goggles, face shield, or safety glasses)

### Non-Agricultural Use Requirements

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Entry Restrictions for Non-WPS Uses:** Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry, well-ventilated place. Store in original container only. In case of leak or spill, contain material and dispose as waste

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

#### Nonrefillable containers 5 gallons or less :

**Container Handling :** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows : Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows : Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

#### Nonrefillable nonrigid containers:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

#### Refillable rigid containers larger than 5 gal :

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate

### Storage and Disposal (Cont.)

collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

#### Nonrefillable rigid containers larger than 5 gal:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

### Product Information

GameOn™ provides postemergence control of annual and perennial broadleaf weeds in established turfgrass and ornamental grasses in: golf courses (including fairways, aprons and roughs, tee boxes), industrial sites, cemeteries, commercial sod farms, and unimproved turfgrass areas. Unimproved turfgrass areas are defined as sites that are not regularly maintained, may only be mowed and treated 1-3 times per year.

Not for use in residential turf.

Susceptible weeds emerged at the time of application will be controlled. A minimum spray volume of 10 gallons or more per acre (gpa) and uniform coverage are required for optimum performance. GameOn is rainfast within two hours after application.

GameOn rapidly stops growth of susceptible weeds. Typical symptoms (discoloration) of dying weeds may be noticeable in 1-3 days after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent upon weed sensitivity, weed size, turf competition, growing conditions at and following treatment, and spray coverage.

### Use Precautions

- To minimize the potential for turfgrass injury on sensitive grasses, such as bentgrass and bermudagrass, avoid additional applications within 3 weeks of a previous application.
- For optimum results, do not mow turfgrass for 72 hours after application.
- Do not allow sprays of GameOn to contact exposed suckers or exposed roots of shallow rooted trees and shrubs or injury may occur.
- The use of this herbicide in the spring when bermudagrass is breaking dormancy may delay green up of the turfgrass.

### Use Restrictions

- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- **DO NOT use on residential turf.**
- Grass clippings from mowing within 2 weeks after application must be left onsite or discarded.
- Do not apply GameOn directly to, or otherwise permit GameOn to come into contact with desirable flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious injury may occur. Do not permit spray mists containing GameOn to drift onto desirable broadleaf plants.
- Do not directly apply to desirable legumes.
- **Chemigation:** Do not apply this product through any type of irrigation system
- Do not make more than two broadcast applications per year to the same treatment site, excluding spot applications.
- **Retreatment Interval:** 21 days

- Do not apply more than 4.0 pints of GameOn per acre per application.
- Do not apply more than 3.0 lb acid equivalent of 2,4-D per acre per year.
- Do not use on centipedegrass, St. Augustinegrass, dichondra or on lawns where desirable clovers are present.
- Use this product strictly in accordance with the drift and runoff precautions on this label in order to minimize off-site exposure.

### Herbicide Resistance Management

GameOn contains 2,4-D Choline Salt, fluroxypyr-meptyl, and halauxifen-methyl herbicides that have an auxin mode of action Group 4 herbicide based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of development weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- If using post-emergence herbicides or tank mixes, control weeds early when they are relatively small.
- Apply full rates of GameOn for the most difficult to control weed in the field at the specified time to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your local company representative, local retailer, or county extension agent.
- Contact your local company representative, crop advisor, or extension agent to find out if suspected resistant weeds to these MOAs have been found in your region. Do not assume that each listed weed is being controlled by multiple mode of action. Products with multiple active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredient in this product.
- If resistance is suspected, treat weed escapes with an herbicide having a mode of action other than Group 4 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum herbicide with other mode of action as a foundation in a weed control program, if appropriate.
- Utilize sequential applications of herbicides with alternative modes of action.
- Rotate the use of this product with non-Group 4 herbicides.
- Avoid making more than two sequential applications of GameOn and any other Group 4 herbicides within a single growing season unless mixed with an herbicide with a different mode of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields to reduce weed seed production.

### Spray Drift Management

#### Avoid Injurious Spray Drift

To minimize spray drift, apply GameOn in a total spray volume of 10 gallons or more per acre using spray equipment designed to produce large-droplet, low pressure sprays. Refer to the spray equipment manufacturer's recommendations for detailed information on nozzle types, arrangement, spacing and operating height and pressure. Operate equipment at spray pressures no greater than is necessary to produce a uniform spray pattern. Operate the spray boom no higher than is necessary to produce a uniformly overlapping pattern between spray nozzles.

#### Precautions:

- Make applications only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure crops, whether dormant or actively growing.
- When applying GameOn, use low pressure equipment capable of producing sprays of uniform droplet size with a minimum of fine spray droplets.
- Under adverse weather conditions, fine spray droplets that do not settle rapidly onto target vegetation may be carried a considerable distance from the treatment area.
- A drift control or spray thickening agent may be used with this product to improve spray deposition and minimize the potential for spray drift. If used, follow all use directions and precautions on the product label.

#### Restrictions:

- Do not apply with hollow cone-type insecticide nozzles or other nozzles that produce a fine-droplet spray.
- Apply GameOn with a nozzle class that ensures medium or coarser spray.
- Do not apply GameOn directly to, or allow spray drift to come into contact with broadleaf plants directly through foliage. Do not apply GameOn directly to, or otherwise permit GameOn to come into contact with desirable flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious injury may occur.
- Do not permit spray mists containing GameOn to drift onto desirable broadleaf plants where sensitive broadleaf plants may be planted in the same season.

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

#### Droplet Size

Use only as a medium, coarse or coarser spray (ASABE S-572 standard) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

#### Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a medium droplet spray, leave one swath unsprayed at the downwind edge of the treated area.

If the sprayer is outfitted with Lesco Chemlawn Gun, Hypro Ultra-Low-Drift (ULD), TeeJet Turbo FloodJet (TF-VS), or TeeJet Air Induction (AI) nozzles no downwind buffer is required.

#### Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

#### Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include cotton, okra, flowers, grapes, fruit trees, soybeans, ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that may not be visible may injure susceptible broadleaf plants.

#### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

#### Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.



## Mixing Directions

Prior to using GameOn, invert the GameOn container several times to ensure product uniformity. Vigorous container shaking shall result in excessive foam to occur.

### GameOn – Alone

1. Fill the tank with 1/2 – 3/4 of the total amount of water.
2. Start agitation.
3. Add the required amount of GameOn.
4. Continue agitation while filling the spray tank to the required volume.
5. To ensure a uniform spray mixture, continuous agitation is required during application. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying. Apply mixture immediately after it is prepared.

**Note:** Adding oil, wetting agent, or other surfactant to the spray mixture may increase effectiveness on weeds, but also may reduce selectivity to turfgrass resulting in turf damage.

### GameOn – Tank Mix

If a broader spectrum of weed control is needed, GameOn may be tank mixed with labeled rates of other pesticides provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product.

### Tank Mixing Precautions:

- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### Tank Mixing Restrictions:

- Do not exceed specified application rates for respective products or maximum allowable application rates for any active ingredient in the tank mix.
- Do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment have been adequately cleaned. (See Equipment Clean-Out Procedures.)
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.

**Tank Mix Compatibility Testing:** When tank mixing GameOn with other materials, a jar test using relative proportions of the tank mix ingredients should be conducted prior to mixing ingredients in the spray tank. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Vigorous, continuous agitation during mixing, filling and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture.

### Mixing Order for Tank Mixes:

1. Fill the spray tank to at least 1/2 of the total spray volume required with water.
2. Start agitation and maintain throughout mixing.
3. Add different formulation types in the following order while allowing each product to completely mix and disperse before adding another:
  - a. Water Soluble Packets
  - b. Dry flowables
  - c. Wettable Powders
  - d. Liquid Suspensions
  - e. Emulsifiable Concentrates
  - f. Solutions
  - g. Adjuvants
  - h. Micronutrients
4. Finish filling the spray tank. Maintain continuous agitation during mixing and throughout application. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying. Apply mixture immediately after it is prepared.

If application or agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is

particularly useful for this purpose. Settled material may be more difficult to resuspend than when originally mixed.

### Restriction:

- Do not allow tank mixes to set overnight, if settled material cannot be resuspended with the spray tank agitation system before applications.

### Application in Liquid Fertilizer Solutions

GameOn may be applied in spray solutions containing nitrogen fertilizer. Run a tank mix compatibility test before mixing GameOn in fertilizer solution. Additional adjuvants are not needed when using GameOn in tank mix liquid nitrogen fertilizer solutions. Temporary turfgrass injury may result when liquid nitrogen fertilizer is used as the spray carrier. Foliar applied liquid nitrogen fertilizer may cause foliar leaf burn, yellowing or reduced growth due to the activity of the liquid fertilizer on the turfgrass.

### Clean-Out Procedures for Spray Equipment

1. Drain any remaining spray mixture from the application equipment, then wash out tank, boom, and hoses with clean water. Drain again.
2. Hose down the interior surfaces of the tank while filling the tank 1/2 full of water.
3. Add commercial tank cleaner, such as household ammonia, at a rate of 1 gallon per 100 gallons of water. Recirculate for 10 – 20 minutes and spray out the mixture through the boom.
4. Remove all spray nozzles and screens and clean separately.
5. If spray equipment will be used for pesticide application to crops sensitive to GameOn, repeat steps 1 through 3. Additional steps may also be required to remove all traces of GameOn including replacing hoses or other fittings that may contain adsorbed actives.
6. Thoroughly clean exterior surfaces of spray equipment.

**Note:** Rinsate may be disposed of on site according to label use directions or at an approved waste disposal facility. Reduced results may occur if water containing soil is used, such as visibly muddy water or water from ponds and ditches that is not clear.

## Directions for Use

GameOn is recommended for use on the following turfgrass species:

### Established Cool Season Turfgrass

Common Name	Scientific Name
bentgrass <sup>1</sup>	<i>Agrostis species</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewing	<i>Festuca rubra var. commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheeps	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
fescue, hard	<i>Festuca trachyphylla</i>
ryegrass, perennial	<i>Lolium perenne</i>

### Established Warm Season Turfgrass

Common Name	Scientific Name
bermudagrass <sup>1</sup>	<i>Cynodon dactylon</i>
fescue, tall (growing in warm season areas)	<i>Festuca arundinaceae</i>
zoysiagrass	<i>Zoysia japonica</i>

<sup>1</sup>Do not treat to bentgrass or bermudagrass with GameOn when the mowing height is less than 1/2 inch. Hybrid bermudagrass can be more sensitive to injury than common bermudagrass. Do not apply more than 3 pints of GameOn per acre to these grasses unless temporary turfgrass injury can be tolerated. To minimize turfgrass injury, additional applications should be made at least four weeks apart. Avoid swath overlaps. The use of this herbicide in the spring when warm season turfgrass is breaking dormancy may delay green up of the turfgrass.

GameOn may discolor and/or stunt turfgrass that is not well established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply GameOn only to healthy, well-established turfgrass that has a well-anchored root system.

### Treatment on Turfgrass Species Not Listed on the Label for GameOn

Users who wish to use GameOn on turfgrass species not listed on this label may determine the suitability for use by treating a small area at a specified rate. Prior to treatment of larger areas, observe the treated area for any sign of herbicidal injury during 40 days of normal growing conditions to determine if the target turfgrass species is tolerant. The user assumes responsibility for any plant damage or other liability resulting from use of GameOn on turfgrass species not listed on this label.

## Weeds Controlled

artichoke, Jerusalem <sup>1</sup>	flixweed	pepperweed <sup>1</sup>
aster, many-flower	galinsoga	pigweeds <sup>1</sup>
Austrian fieldcress <sup>1</sup>	garlic, wild <sup>1</sup>	plantain (such as broadleaf, narrowleaf and buckhorn)
bedstraw	geranium, Carolina	poorjoe
beggarticks <sup>1</sup>	goldenrod	prickly sida
bindweed (hedge, field and European) <sup>1</sup>	ground ivy	primrose, common
bittercress, smallflowered	hawkweed, orange	prostrate knotweed
bitterweed	healall	purslane, common
black medic	hemp, wild	pusley, Florida
broadleaf plantain	henbit	radish, wild
broomweed	horseweed (marestail)	ragweed, common
buckhorn plantain	ironweed, western	ragweed, giant
bullnettle <sup>1</sup>	ivy, ground	ragwort, tansy <sup>1</sup>
burdock, common	jewelweed	rape, wild
buttercup, smallflowered <sup>1</sup>	jimsonweed	rocket, yellow
carpetweed	knotweed <sup>1</sup>	salsify, common <sup>1</sup>
carrot, wild	kochia	salsify, western <sup>1</sup>
catnip	lambsquarters, common	shepherdspurse
Chickweed, mouse-ear	lettuce, prickly <sup>1</sup>	sicklepod
Chickweed, common	lettuce, wild	smartweed (annual species) <sup>1</sup>
chicory	loco, bigbend	sneezeweed, bitter
cinquefoil, common	lupines	sowthistle, annual
cinquefoil, rough	mallow, little <sup>1</sup>	spotted spurge
clover	mallow, Venice <sup>1</sup>	sunflower
cocklebur, common	marshelder	sweetclover
coffeeweed	morningglory, annual	tansymustard
copperleaf, Virginia	morningglory, ivy	thistle, bull
cress, hoary <sup>1</sup>	morningglory, woolly	thistle, Canada <sup>1</sup>
croton, Texas	mousetail	thistle, musk <sup>1</sup>
croton, woolly	mustards (except blue mustard)	thistle, Russian (tumbleweed) <sup>1</sup>
dandelion	narrow-leaved plantain	velvetleaf
deadnettle, purple	nettles (including stinging) <sup>1</sup>	vervains <sup>1</sup>
docks <sup>1</sup>	onion, wild <sup>1</sup>	vetches
dogbanes <sup>1</sup>	oxalis	waterplantain
doveweed	parsnip, wild	white clover
eveningprimrose, cutleaf	pennycress, field	wood sorrel
fleabane	pennywort	wormwood

<sup>1</sup> May require application to small weeds, repeat application and /or use of the higher specified rates of this product

## USE SITES

### Turfgrass Excluding Turfgrass Grown for Sod Farms

#### Application Timing and Rates

Best weed control results are obtained when GameOn is applied to small, actively growing weeds. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control.

Apply GameOn to newly seeded or sprigged turfgrass only after the second mowing or after the turfgrass seedlings have tillered and developed an adequate secondary root system. Reseeding is not recommended for at least three weeks after application.

Apply GameOn to turfgrass at the rate of 3 to 4 pints per acre as a ground broadcast treatment or spot treatment. Use higher rates from 3.5 to 4 pints per acre for weeds that are more mature or difficult to control. Calibrate application equipment prior to use according to manufacturer's directions. Avoid skips or overlaps as poor weed control or plant injury may occur.

pt/acre	fl oz/acre	fl oz/1000 sq ft	Tbsp./1000 sq ft
3	48	1.1 (33 mL)	2.2
3.5	56	1.3 (36 mL)	2.6
4	64	1.5 (44 mL)	3

#### Spray Coverage

Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. Do not broadcast apply in less than 10 gallons of total spray volume per acre. For best results and to minimize spray drift, apply in a spray volume of 10 gallons or more per acre. As turfgrass canopy and weed density increase, increase spray volume to obtain optimum weed control. Use only nozzle types and spray equipment designed for herbicide application. To reduce spray drift, follow precautions under Spray Drift Management.

#### Spot Treatment of Ornamental Turfgrass Using Portable Sprayers

Mix 1.3 fl oz of GameOn in enough water to make 1 gallon of spray and apply at any time broadleaf weeds are susceptible by wetting foliage of undesirable plants to point of runoff. This is enough spray to treat approximately 1000 sq ft of turf.

#### Restrictions:

- **DO NOT use on residential turf.**
- For optimum results, do not mow turfgrass for 72 hours after application.
- DO NOT collect grass clippings for use in mulch or compost until 2 weeks after application.
- **Retreatment Interval:** Do not reapply within 3 weeks of a previous application.
- Do not reseed for 3 weeks after application.
- Do not use GameOn on golf course putting greens
- Do not make more than two broadcast applications per year to the same treatment site, excluding spot applications.

## Turfgrass Grown for Sod Farms

**Agricultural Use Requirements:** When used in sod farms, follow PPE and re-entry instructions in the Agricultural Use Requirements section of this label.

Application Timing	GameOn (pint/acre)	Use Directions
sod farms (postemergence)	3 - 4	Apply 3 to 4 pints of GameOn per acre to control broadleaf weeds in turfgrass listed above except as noted in the Restrictions of this section. Deep-rooted perennials such as bindweed and Canada thistle may require repeat applications. Avoid mowing sod farms for 1 to 2 days before or after application. Delay irrigation until the day following application.

### Precautions:

- **Re seeding:** Delay reseeding at least 28 days following application. Preferably, with spring application, reseed in the fall and, with fall application, reseed in the spring.

### Restrictions:

- **Retreatment Interval:** Do not reapply to a treated area within 28 days of a previous application.
- Do not use on creeping grasses such as bentgrass except as a spot treatment.
- Do not apply to bermudagrass on sod farms.
- Do not use on injury-sensitive southern turfgrass such as St. Augustinegrass, centipedegrass or bermudagrass.
- Do not use on dichondra or other herbaceous groundcovers. Legumes may be damaged or killed.
- Do not apply more than 4.0 pints of GameOn per acre per application and more than a total of 8 pints of GameOn per year

## Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent permitted by law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

## Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

## Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

## Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®™Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

Label Code: CD02-434-020

Initial Publication

LOES Number: 010-02353

EPA accepted 06/06/19