### LebanonTurf

# Country Club Blended Fertilizer

## 19-0-19 SIUM SULFATE

GUARANTEED ANALYSIS	
Total Nitrogen (N)	19%
4.9% Water Insoluble Nitrogen*	
6.1% Urea Nitrogen	
8.0% Other Water Soluble Nitrogen*	
Soluble Potash (K <sub>2</sub> 0)	19%
Sulfur (S)	7%
7.0% Combined Sulfur (S)	

**Derived From:** methylene ureas, sulfate of potash, urea, ferrous oxide, ferrous sulfate.

\*12.9% Slowly available Nitrogen from Methylene Ureas.

**GUARANTEED MINIMUM ANALYSIS** 

 Total Nitrogen (N)
 19%

 4.9% Water Insoluble Nitrogen\*
 6.1% Urea Nitrogen

 8.0% Other Water Soluble Nitrogen\*
 19%

 Soluble Potash (K20)
 19%

 Sulfur (S)
 7%

 7.0% Combined Sulfur (S)
 2.0%

 Iron (Fe)
 2.0%

 0.01% Water Soluble Iron (Fe)

**Derived From:** methylene ureas, sulfate of potash, urea, ferrous oxide, ferrous sulfate.

\*12.9% Slowly available Nitrogen from Methylene Ureas.

U.S.A

CANADA

NOTICE: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

0.01% Water Soluble Iron (Fe)

### Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.7 lbs per 1,000 sq. ft. or 196 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.6 lbs. per 1,000 sq. ft. or 100 lbs. per 3,000 sq. ft. or 100 sq. ft. o

lbs per 1,000 sq. ft. or 109 lbs. per acre.			
SUGGESTED SPREADER SETTINGS			
Spreader 0.9	9#N/1,000 sq. ft.	0.5#N/1,000 sq. ft.	
Lebanon Turf	41/2	3	
Andersons AccuPr	o K	G	
Earthway Rotary	15	13	
Gandy	26	23	
Lely	5 II	31/2 II	
Lesco (letter/nume	ric) I/17	F/13	
PennMulch HVO	K	G	
ProScape SS	K	G	
Spyker	41/2	3	
Vicon	24	21	

These settings were calibrated and field tested.
However, age and condition of spreader, speed of
operation and evenness of terrain may require slightly
different settings for desired coverage.

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

Hazard Statements:
Causes skin irritation.
May be harmful if inhaled.
Causes eve irritation.

Precautionary Statements for handling:

Wash hands and exposed skin thoroughly after handling.
Wear protective gloves/protective clothing/eye protection/face protection.
If in eyes, rinse with water and seek medical attention if symptoms persist.
Take off contaminated clothing and wash before reuse.
IF ON SKIN: Wash with plenty of soap and water.
IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell

If skin irritation occurs: Get medical advice/ attention. If eye irritation persists: Get medical advice/attention. Keep out of reach of children

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

### WARRANTY

**NOTE** To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m<sup>2</sup>, applied every 4 to 8 weeks during the local growing season.

Application Rates: To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 1.6 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N)

Vicon 24 21
These settings were calibrated and field tested.
However, age and condition of spreader, speed of
operation and evenness of terrain may require slightly
different settings for desired coverage.



F699 **22-37174** 

E/21



For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation
1600 E. Cumberland St • Lebanon, PA 17042

50 Lbs. (22.7 kg) Net Weight

800-233-0628 • (717) 273-1685

## 19.0.19 SETHER EXPORTANCE OF THE SULFATE OF THE SUL

### **GUARANTEED ANALYSIS**

Total Nitrogen (N)	19%
4.9% Water Insoluble Nitrogen*	
6.1% Urea Nitrogen	
8.0% Other Water Soluble Nitrogen*	
Soluble Potash (K <sub>2</sub> 0)	19%
Sulfur (S)	7%
7.0% Combined Sulfur (S)	
Iron (Fe)	2.0%
0.01% Water Soluble Iron (Fe)	

**Derived From:** methylene ureas, sulfate of potash, urea, ferrous oxide, ferrous sulfate.

\*12.9% Slowly available Nitrogen from Methylene Ureas.

### **GUARANTEED MINIMUM ANALYSIS**

Total Nitrogen (N)	19%
4.9% Water Insoluble Nitrogen*	
6.1% Urea Nitrogen	
8.0% Other Water Soluble Nitrogen*	
Soluble Potash (K <sub>2</sub> 0)	19%
Sulfur (S)	7%
7.0% Combined Sulfur (S)	
Iron (Fe)	2.0%
0.01% Water Soluble Iron (Fe)	

**Derived From:** methylene ureas, sulfate of potash, urea, ferrous oxide, ferrous sulfate.

\*12.9% Slowly available Nitrogen from Methylene Ureas.

U.S.A

CANADA

NOTICE: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

### **Application Rates:**

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.7 lbs per 1,000 sq. ft. or 196 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.6 lbs per 1,000 sq. ft. or 109 lbs. per acre.

SUGGESTED SPREADER SETTINGS			
Spreader 0.9#N/	1,000 sq. ft.	0.5#N/1,000 sq. ft.	
Lebanon Turf	41/2	3	
Andersons AccuPro	K	G	
Earthway Rotary	15	13	
Gandy	26	23	
Lely	5 II	31/2 II	
Lesco (letter/numeric)	I/17	F/13	
PennMulch HVO	K	G	
ProScape SS	K	G	
Spyker	41/2	3	
Vicon	24	21	

These settings were calibrated and field tested.
However, age and condition of spreader, speed of
operation and evenness of terrain may require slightly
different settings for desired coverage.

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

### WARNIN



### **Precautionary Statements for handling:**

Wash hands and exposed skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection. If in eyes, rinse with water and seek medical attention if symptoms persist. Take off contaminated clothing and wash before reuse. IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell.

If skin irritation occurs: Get medical advice/ attention. If eye irritation persists: Get medical advice/attention.

Keep out of reach of children

**Hazard Statements:** 

Causes skin irritation.

Causes eye irritation.

May be harmful if inhaled.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

### WARRANTY

**NOTE** To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m<sup>2</sup>, applied every 4 to 8 weeks during the local growing season.

### **Application Rates:**

To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 1.6 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N) Equivalent to 0.9 kg of product per 100 sq. meters.

	J - 1	
SUGGESTED SPREADER SETTINGS		
Spreader 0.43 k	g #N/100 m².	0.22 kg #N/100 m <sup>2</sup> .
Lebanon Turf	41/2	3
Andersons AccuPro	K	G
Earthway Rotary	15	13
Gandy	26	23
Lely	5 II	3½ II
Lesco (letter/numeric)	I/17	F/13
PennMulch HVO	K	G
ProScape SS	K	G
Spyker	41/2	3
Vicon	24	21

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



FAQQ

22-37174



For technical assistance or more information about our products visit www.LebanonTurf.com

### Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight