

PEEL CORNER
OF BOOKLET

Manufactured By:



230 FM 87

BONHAM, TEXAS 75148

EPA Reg. No. 7401-311 • EPA Est. No. 7401-TX-01

Visit Us At: www.fertilome.com

Product Questions? 855-270-4776

10363-0315-CL



ferti·lome[®]

Fire Blight Spray

- For Apples, Pears, Pyracantha, Chrysanthemums, Philodendron, Dieffenbachia, And Roses
- 1 Tablespoon Makes 2 1/2 Gallons Of Spray

ACTIVE INGREDIENTS:

Streptomycin Sulfate* . . .22.4%;

INERT INGREDIENTS: . . .77.6%

*Contains 17% Streptomycin derived from 22.4% Streptomycin Sulfate.

Keep Out Of Reach Of Children

CAUTION

See Back Panel For Additional
Precautionary Statements

NET CONTENTS 2 Ozs.

PRECAUTIONARY STATEMENTS • HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies and drainage systems.

FIRST AID - Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **In the event of a medical or chemical emergency, contact ChemTel North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585**



If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 – 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.• Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

1. WHAT IS THIS PACKAGE FOR?

Use this product to control *Fire Blight and Bacterial Wilt, Stem Rot, Leaf Spot, and Crown Gall* on fruit trees, flowers and shrubs.

2. HOW DO YOU USE IT?

FRUIT: APPLES AND PEARS (Areas Other Than West Coast): To control *Fire Blight* in home plantings, apply 1 tablespoon in 2½ gallons of water (equivalent to 100 ppm or 4 ozs. per 50 gallons of water) in foliar and blossom sprays. Make first spray at the start of blossoming period. Continue spray applications every 3 to 4 days during bloom time. Apply additional sprays every 5 to 7 days after blossom period when weather conditions favor spread of Fire Blight. Do not apply when fruit is visible.

PYRACANTHA: To control *Fire Blight*, apply a solution of 1 level tablespoon of **ferti•lome® FIRE BLIGHT SPRAY** per 2½ gallons of water (100 ppm concentration). Begin spraying at start of blossom period. Repeat at 3 to 4 day intervals during bloom and at 5 to 7 days intervals after bloom if weather favors disease spread.

CHRYSANTHEMUMS (Cuttings): To control *Bacterial Wilt*, soak cuttings for 4 hours in a solution containing ½ tablespoon of **ferti•lome® FIRE BLIGHT SPRAY** per 2½ gallons of water (50 ppm concentration) and plant as usual.

DIEFFENBACHIA: To control *Bacterial Stem Rot*, spray plants at 5 to 7 day intervals with a solution of 1 level tablespoon of **ferti•lome® FIRE BLIGHT SPRAY** in 2½ gallons water (100 ppm concentration) or soak cuttings for 20 minutes in a solution of 2 level tablespoons in 2½ gallons of water (200 ppm concentration) and plant in sterile rooting medium.

PHILODENDRON: To control *Bacterial Stem Rot*, remove all decayed leaves and spray with a solution of 2 level tablespoons of **ferti•lome® FIRE BLIGHT SPRAY** in 2½ gallons of water (200 ppm concentration) as preventative or at first sign of water soaked leaves. Repeat at 4 to 5 day intervals.

ROSES: To control *Crown Gall*, remove infested plant and cut out gall tissue. Soak root system and cut surfaces of infested area for 15 minutes in a solution of 2 level tablespoons of **ferti•lome® FIRE BLIGHT SPRAY** per 2½ gallons water (200 ppm concentration). Replant in soil free from disease organisms, and apply ½ tablespoon per 2½ gallons water (50 ppm concentration) as foliage spray at weekly intervals.

3. HERE ARE THE RESULTS YOU SHOULD EXPECT!

When used according to directions, this product will control *Fire Blight and Bacterial Wilt, Stem Rot, Leaf Spot, and Crown Gall* on fruit trees, flowers and shrubs.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container. Keep in area inaccessible to children and pets.

PESTICIDE DISPOSAL: If empty: Non-refillable container. Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. To the extent consistent with applicable law, Voluntary Purchasing Groups, Inc. (VPG) warrants that this product conforms to the chemical description on the label and that the limit of any liability incurred shall be the purchase price paid by the user or buyer.

Manufactured By:
VOLUNTARY PURCHASING GROUPS, INC.
230 FM 87 • BONHAM, TEXAS 75418
EPA Reg. No. 7401-311 • EPA Est. No. 7401-TX-01
Visit Us At: www.fertilome.com
Product Questions? 855-270-4776



This page left blank intentionally.

This page left blank intentionally.