

MCPA, Triclopyr, MCPP-p	GROUP	4	HERBICIDE
Carfentrazone	GROUP	14	HERBICIDE

LESCO®

 **Momentum®
XPD**

Herbicide

For selective weed control in ornamental lawns and turf grasses

ACTIVE INGREDIENTS:	% by Weight
MCPA, 2-ethylhexyl ester	28.50%
Mecoprop-p, acid	5.48%
Triclopyr, butoxyethyl ester	4.23%
Carfentrazone-ethyl	0.49%
OTHER INGREDIENTS:	61.30%
TOTAL:	100.00%

LESCO® Momentum XPD Herbicide is an Emulsifiable Concentrate which contains:
1.50 lb. MCPA acid equivalent per gallon or 18.28%
0.45 lb. Mecoprop-p acid equivalent per gallon or 5.48%
0.25 lb. Triclopyr acid equivalent per gallon or 3.04%
0.04 lb. Carfentrazone-ethyl per gallon or 0.49%

Contains petroleum distillates


For Medical Emergencies, Call 877-325-1840
For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMREC 800-424-9300

**KEEP OUT OF REACH OF CHILDREN
WARNING / AVISO**

SI USTED NO ENTIENDE LA ETIQUETA, BUSQUE A ALGUIEN PARA QUE SE LA EXPLIQUE A USTED EN DETALLE. (IF YOU DO NOT UNDERSTAND THIS LABEL, FIND SOMEONE TO EXPLAIN IT TO YOU IN DETAIL.)
SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

DISTRIBUTED BY: LESCO, Inc
24500 Center Ridge Rd. Suite 350
Westlake, OH 44145

EPA REG. NO. 228-760-10404
EPA EST. NO. indicated by the first two letters of the batch number on this package
(CH) 228-IL-001, (ST) 072344-MO-004



NET CONTENTS:
2.5 gal (9.46 L)

SiteOne.com/LESCO

10869225

FIRST AID

IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately.• Do not give any liquid to the person.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor immediately for treatment advice.
NOTE TO PHYSICIAN: Contains petroleum distillate. Vomiting may cause aspiration pneumonia.	

HOTLINE NUMBER

Have the product container or label with you when calling Poison Control 1-800-222-1222 or a doctor or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12:00 PM Pacific Standard Time.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING / AVISO

Causes substantial but temporary eye injury. Harmful if swallowed. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequent repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

All mixers, loaders, applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks,
- protective eyewear,
- chemical-resistant gloves made of barrier laminate, butyl rubber (>14 mil.), nitrile rubber (>14 mil.) or Viton (>14 mil.), and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or other exposure to the concentrate.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and/or absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROL STATEMENT:

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.607(d-e), the handler PPE may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to

treated areas. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

GROUNDWATER ADVISORY STATEMENT: This product has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

FISH ADVISORY STATEMENT: This product may be hazardous to aquatic organisms, particularly in clear, shallow water bodies that are adjacent to treated areas. Transport to water by runoff or spray drift of this product in areas where surface water is present, or intertidal areas below the mean high water mark, should be avoided. Do not contaminate water when disposing of equipment wash water or rinsate.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

This pesticide must be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposure.

Do not apply this product in a way that will contact workers, other persons, or pets, either directly or through drift. Keep people and pets out of the area during application. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical-resistant footwear plus socks
- protective eyewear
- chemical-resistant gloves of barrier laminate, butyl rubber (>14 mil.), nitrile rubber (>14 mil.) or Viton (>14 mil.), and
- chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

This product is a combination of 4 active ingredients and provides fast and effective weed control of troublesome weeds in Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Roughs and Native Areas); also for use on Sod Farms. Additionally, this product is for use in industrial areas, railroads and fence rows and other similar turf areas.

USE RESTRICTIONS

- Do not apply this product through any type of irrigation system.
- Do not apply this product aerially.
- Do not use on golf course greens, tees, or aprons.
- Do not use in or near greenhouses.
- Do not graze or hay treated areas.
- Do not pour spray solutions near desirable plants.
- Do not use on carpetgrass, St. Augustinegrass, dichondra, or on lawns where desirable clovers are present unless removal is desired.
- Do not use adjuvants, oils, wetting agents, fertilizers, surfactants or other pesticides to the spray solution unless your experience indicates that the tank mixture will not result in turf injury.
- The maximum rate per application is 4.5 pts per acre (0.84 lbs MCPA ae, 0.25 lbs Mecoprop-p, 0.14 lbs Triclopyr ae, 0.02 lbs Carfentrazone-ethyl per acre), and the maximum annual rate is 9 pints per acre per year (1.69 lbs MCPA ae, 0.5 lbs Mecoprop-p, 0.28 lbs Triclopyr ae, 0.04 lbs Carfentrazone-ethyl per acre per year).
- Do not apply more than 2 broadcast applications to the same treatment site per year, excluding spot treatments.
- The minimum retreatment interval between broadcast applications is 30 days.
- This product is persistent and may be present in treated plant materials for over 30 days after application. Do not remove grass clippings off-site for compost distribution or mulching until 30 days after application.
- Only use for sites, pests, and application methods specified on this labeling.

Endangered Species: It is a Federal offense to use any pesticide in a manner that results in the death of an endangered species. Use of this product may pose a hazard to endangered or threatened species. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county in which you are applying the product. To obtain Bulletins, no more than six months before using this product, consult <http://www.epa.gov/espp/> or call 1-800-447-3813. You must use the Bulletin valid for the month in which you will apply the product.

USE PRECAUTIONS

- Failure to observe all precautions may result in injury to turfgrass and/or susceptible plants.
- Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
- Avoid fine mists.
- Use lawn-type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees.
- Use caution when applying to ensure no spray solution collects on the tires of application equipment.
- Use caution when daily high air temperature exceeds 90 degrees.
- For optimum results, do not apply if rainfall is expected within 4 hours of the application.
- The suitability of this product on non-recommended turf species may be determined by treating a small area at 3 pints per acre.
- The treated area should be observed for any sign of turf injury for a period of 14 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.
- Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

RESISTANCE MANAGEMENT

For resistance management, this product contains Group 4 herbicides (MCPA, Mecoprop-p and Triclopyr) and a Group 14 herbicide (Carfentrazone). Any weed population may contain or develop plants naturally resistant to this herbicide and other Group 4 and 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 4 and 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and that considers mechanical control methods, cultural (e.g. timing to favor the desirable plants and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Report any incidence of non-performance or suspected resistance of this product against a particular weed to your local LESCO, Inc. representative, local retailer, or county extension agent.
- For further information or to report suspected resistance, 1-800-347-4272.

SPRAYING INSTRUCTIONS

Spray Distribution: The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator.

Spray Volumes: Apply in 10 to 240 gallons of water per acre (0.2 to 5.5 gallons of water per 1,000 square feet).

Spot Spray with Hand Sprayers: Calibration and proper uniformed applications are essential when using this product. Over application or rates above those specified, including excessive overlaps of this product, can cause injury. Calibrate sprayer to deliver 1 gallon of spray solution per 1,000 square feet of treated area. Adjust application rate accordingly if utilizing a higher or lower spray gallonage.

Lower Volume Equipment: Use as little as 10 gallons of water per acre. Use only application equipment that can spread droplets uniformly, wetting each weed surface.

MIXING AND CLEANING INSTRUCTIONS

Fill spray tank with water. Start agitation and slowly add the specified amount of concentrate. Maintain continuous agitation after mixing and during application. If this product remains in the tank for extended periods of time, some separation may occur. Reagitate before use.

TANK MIXING: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Clean application equipment carefully before and after application. Use a strong detergent or tank cleaner to rinse all parts of the sprayer thoroughly.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- User must apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to select the nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boomless Ground Applications:

- Applicators are required to select the nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Carfentrazone-ethyl is a contact herbicide. Avoid any drift conditions that would allow the product to contact desirable vegetation. Carfentrazone-ethyl is not volatile; however, mist from spray drift may cause injury to sensitive plants.

The interaction of equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses, or to applications of dry materials. Where states have more stringent regulations, they must be observed.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Spray Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Handheld Technology Applications

- Take precautions to minimize spray drift

WEEDS CONTROLLED

Artichoke*	Florida pusley*	Pennywort*	Spotted knapweed*
Aster*	Galinsoga	Pepperweed	Spurge*
Bedstraw*	Goatsbeard	Pigweed	Stinging nettle*
Bindweed*	Goldenrod*	Plantain	Thistle*
Black medic*	Ground ivy*	Poison hemlock*	Toadflax*
Broadleaf plantain*	Hawkweed*	Poison ivy*	Velvetleaf*
Buckhorn plantain*	Healall*	Poison oak*	Veronica*
Burdock	Henbit	Pokeweed*	Vetch*
Buttercup*	Horsenettle*	Poorjoe*	White clover*
Canada thistle*	Horseweed*	Prickly lettuce	Wild aster*
Carpetweed*	Jimsonweed*	Prickly sida*	Wild carrot
Catnip*	Kochia*	Primrose*	Wild garlic*
Catsear	Knawel*	Puncturevine*	Wild geranium*
Chickweed	Knotweed*	Purslane*	Wild lettuce*
Chicory	Lambsquarters	Ragweed	Wild mustard*
Cinquefoil	Lespedeza*	Red sorrel*	Wild onion*
Clover	Mallow*	Redstem filaree*	Wild radish
Cocklebur	Marshelder	Sheep sorrel*	Wild violet*
Coffeeweed	Matchweed*	Shepherd's-purse	Yarrow*
Croton	Mouse-ear chickweed	Smartweed*	Yellow rocket
Dandelion	Mustard	Sowthistle*	Yellow woodsorrel*
Dock	Oxeye daisy*	Speedwell*	
Dogfennel*	Parsley-piert*	Spiderwort*	

For complete control, some of these species may require repeat spot applications and/or use of the highest specified rate.

*Not registered for use by the state of California.

ORNAMENTAL LAWNS AND TURF AND SOD FARMS

Apply 2.75 to 4.5 pts per acre of this product to control weeds growing in Kentucky bluegrass, annual bluegrass, annual ryegrass, perennial ryegrass, tall fescue, red or fine leaf fescues, creeping bentgrass, colonial bentgrass, hybrid Bermudagrass, common Bermudagrass, centipedegrass, zoysiagrass, bahiagrass and buffalograss. See below table for details on application rates.

Generally, the lower application rates will provide satisfactory control of sensitive weeds. The higher application rates will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages. For all applications: (1) do not overlap spray patterns; and (2) use reduced rates if turfgrass is stressed from heat, drought, etc.

Note: Bronzing with spotting and speckling on the turfgrass foliage may occur following application. Treated turfgrass will recover with new growth. Discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, avoid applications to highly actively growing turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress.

Table 1. Use Rates			
TURFGRASS*	Use Rate (Fluid Ounces)	Use Rate (Pints)	Notes
Cool-season Turf			
Kentucky bluegrass	48 to 72 fl. oz. per acre	3 to 4.5 pints per acre	Use reduced rates if grass is stressed from heat, drought or lower mowing heights such as fairways. Some temporary discoloration and growth regulation may occur.
Annual bluegrass			
Annual ryegrass	1.1 to 1.7 fl. oz. per 1,000 sq. ft.		For best results apply when weeds are small in size and actively growing.
Perennial ryegrass			
Tall fescue			
Fine fescue			
Creeping bentgrass	44 to 48 fl. oz. per acre	2.75 to 3 pints per acre	Excluding use on golf greens, tees and aprons.
Colonial bentgrass	1.0 to 1.1 fl. oz. per 1,000 sq. ft.		
Warm-season Turf			
Hybrid Bermudagrass	44 to 56 fl. oz. per acre	2.75 to 3.5 pints per acre	For Centipedegrass use lower rates until tolerance to injury can be determined.
Common Bermudagrass			
Centipedegrass	1.0 to 1.3 fl. oz. per 1,000 sq. ft.		Do not apply to warm-season turf during spring green-up or in the late fall during the transition period between active growth and dormancy.
Zoysiagrass			
Bahiagrass			
Buffalograss			This product can be applied to fully dormant warm season grasses.

*This herbicide has demonstrated turfgrass tolerance. However, not all varieties have been evaluated. Turfgrass managers should treat small areas prior to widespread use.

TIMING FOR TURFGRASS APPLICATIONS

Follow-up applications may be required for dense infestations of perennial and biennial weeds.

Newly Seeded Areas: The application of this herbicide to grass seedlings is not recommended until after the second mowing. Do not apply to newly seeded areas until grass is well established unless discoloration or damage can be tolerated.

Newly Sodded, Sprigged or Plugged Areas: Do not apply to newly sodded, sprigged, or plugged grasses for at least 4 weeks after the sodding, sprigging, or plugging operations.

Seeding: Turf species listed on this label can be seeded into treated areas three weeks after the application.

Mowing: For best results, turf should not be mowed for 2 days before or after application to maximize leaf surface and herbicide uptake and translocation.

Irrigation: For best results do not irrigate or water the turf within 24 hours after application. If rainfall does not occur within 7 days of the application, irrigation of at least $\frac{1}{2}$ " is recommended.

NON-TURF AREAS (INDUSTRIAL AREAS, RAILROADS AND FENCE ROWS)

Industrial areas, railroads and fence rows: Apply 3.5 to 4.5 pints of this product per acre to control listed broadleaf weeds. Apply in 10 to 240 gallons of water per acre (0.2 to 5.5 gallons of water per 1,000 square feet).

Apply any time between the time when plants come into full leaf (spring) to when the plants begin to go dormant. Best results are obtained

when weeds are young and actively growing. Do not cut weeds until the herbicide has had sufficient time to translocate to the roots. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 1.5 fluid ounces of this product per gallon of water and spray to thoroughly wet all foliage.

Limitations:

Applications to non-turf areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production.

STORAGE AND DISPOSAL

Do not contaminate water, food, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store above 28°F or agitate before use.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

LESCO is a registered trademark of LESCO, Inc.

All other trademarks are the property of their respective owners.

10869225 V0