

LESCO[®]

TwoSome™ Fungicide

For the prevention and control of fungal diseases on ornamental plants grown in containers, fields or greenhouses, as well as fungal turf diseases on greens, fairways, tees and similar turf sites. This product may be applied to ornamentals by chemigation or ground spray equipment, and to turf sites using ground spray equipment.

For use by individuals/firms licensed or registered by the State to apply ornamental or turf pest control products. This product is NOT for residential use.

ACTIVE INGREDIENTS:

Thiophanate-methyl: Dimethyl [(1,2-phenylene)bis(iminocarbonothioyl)]bis(carbamate)*	19.65%
Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide**	19.65%
OTHER INGREDIENTS:	60.70%

TOTAL: 100.00%

*Equivalent to 1.87 pounds Thiophanate-methyl per gallon.

**Equivalent to 1.87 pounds Iprodione per gallon.



▲
PEEL HERE TO OPEN

KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN
 See inside booklet for additional PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.
For Medical Emergencies Only, Call (877) 325-1840.

FIRST AID

IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

Distributed by: **LESCO, Inc.**
 1385 East 36th Street
 Cleveland, OH 44114-4114

#10164021
NET CONTENTS: 1 gal. (3.78 L)

EPA Reg. No. 228-630-10404
 EPA Est. No. indicated by the first two letters
 of the batch number on this package
 (AL) 228-IL-002, (SA) 228-IL-004

13143000

RV052014[1]

LESCO[®]

TwoSome[™] Fungicide

For the prevention and control of fungal diseases on ornamental plants grown in containers, fields or greenhouses, as well as fungal turf diseases on greens, fairways, tees and similar turf sites. This product may be applied to ornamentals by chemigation or ground spray equipment, and to turf sites using ground spray equipment.

For use by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products. This product is NOT for residential use.

ACTIVE INGREDIENTS:

Thiophanate-methyl: Dimethyl [(1,2-phenylene)bis(iminocarbonothioyl)]bis(carbamate)*	19.65%
Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide**	19.65%
OTHER INGREDIENTS:	60.70%

TOTAL: 100.00%

*Equivalent to 1.87 pounds Thiophanate-methyl per gallon.

**Equivalent to 1.87 pounds Iprodione per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN**

See inside label booklet for **FIRST AID** and **PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.
For Medical Emergencies Only, Call (877) 325-1840.

#10164021

NET CONTENTS: 1 gal. (3.78 L)

TM

Distributed by: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114

EPA Reg. No. 228-630-10404
EPA Est. No. indicated by the first two letters
of the batch number on this package
(AL) 228-IL-002, (SA) 228-IL-004

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION/PRECAUCIÓN

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Mixers, loaders, others exposed to the concentrate and cleaners/repairers of equipment must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Chemical-resistant apron and
- Chemical-resistant footwear plus socks

Applicators using hand-held equipment must wear:

- Coveralls or long-sleeved shirt and long pants
- Chemical-resistant gloves
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposures and
- Dust-mist filtering respirator (NIOSH-approved respirator with any R, P or HE filter)

Applicators using mechanical ground equipment must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Applicators using truck-mounted equipment with a handgun at the end of the hose (i.e., for commercial turf-grass or ornamental applications) and all the other handlers not specified above must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This chemical can contaminate surface waters through aerial and ground spray applications. Under some conditions, it may also have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips and areas overlying tile drainage systems that drain to surface water.

This product is toxic to invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Cover, incorporate or clean up spills. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

If applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application. Do not apply this product when the wind direction is toward aquatic area.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours for ornamental uses. The restricted-entry interval for all other WPS uses is 24 hours.

(continued)

AGRICULTURAL USE REQUIREMENTS (cont.)

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposures

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than the applicator) or pets on the treatment area during application. Do not enter or allow others to enter the treated area until sprays have dried.

APPLICATION RESTRICTIONS

This product is for use on ornamentals by chemigation or ground spray equipment and on turf using ground spray equipment.

This product is not for residential use.

For use by certified applicators or those under their immediate supervision.

Do not apply with fixed wing or rotary aircraft.

Not for use on turf being grown for sale or other commercial use as sod. Do not apply to home orchards/backyard fruit trees after fruit set.

Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.

For golf courses only, do not apply to turf cut higher than 1" on golf holes where water bodies are present.

Do not apply this product when the wind direction is toward aquatic areas.

CHEMIGATION INSTRUCTIONS

Apply this product only through pressurized drench (flood), sprinkler (overhead), or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Requirements for Chemigation Systems Connected to Public Water Systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Chemigation System Requirements

Systems utilizing a pressurized water and pesticide system must meet the following requirements:

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Chemigation Precautions

Do not apply in greenhouses when temperatures are above 95°F.

Do not allow the spray mixture to stand in the tank for more than 12 hours.

For best results do not water treated areas within 24 hours of application.

Chemigation Mixture Preparation and Application

Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system and flush thoroughly with clean water.

See the label instructions in the "Ornamentals Section" of this label for dilution rates and timing of application. Shake product container well before pouring into spray tank.

When applying this product via chemigation, a pesticide supply tank should be used. Prepare a suspension of this product in a mix tank by filling the tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation and slowly add the specified amount of this product. Complete the mix by adding the remaining volume of water.

Apply this product with liquid fertilizers that are chemically neutral. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, tend to degrade the pesticide resulting in reduced performance, and must be avoided.

The spray mixture must be continuously agitated during application.

Apply this product continuously for the duration of the water application.

Except in cases of severe infestation, allow one week between applications of this and other control products. (This guideline can be relaxed where a severe insect or disease attack requires immediate treatment.)

Instructions for Sprinkler (Overhead) Chemigation

Observe the requirements in the "Chemigation System Requirements" section.

Follow instructions in the "Chemigation Mixture Preparation and Application" section.

Set the sprinkler system to deliver 0.1 to 0.25 inch of water per acre. Volumes of water higher than this reduce efficacy. Start sprinkler and uniformly inject the suspension of this product into the irrigation water line so as to deliver the desired rate per acre. Inject the suspension of this product with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing.

Maintain continuous agitation in the mix tank during mixing and application to assure uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting larger volumes of a more dilute suspension per unit time.

Application of more than specified quantities of irrigation water per acre tend to result in decreased product performance.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing product cannot be flushed and must be dismantled and drained. In a center pivot system, block the nozzle set nearest the well pivot injection unit to prevent spray being applied to this area.

Where sprinkler distributions do not overlap sufficiently, unacceptable disease control will result.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.

Check local restrictions and requirements regarding sprinkler irrigation application as they might vary from state to state.

NOTE: When treatment with this product has been completed, do not irrigate area for 24 to 48 hours to prevent washing the chemical off the crop.

Instructions for Pressurized Drench (Flood) Chemigation

Observe the requirements in the "Chemigation System Requirements" section.

Follow instructions in the "Chemigation Mixture Preparation and Application" section.

Systems using a gravity-flow pesticide-dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

Instructions for Drip (Trickle) Chemigation

Observe the requirements in the “Chemigation System Requirements” section.

Follow instructions in the “Chemigation Mixture Preparation and Application” section.

TURF APPLICATIONS

For use only by certified applicators or those under their immediate supervision.

This product prevents and controls the following turf diseases:

- Anthracnose (*Colletotrichum graminicola*)
- Brown Patch (*Rhizoctonia solani*)
- Dollar spot sp.
- Leaf spot (such as Helminthosporium Leaf Spot caused by *Drechslera* spp.)
- Pink snow mold

Turf Use Restrictions

- DO NOT apply this product to turf through any type of irrigation system (chemigation).
- DO NOT make more than six applications of this product in a year.
- DO NOT use less than a 14-day retreatment interval.
- DO NOT graze treated areas, and do not feed clippings to livestock.

Preparation

Use with 1 to 4 gallons of water per 1,000 ft² for tees and greens. For larger turf areas, such as Fairways, roughs, Sod farms, public do not use water volumes below 1 gal./1,000 ft².

Shake the product container well before adding to the spray mix.

While filling the spray tank with the desired amount of water, start agitation and add the specified amount of this product to the tank (see the table below for rates).

Thoroughly rinse the product container and pour the rinsate into spray tank.

Application

Do not allow the spray mixture to stand in the tank for longer than 12 hours prior to application.

Agitate the spray mixture continuously during application.

Apply with properly calibrated spray equipment using standard pressures to minimize spray drift.

For Disease Prevention:

- Apply using the Light Pressure rates when weather conditions are favorable for the development of disease.
- Repeat applications every 14 to 21 days as necessary.

For Disease Control:

- Apply using the appropriate rate based on the level of infestation (Light/Medium/Heavy) and repeat applications every 14 days until the disease is brought under control. Once the disease is controlled, follow instructions for disease prevention above.

For best results do not water or mow treated areas within 24 hours of application.

Turf Application Rates

Use Site	Disease Pressure	Single Application Rate (fl. oz. product/1,000 sq. ft.?)	Maximum Annual Application Rate (fl. oz. product/1,000 sq. ft.?)
Non-Residential Turf	Light	1.0 to 2.0	16.0
	Medium	2.0 to 3.0	
	Heavy	3.0 to 4.0	
Golf Courses – Tees/Greens/Aprons	Light	1.0 to 2.0	24.0
	Medium	2.0 to 3.0	
	Heavy	3.0 to 4.0	
Golf Courses – Fairways (except Florida)	Light	1.0 to 2.0	8.0
	Medium	2.0 to 3.0	
	Heavy	3.0 to 4.0	
Golf Courses – Fairways (Florida only and only during overseeding)	Light	1.0 to 2.0	4.0
	Medium	2.0 to 3.0	
	Heavy	3.0 to 4.0	

ORNAMENTAL APPLICATIONS

Do not apply to home orchards/fruit trees after fruit set.

This product will control fungal diseases on the following ornamentals:

Ageratum	Columbine	Episole	Lilies	Poppy
Ajuga	Conifers	Euronymous	Lipstickvine (Aeschynanthus)	Pothos ¹
Almond – Ornamental	Coral bells (Heuchera)	Ficus	Marigold	Primrose
Alyssum	Crape myrtle	Forsythia	Monarda (Bee Balm)	Privet
Andromeda	Crassula	Gazania	Palm	Protea
Aphelandra	Croton	Geranium	Pansy	Pyracantha
Artemesia	Cyclamen ²	Gladiolus	Peach – Ornamental	Rhododendron
Aster	Daffodils	Gloxinia	Peperomia	Rose
Azalea	Dahlia	Gypsophila	Periwinkle	Rose Tree of China
Boxwood	Delphinium	Hawthorn	Petunia ¹	Salvia
Cactus	Deutzia	Holly	Philodendron	Schefflera
Calendula	Dianthus	Hoya	Phlox	Snapdragon
Carnation	Dieffenbachia	Hydrangea	Pilea	Stative
Cherry – Ornamental	Dizygotheca	Impatiens ¹	Pine	Tree Ivy
Chrysanthemum ³	Dogwood	Iris	Pittosporum	Viburnum
Cineraria	Dracena	Juniper	Plum – Ornamental	Violet
Coleus	English ivy	Kalanchoe	Poinsettia ²	Zinnia

¹Do not use this product as a drench on impatiens, petunias or pothos.

²Residue will be noticeable on foliage if plants are treated prior to sale when in bloom.

³On chrysanthemums, do not make repeat applications at high drench rates.

NOTE: DO NOT use this product on *Spathiphyllum* or New Guinea Impatiens.

Apply this product by ground spray or chemigation only to control the following diseases:

Foliar Applications

Aerial web blight	Botrytis blight	Didymellina leaf spot	Ink spot	Ray blight
Alternaria leaf blight	Botrytis gray mold	Diplodia tip blight	Ovulinia	Rhizoctonia blight
Anthraconose	Cercospora leaf spot	Entomosporium leaf spot	Phomopsis blight	Scab
Ascochyta blight	Corynespora leaf spot	Fusarium leaf spot	Powdery mildew	Septoria leaf spot
Black Spot ¹	Cylindrocladium ²	Helminthosporium leaf spot	Ramularia leaf spot	

¹Roses only

²Not effective on *Cylindrocladium spathiphylli*

Drench Applications (after transplant)

Alternaria leaf blight	Cylindrocladium rot	Fusarium root rot	Ink spot	Sclerotinia stem, crown and root rot
Botrytis blight	Fusarium crown rot	Fusarium stem rot	Rhizoctonia root rot	Thielaviopsis rot

Preparation

Mix 33 to 84 fl. oz. of this product with 100 gallons of water. Use the lower rates listed for disease prevention or when pest pressure is low. When pest pressures are moderate to high apply using the higher range of the rates listed.

Application

Foliar spray:

When conditions are favorable for disease development, apply to foliage up to the point of runoff. Make repeat applications every 7 to 14 days as necessary for control.

DO NOT apply more than 192 fl. oz. of this product per acre in a single application. Do not make more than 4 applications per year.

Drench:

Mix 13.5 fl. oz. per 100 gallons of drench solution. After transplant into propagation beds or containers, apply 0.5 to 2 pints of drench solution per square foot depending on media type and depth. To maintain control, make repeat applications every 2 weeks as necessary while conditions remain favorable for disease development. This product should be used as part of a managed rotational program.

Do not apply more than 37 fl. oz./1,000 ft² per year.

Do not make more than 6 applications per year.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, clean, dry place. Do not store at temperatures exceeding 120°F or at temperatures below 32°F.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, INC., BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

RV052014[1]

LESCO is a registered trademark of LESCO Technologies, LLC.
TwoSome and the sweeping design are trademarks of LESCO Technologies, LLC.
All other trademarks are the property of their respective owners.