

LESCO®
MOMENTUM®
HERBICIDE

**FOR SELECTIVE BROADLEAF AND CERTAIN
GRASSY WEED CONTROL**

Contains 2,4-D, Quinclorac and Dicamba

**Controls: Dandelion, Chickweed, Black Medic,
Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle,
Large and Smooth Crabgrass and many other
species of broadleaf and certain grassy weeds;
some of which are listed on this label.**

ACTIVE INGREDIENTS:

Dimethylamine Salt of
2,4-Dichlorophenoxyacetic Acid* 13.24%
Quinclorac (3,7-Dichloro-8-
quinolinecarboxylic Acid)** 8.25%
Dicamba (3,6-Dichloro-o-Anisic Acid)*** 1.38%

OTHER INGREDIENTS: 77.13%
TOTAL: 100.00%

By Isomer Specific AOAC Method, Equivalent to:

* 2,4-Dichlorophenoxyacetic Acid
..... 11.34%, 1.00 lbs./gal.

** 3,7-Dichloro-8-quinolinecarboxylic Acid.....
..... 8.25%, 0.75 lbs./gal.

*** 3,6-Dichloro-o-Anisic Acid.....
..... 1.38%, 0.125 lbs./gal.

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

**SEE INSIDE BOOKLET FOR FIRST AID
AND PRECAUTIONARY STATEMENTS**

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840**

#091598NR
NET CONTENTS: 64 oz (1.89 L)

TM

EPA Reg. No. 228-531

EPA Est. No. 228-IL-1

Manufactured By: **Nufarm Americas Inc.** • 11901 S. Austin Avenue • Alsip, IL 60803

LESCO is a registered trademark of LESCO Technologies, LLC.
Momentum is a registered trademark and used by LESCO, Inc. with permission of the trademark owner.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material.

All mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks, and
- Chemical-resistant gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

The chemicals in this product have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls worn over short-sleeve shirt and short pants, chemical-resistant footwear plus socks, chemical-resistant gloves made of any waterproof material, protective eyewear and chemical-resistant headgear for overhead exposure.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind directions, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

For ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

PRODUCT INFORMATION

This product combines Quinclorac, 2,4-D and Dicamba to provide broad-spectrum control of grassy and broadleaf weeds.

- Postemergent control of a comprehensive list of both grassy and broadleaf weeds.
- Excellent clean-up product for areas that did not receive a preemergent grass herbicide application in the spring.
- Controls newly germinated 1 to 2-leaf crabgrass, to 1-tiller crabgrass and when crabgrass has matured to 5-tillers or greater.
- Quinclorac contributes grassy weed control and is absorbed by foliage and roots and translocated throughout the plant.

Susceptible grasses demonstrate stunting, chlorosis, and gradual reddening followed by necrosis and death.

USE SITES

This product may be applied postemergence to residential and non-residential turfgrasses. Examples of such sites include, but are not limited to: grounds or lawns around residential and commercial establishments, multi-family dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses and sod farms.

Highly Tolerant Species	
Bluegrass, Annual (Poa annua)	Fescue, Tall
Bluegrass, Kentucky	Ryegrass, Annual
	Ryegrass, Perennial
Moderately Tolerant Species	
Bentgrass, Creeping ^{1,3}	Fescue, Chewing's
Bermudagrass, Common ¹	Fescue, Fine ²
Bermudagrass (Hybrids) ¹	Fescue, Hard
Bluegrass, Rough (Poa trivialis)	Fescue, Red
	Zoysiagrass
Susceptible (Do Not Use on these grass species)	
Bahiagrass	Centipedegrass
Bentgrass, Colonial	Dichondra
Bentgrass, Seaside	Paspalum, Seashore
Buffalograss	St. Augustinegrass
Carpetgrass	
1) Yellowing that may occur on these species can be reduced by the addition of chelated iron or sprayable soluble nitrogen.	
2) Only apply to fine fescue when it is part of a blend.	
3) Application to established creeping Bentgrass must be applied in 2 to 3 split applications at 1.1 to 1.9 ounces per 1,000 sq. ft.	
Do not exceed 1.5 pounds a.i. Quinclorac per acre per year. Sequential application(s) should be timed 21 days apart.	
Broadleaf weed control may be reduced due to sequential applications. Do NOT use on golf course greens, tees and collars.	
Do NOT use on lawns with desirable clovers or legumes or on ornamentals.	

Warm season grasses such as Common Bermuda and Zoysia may see temporary discoloration. Use reduced rates if grass is stressed from heat or drought. Do not apply during growth stages from dormancy to green-up and from green-up to dormancy.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed the maximum labeled rate for the turf type where the product is to be applied. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

TIMING

Apply this product to actively growing weeds as a postemergence broadcast or spot spray. Follow-up applications may be required for dense infestations of broadleaf and grassy weeds. Under certain conditions, application of this product made to annual grasses at 2 to 4-tiller may not provide complete control.

NEWLY SEEDED AREAS:

The application of this product to grass seedlings is not recommended until after the third mowing.

NEWLY SODDED, SPRIGGED, OR PLUGGED AREAS:

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations. Delay applications for 4 weeks after seeding and emergence of Kentucky Bluegrass, Perennial Ryegrass and Fine Fescues.

SEEDING:

Delay applications for 4 weeks after seeding and emergence of turf species listed on this label.

MOWING:

Do not mow for two days before or two days after application. Clippings for the first three mowings should be left in the application area. Do not use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

IRRIGATION AND RAINFALL

If soil moisture is not sufficient prior to application, irrigation may improve weed control. For best results, DO NOT water or irrigate for 24 hours after application. If rainfall does not occur in 2 to 7 days after application, irrigation of at least one-half inch is required.

Do not apply this product by air or through any type of irrigation equipment.

MIXING INSTRUCTIONS

Begin with a clean spray tank. Fill the spray tank with one-half the required amount of clean water. Slowly add this product while agitating, then complete filling the tank with water. Maintain continuous agitation until spraying is complete. If left standing for extended periods of time, re-agitate to assure uniformity of the spray mixture.

Adding adjuvants may cause slight leaf burn, but turf vigor is not reduced. Delaying applications when relative humidity and temperatures are high may help to avoid potential for leaf burn and turfgrass damage. Low mowing heights may also increase the possibility of turf injury. The addition of chelated iron or sprayable solution nitrogen fertilizer will reduce slight yellowing.

For best results, the addition of methylated seed oil is recommended when it meets all of the following criteria:

- be non-phytotoxic
- contain only EPA-exempt ingredients
- provide good mixing quality in the jar test
- be successful in local experience

Do not include additives when tank mixing with emulsifiable concentrate (EC) products as this may cause phytotoxicity. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

This product may be tank mixed with EPA-registered preemergent herbicides (if compatible) for extended residual control.

The following compatibility test should always be performed prior to full-scale tank mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product).
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

Sprayer Cleaning: Clean application equipment thoroughly before and after application to prevent cross-contamination. Use a strong detergent or approved spray tank cleaner and rinse thoroughly.

SPRAYING INSTRUCTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Do not apply at wind speeds greater than 15 mph.

Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Avoid broadcast applications when air temperature exceeds 90°F. When using small, spot treatment applications in temperature over 90°F, turf injury may occur.

SPRAY VOLUMES:

Higher water volumes are generally required to control grassy weeds. Use 20 to 300 gallons per acre (0.5 to 6.9 gallons per 1,000 square feet) and a spray pressure of 20 to 40 psi. Higher water volumes should be used for control of dense weed populations to ensure weed foliage is completely covered.

LOWER VOLUME EQUIPMENT:

Use at least 0.5 gallons of water per 1,000 square feet. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

NOTE:

For all grasses (1) do not overlap spray patterns; (2) use reduced rates if grass is stressed from heat, drought, etc.; and (3) follow CDA equipment spray instructions.

Make no more than 2 applications per year.

POSTEMERGENT BROADLEAF WEED CONTROL

This product will control or suppress the following list of broadleaf weeds. For best results, apply this product when weeds are actively growing and in early stages of growth. More mature weeds will be more difficult to control and may require a second application. Mature, drought-stressed weeds will be more difficult to control so adequate soil moisture is preferred. Adverse or extreme environmental conditions such as poor soil conditions, high temperatures, drought and cultural conditions may affect the performance of this product.

Do not broadcast apply this product above 90°F. Spot treatments above 90°F may result in some turf injury.

BROADLEAF WEEDS CONTROLLED

Alder	Common sowthistle
Annual yellow sweet clover	Corn chamomile
Artichoke	Creeping Jenny
Austrian fieldcress	Crimson clover
Bedstraw	Croton
Beggartick	Cudweed
Biden	Curly dock
Bindweed	Curly indigo
Bird vetch	Dandelion
Bitterweed	Dead nettle
Bitter wintercress	Dock
Black-eyed Susan	Dollar weed
Black medic	Dogbane
Black mustard	Dogfennel
Black-seed plantain	Elderberry
Blessed thistle	English daisy
Blue lettuce	Fall dandelion
Blue vervain	False dandelion
Boxelder	False flax
Bracted plantain	False sunflower
Brassbuttons	Fiddleneck
Bristly oxtongue	Field bindweed
Broadleaf dock	Field pansy
Broadleaf plantain	Fleabane (daisy)
Broomweed	Flixweed
Buckhorn	Florida betony
Buckhorn plantain	Florida pusley
Bulbous buttercup	Frenchweed
Bull nettle	Galinsoga
Bull thistle	Garlic mustard
Burdock	Goathead
Burning nettle	Goatsbeard
Bur ragweed	Goldenrod
Burweed	Ground ivy
Buttercup	Gumweed
Canada thistle	Hairy bittercress
Carolina geranium	Hairy fleabane
Carpenterweed	Hawkweed
Catchweed bedstraw	Healall
Catsear	Heartleaf drymary
Catnip	Hedge bindweed
Chickweed	Hedge mustard
Chicory	Hemp
Cinquefoil	Henbit
Common chickweed	Hoary cress
Common mullein	Horsetail

(continued)

BROADLEAF WEEDS CONTROLLED *(continued)*

Indiana mallow	Smooth pigweed
Ironweed	Sneezeweed
Jewelweed	Southern wild rose
Jimsonweed	Sowthistle
Kochia	Spanishneedle
Knawel	Spatterdock
Knotweed	Speedwell
Lambsquarter	Spiny amaranth
Lespedeza	Spiny cocklebur
Locoweed	Spotted catsear
Lupine	Spotted knapweed
Mallow	Spotted spurge
Marshelder	Spurge
Matchweed	Spurweed
Mexicanweed	Stinging nettle
Milk vetch	Stinkweed
Milkweed bloodflower	Stitchwort
Mugwort	Strawberry clover
Morningglory	Sumac
Mousear chickweed	Sunflower
Musk thistle	Sweet clover
Mustard	Tall nettle
Narrowleaf plantain	Thistle
Narrowleaf vetch	Tick trefoil
Nettle	Toadflax
Orange hawkweed	Trailing crownvetch
Oxalis	Tumble mustard
Oxeye daisy	Tumble pigweed
Parsley-piert	Tumbleweed
Parsnip	Velvetleaf
Pearlwort	Venice mallow
Pennycress	Veronica
Pennywort	Vervain
Peppergrass	Vetch
Pepperweed	Violet, wild
Pigweed	Virginia buttonweed
Pineywoods bedstraw	Virginia creeper
Plains coreopsis	Virginia pepperweed
Plantain	Wavyleaf bullthistle
Poison hemlock	Western clematis
Poison ivy	Western salsify
Poison oak	White clover
Pokeweed	White mustard
Poorjoe	Wild aster
Povertyweed	Wild buckwheat
Prostrate spurge	Wild carrot
Prostrate vervain	Wild four-o'-clock
Puncture vine	Wild garlic
Purslane	Wild geranium
Ragweed	Wild lettuce
Red clover	Wild marigold
Redroot pigweed	Wild onion
Red sorrel	Wild parsnip
Redstem filaree	Wild radish
Rough cinquefoil	Wild rape
Rough fleabane	Wild strawberry
Roundleafed marigold	Wild sweet potato
Rush	Wild vetch
Russian pigweed	Willow
Russian thistle	Witchweed
St. Johnswort	Woodsorrel
Scarlet pimpernel	Woolly croton
Scotch thistle	Woolly morningglory
Sheep sorrel	Woolly plantain
Shepherdspurse	Wormseed
Slender plantain	Yarrow
Smallflower galinsoga	Yellow rocket
Smartweed	Yellowflower pepperweed
Smooth dock	and other broadleaf weeds.

POSTEMERGENT CONTROL OF GRASSY WEEDS

This product can provide control and suppression of certain grassy weeds. For best results, apply this product when weeds are actively growing and in early stages of growth. More mature grasses will be more difficult to control and may require a second application. Mature, drought-stressed grassy weeds will be more difficult to control so adequate soil moisture is preferred. Adverse or extreme environmental conditions such as poor soil conditions, high temperatures, drought and cultural conditions may effect the performance of this product. Do not broadcast apply this product above 90°F. Spot treatments above 90°F may result in some turf injury.

GRASSY WEEDS CONTROLLED IN ORNAMENTAL LAWNS AND TURF

COMMON NAME	COMMENTS
Barnyardgrass Crabgrass, Large Crabgrass, Smooth Foxtail, Giant Foxtail, Green Foxtail, Yellow Signalgrass, Broadleaf	Under certain conditions annual grasses at the 2 to 4-tiller stage may not be completely controlled and a sequential application may be needed at 14 to 21 days. Do not exceed two applications per year.

USE RATES

This product may be applied postemergence to residential and non-residential turfgrasses according to the following rate chart. Examples of such sites include, but are not limited to: grounds or lawns around residential and commercial establishments, multi-family dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses and sod farms. Not for use on sod farms in Arizona. For use in New York by spot treatment only.

APPLICATION GUIDELINES			
APPLICATION	RATE	SPRAY VOLUME	MAXIMUM APPLICATIONS
Single	7 to 8 pints/A or 112 to 128 ounces/A	20 to 300 gal/A or 0.5 to 6.9 gal/1,000 sq. ft.	2 per year
Spot Treatment	2.6 to 2.9 fluid ounces per 1,000 sq. ft.	1 gal/1,000 sq. ft.	

Application Guidelines:

- Ensure that spray volumes are adequate to completely cover weeds, especially when dense weed infestations make it difficult to completely cover the foliage of target weeds.
- Early applications of this product will not control weeds germinating later in the season so a second application may be necessary.
- If turf injury is observed after the first application, delay the second application until turf has recovered.
- Dual Applications: A single application should control less mature, light weed populations; however, a second application may be required to control heavy, mature broadleaf weed infestations. Under certain conditions, annual grasses in the 2 to 4-tiller stages may require a sequential application. If a second application is required, allow 14 to 21 days between applications.
- For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration. **DO NOT** burn unless allowed by state and local ordinance. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations of liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV062812)

LESCO is a registered trademark of LESCO Technologies, LLC.

Momentum is a registered trademark and used by LESCO, Inc. with permission of the trademark owner

NOTES

NOTES

NOTES

LESCO® Momentum® Q Herbicide

FOR SELECTIVE BROADLEAF AND CERTAIN GRASSY WEED CONTROL

Contains 2,4-D, Quinclorac and Dicamba

Controls: Dandelion, Chickweed, Black Medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle, Large and Smooth Crabgrass and many other species of broadleaf and certain grassy weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* 13.24%
Quinclorac (3,7-Dichloro-8-quinolinecarboxylic Acid)** 8.25%
Dicamba (3,6-Dichloro-o-Anisic Acid)*** 1.38%

OTHER INGREDIENTS: 77.13%
TOTAL: 100.00%

By Isomer Specific AOAC Method, Equivalent to:

* 2,4-Dichlorophenoxyacetic Acid 11.34%, 1.00 lbs./gal.

** 3,7-Dichloro-8-quinolinecarboxylic Acid 8.25%, 0.75 lbs./gal.

*** 3,6-Dichloro-o-Anisic Acid 1.38%, 0.125 lbs./gal.

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE LABEL BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or on clothing.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

The chemicals in this product have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration. **DO NOT** burn unless allowed by state and local ordinance. If burned, stay out of smoke.

EPA Reg. No. 228-531

Manufactured by: Nufarm Americas Inc. • 11901 S. Austin Avenue • Alsip, IL 60803

LESCO is a registered trademark of LESCO Technologies, LLC.

Momentum is a registered trademark and used by LESCO, Inc. with permission of the trademark owner.

EPA Est. No. 228-IL-1

NET CONTENTS:
64 oz (1.89 L)

PULL HERE TO OPEN
PRESS TO RESEAL

(RV062812) 12796000 [1]