



# Stonewall® 0.43% Plus Fertilizer

Read entire label before using this product. Use only according to label instructions.

NOTICE: Before using this product, read the Directions for Use, Storage and Disposal Instructions, and Warranty Statements. If the Warranty statements are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

Contains LESCO® Poly Plus® Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding.

ACTIVE INGREDIENT: Prodiamine: (N3, N3-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine).....0.43%
INERT INGREDIENTS:.....99.57%
Total:.....100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION
See the following sections Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.

FIRST AID
If swallowed: Call a poison control center or doctor immediately for treatment advice.
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes.
If on skin or clothing: Take off contaminated clothing.
If inhaled: Move person to fresh air.

HOTLINE NUMBER
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin.

ENVIRONMENTAL HAZARDS
This product has low solubility in water. At the limits of solubility, this product is not toxic to fish.

STORAGE AND DISPOSAL
Do not contaminate water, foodstuffs, feed or seed by storage or disposal.
PESTICIDE STORAGE: Store this product only in its original container in a dry, cool, secured storage area out of reach of children.

DIRECTIONS FOR USE
-It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
-This product is a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses.

USE PRECAUTIONS
The following use precautions apply to the use of this product in turf grasses and lawns:
-Application of this product may thin emerged annual bluegrass and newly over-seeded grasses.

APPLICATION DIRECTIONS
ESTABLISHED TURF
This product is a selective pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grass and lawns.



Region Date
Upper North After May 10
North After April 20
Transition After March 20
South Jan 1 - March 20

RATES OF APPLICATION
This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year.

MAXIMUM ANNUAL RATES
This product is recommended for use on the turf grass species listed below. Do not exceed the maximum yearly rate:

Table with 3 columns: Turf Species, lb LESCO Stonewall 0.43% Plus Fertilizer / acre, lb AI per Acre. Includes species like Bermudagrass, Bahiagrass, Centipedegrass, etc.

1. These are the maximum rates per calendar year by species limitations.
2. May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 121 lb/Acre.

SUGGESTED SPREADER SETTINGS\*
Recommended Rates in lb per 1,000 sq ft
Spreader 3.44 lb 4.02 lb 5.34 lb 8.01 lb

\*IMPORTANT: These settings are only approximate. Age, condition of the spreader, and operator speed can cause wide variation.

APPLICATION TIMING AND INFORMATION
This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top, or directed application.

24-0-11
GUARANTEED ANALYSIS
TOTAL NITROGEN (N) .....24.00%
24.00% Urea Nitrogen\*
SOLUBLE POTASH (K2O)..... 11.00%
MAGNESIUM (Mg) .....0.50%
0.01% Water Soluble Magnesium (Mg)
IRON (Fe) .....1.00%
0.01% Water Soluble Iron (Fe)
MANGANESE (Mn) .....0.50%
0.01% Water Soluble Manganese (Mn)
DERIVED FROM: Polymer Coated Urea, Urea, Muriate of Potash, Iron Sucrate, Manganese Sucrate, Magnesium Sucrate.

WEEDS CONTROLLED
When used in accordance with this label, this product will provide control of the following weeds:
Barnyardgrass, Goosegrass1, Lovegrass, Speedwell, Bluegrass, Annual (Poa annua), Henbit, Panicum (Texas, Fall, Brown Top), Sprangletop, etc.

1. In many areas a single application of 150 to 349 lb/acre of this product will control the above listed weeds and goosegrass.
2. Suppression only. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species.

WHEN TO APPLY AFTER OVERSEEDING
Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer.

WHEN TO OVERSEED AFTER APPLICATION
This product will inhibit the germination of turf species if overseeded too soon after application.

Table with 3 main columns: LESCO Stonewall 0.43% Plus Fertilizer Rate (lb/Acre), Interval Before Overseeding, and sub-columns for North, Transition, South.

LANDSCAPE ORNAMENTALS
(Including Established Perennial and Wildflower Plantings)
This product may be applied for residual pre-emergence weed control in ornamentals.

TOLERANT ORNAMENTAL SPECIES
This product will not harm most established ornamental trees, shrubs, and flowers.
Table with 4 columns: Scientific Name, Common Name, Scientific Name, Common Name.

Table with 4 columns: Scientific Name, Common Name, Scientific Name, Common Name. Lists various plant species and their compatibility with the fertilizer.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY
NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114.

#702474