

Dimension[®]

DIMENSION[®] SPECIALTY HERBICIDE

SPECIALTY HERBICIDE

Dimension[®] specialty herbicide provides preemergence and early postemergence control of crabgrass, along with preemergence control of more than 45 other grassy and broadleaf weed species. It offers proven control wherever it's applied — residential and commercial lawns; golf course and sports turf; landscape ornamentals; container and field-grown nursery ornamentals; commercial sod farms; and noncrop and industrial sites.

Not only is Dimension effective for preemergence control of crabgrass, but it also provides early postemergence control when treated prior to tillering — up to five leaves. The five-leaf stage generally corresponds to the time when crabgrass seedlings are easily visible in a lawn or turf. The benefit of early postemergence control is the longer application window it offers compared with products that only provide preemergence control.

Dimension 2EW provides the same proven results you've come to expect from previous formulations, with the flexibility to be applied over the top of more than 440 landscape, 140 field- and 125 container-grown ornamental plants. New ornamentals for over-the-top application are continually being added to the label. In addition to being labeled for use on many popular ornamental species, the



concentrated 2EW formulation has a lower odor, is easy to measure and has low use rates.

You can count on any of the convenient formulations to work without ever staining surroundings. Available sprayable formulations include Dimension 40WP, Dimension 2EW and Dimension EC. Straight granular and on-fertilizer formulations are also available.

How Dimension Works

Dimension[®] specialty herbicide is a member of the pyridine family of chemistry. When the active ingredient, dithiopyr, is absorbed by the roots of susceptible plants, they swell and stop growing.

Activity is limited to points of growth, such as root tips. And because Dimension offers early postemergence crabgrass activity, the window for application is two to eight weeks longer than other herbicides currently on the market that only provide preemergence control.

Weeds Controlled

Dimension will control or suppress more than 45 annual grassy and broadleaf weeds, including:

- Annual bluegrass (*Poa annua*)
- Barnyardgrass
- Chickweed
- Crabgrass (preemergence and early postemergence)
- Goosegrass
- Henbit
- Kikuyugrass
- Oxalis
- Spurge

Turf Recommendations

Dimension can be used on established seeded, sodded or sprigged turf. Be sure the turf has a well-developed root system and a uniform stand, and has received at least two mowings before applying Dimension the first time. Always apply at the recommended rate according to label directions depending on your geography. In preemergence applications, Dimension[®] specialty herbicide may be applied up to eight weeks before crabgrass germination; in early postemergence applications, it may be applied up to tillering. On golf courses, it can be used on fairways, roughs