A Horticultural Vinegar Biopesticide-for Non-Selective Control of Herbaceous Broadleaf Weeds and Weed Grasses which Surround Food Crops, Non-food Crops and Non-production Agricultural, Farmstead, Right-of-Way and Institutional Land Sites. FIRST AID

IF IN EYES: Hold eyelid open and flush with a steady, gentle stream of water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center of doctor for advice. IF ON SKIN OR CLOTHING: Take off contaminated clothing, Rinse skin

immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice. IF SWALLOWED: Call a poison control center or doctor immediately for

treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center or doctor. Do not give anything by mouth to an unconscious person. IF INHALED: Move person to fresh air. If person is not breathing, call 91 or ambulance, then give artificial respiration, preferably mouth to mouth, if possible, Call a poison control center or doctor for further treatment advice. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the

use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact -800-858-7378 for emergency medical treatment.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER-PELIGRO

Corrosive-causes irreversible eve damage. Wear goggles or face shield when handling, Harmful if absorbed through skin, Harmful if swallowed. Do not get in eyes, on skin, or on clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling. Wear personal protection equipment when

handling and/or applying. PERSONAL PROTECTION EQUIPMENT (PPE): Applicators and other handlers must wear appropriate protective eve wear, such as face shield or goggles and face mask (with MSHA/NIOSH approval number prefix such as N95, R-95, or P-95), long sleeved shirt and long pants.

waterproof gloves and shoes plus socks. USER SAFETY REQUIREMENTS: Follow manufacture's instructions for cleaning/maintaining PPE. If no such instructions exist for PPE, use

detergent and hot water. Keep and wash PPE separately from other USER SAFETY RECOMMENDATIONS: Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after

handling this product. Wash the outside of gloves before removing. As soon as possible, wash throughly and change into clean clothing. ENVIRONMENTAL HAZARDS: This pesticide is toxic to birds exposed to spray drift, direct treatment or residues on crops or weeds. Do not apply this product or allow to drift to crops or weeds if birds are actively visiting the treatment area. This pesticide is toxic to fish and aquatic invertebrates. For terrestrial use only. Do not apply directly to

DIRECTIONS FOR USE: It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using. Do not apply this product in a way that will contact workers or other persons. either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe. consult the State or Tribal agency responsible for pesticide regulation. Endangered Species Protection Requirements: This product may have effects on endangered species. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county in which you are applying the product. To obtain Bulletins, no more than six months before using this product, consult http://www.epa.gov/espp/ or call 1-800-447-3813. You must use the Bulletin valid for the month in which you will apply the product. Avoid application to reactivate metals such as aluminum, tin, iron or items

such as fencing or lawn furniture in order to prevent staining, mottling, or otherwise interfering with finished metal surfaces. In case of contact, rinse the sprayed surfaces with water. Avoid repeated applications on concrete, since this product, as all other acids may result in etching of cement.

AGRICULTURAL USE REQUIREMENTS: Use this product only in accordance with its labeling and with the Worker Protection Standard., 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to statements on this label about personal protective equipment (PPE) and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry to treated areas during the restricted entry interval (REI) of 48 hours. PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water) is:

*Coveralls *Waterproof gloves *Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS: The requirements in this box apply to uses of this product that are NOT Within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Do not enter or allow worker entry to treated areas during the restricted entry interval (REI) of 48 hours. Keep unprotected persons out of the treated areas until spray has dried.

Medina Agriculture Products Co., Inc. 20% Vinegar is a guick-acting, non-selective contact herbicide containing an organic acid for which the residues degrade promptly in the soil. Foliar contact results in rapid desiccation and control of annual weeds and grasses, and top growth reduction of herbaceous perennial weeds and grasses. For control of established perennial weeds, retreatment may be necessary. Medina Agriculture Products Co., Inc. 20% Vinegar may be applied up to two

days before harvest.

TYPICAL WEEDS CONTROLLED: ANNUAL BROADLEAF WEEDS: Black medic, Chickweed, Black Nightshade, Cinquefoil (rough), Common groundsel, Hairy nightshade, Lambsquarters, Mustard spp., Oxalis spp., Pigweed spp., Ragweed spp., Shepherdspurse,

PERENNIAL GRASSES: Quackgrass, Witchgrass, Bluegrass,

Smartweed (Ladysthumb), Velvetleaf, PERENNIAL BROADLEAF WEEDS: Amaranth spp. Cinquefoil (silvery). Dandelion, Ground Ivy, Plantain spp., Toadflax, Tufted vetch, Wild carrot, Moss. ANNUAL GRASSES: Crabgrass, Foxtail spp., Italian ryegrass, Poa annua.



For Organic Production

Active Ingredient: Acetic Acid......20.0%* Other Ingredients.....80.0%

*Equivalent to 200 grain vinegar by titration

NOT FOR HUMAN CONSUMPTION

DANGER - KEEP OUT OF REACH OF CHILDREN

PELIGRO - MANTENER FUERA DEL ALCANCE DE LOS NIÑOS If you do not understand the label, find someone to explain it to you in detail. See label side panels for First Aid and additional precautionary statements. Si no entiendes la etiqueta, busca a alguien que te la explique detalladamente. Consulte los paneles laterales de la etiqueta para conccer 🧗 primeros auxilios y declaraciones de precaución adicionales.

Net Contents 128 Fl. oz (1 GAL.) 3.8L

DO NOT SPRAY DIRECTLY ONTO FOOD CROPS Use this product to control weeds that surround all food crops, nonfood crops (ornamentals) and non-production agricultural, farmstead, right-of-way, and institutional land sites. Use to control weeds in interstate freeways/highways, federal and state highways and city/county roads and walkways, railroad right-of-way, tank farms, power stations and easements, and other rights-of-way, industrial sites, and vacant lots. Suitable to keep down weeds on all botanical and private association ornamental gardens, kennels, dog runs, and other animal enclosures and walkways. **DRIFT MANAGEMENT:** Drift is the off site movement of spray away from the intended target.

Managing drift is important for the protection of endangered species. In addition, this product is non-selective so it will severely injure or kill any crop plants or other plants it cor in contact with. Do not apply this product by aerial application. Do not apply when wind conditions favor movement of spray away from the site of application. When applying this product in a broadcast application utilize a hooded sprayer, shielded or other spray equipment that prevents the movement of spray away from the target. Utilize nozzles that are product designed to produce large spray droplets to reduce drift

EARLY SEASON ANNUAL WEED CONTROL: Apply this product when weeds are small (3 to 5 leaf stage) and actively growing. Spray this product on unwanted weeds (vegetation) to the point of wetness. For best results spray in full sunshine at temperatures above 65 degrees Fahrenheit, Since dilution of this product will reduce effectiveness, do not dilute, Avoid spraying landscape plants, ornamentals and other desirable foliage and this product causes

NOTE: ALL CONTACTED VEGETATION WILL BE AFFECTED, DO NOT ALLOW THIS PRODU TO COME IN CONTACT WITH DESIRABLE PLANTS. OVER SPRAY OR DRIFT WILL INJURE OR KILL CONTACTED VEGETATION INCLUDING ANY CROP PLANTS.

FURF AND ORNAMENTALS: Such as but not limited to woody ornamentals, potted crops. foliage plants, greenhouses, shade houses, and any other plant production structure in which the product can be applied without contacting desirable plants. Applications to weeds in turf will kill surrounding turf. BULB VEGETABLES: Such as but not limited to onion, green onion, shallots and garlic, LEAFY VEGETABLES: Such as but not limited to celery, head lettuce, leaf lettuce, spinach. Do not broadcast within the growing crop. Apply only with shielded or hooded spray equipment to weeds growing between crop rows. FRUITING VEGETABLES: Such as but not limited to tomato, bell pepper, chili pepper, eggplant and okra. Do not broadcast within the growing crop. Apply only with shielded or hooded spray equipment to weeds growing between crop rows. POME FRUIT: Such as but not limited to apple and pear. For all fruit crops spray must be directed under the crop canopy. Application equipment must be hooded or shielded. Any contact with crop will cause crop injury, CUCURBIT VEGETABLES: Such as but not limited to cucumber, squash, cantaloupe, watermelon, Do not

broadcast within the growing crop. Apply only with shielded or hooded spray equipment to weeds growing between crop rows.

FOR SPOT TREATMENT APPLICATION: Direct spray with a single nozzle or a hand-gun applicator to thoroughly wet undesirable weed foliage. Re-treatment of perennial weed

growth may be required for control. FOR BROADCAST SPRAY APPLICATIONS: For broadcast spray application to control weeds Application types include pre-plant (prior to crop establishment), preharvest desiccation (to kill crop foliage prior to harvesting the crop) postharvest crop desiccation (remaining crop residues after crop has been harvested). Use 15 to 30 gallons of this product per acre. Thorough coverage is necessary to achieve desirable control. For dense weed populations and/or heavy foliage, use the higher spray volume to endure adequate coverage. Apply with flat fan nozzles and at sufficient pressure to achieve required coverage, but do not exceed a pressure 35 PSI. When applying this product in a broadcast application within a growing crop, utilize a hooded sprayer, shielded, or other spray equipment that prevents the movement of spray away from the target. For pre-plant burn down of young vegetation, use of wide angle nozzles (110 degrees or wider) spaced close together to enable the boom to be positioned

lower and closer to the spray target, reducing the potential for drift. POST-DIRECTED: (Row middles, shielded sprayer or hooded sprayer); For spray application to control weeds, apply 15 to 30 gallons of this product per acre. Thorough coverage is necessary to achieve desirable control. For dense and/or heavy weed populations, use the higher spray volume to ensure adequate coverage. Apply with flat fan nozzles and at the lowest pressure needed to achieve required coverage, but do not exceed 35 PSI. If treating

along the side of crop plants, shielded or hooded sprayers must be used to avoid crop injury

SPRAY EQUIPMENT: Do not apply this product by aerial application. Do not apply when wind conditions favor movement of spray away from the site of application. When applying this product in a broadcast application utilize a hooded sprayer, shielded, or other spray equipment that prevents the movement of spray away from the target. Utilize nozzles that are designed to produce large spray droplets to reduce drift. Spray nozzles need to be plastic or 316 stainless steel as this product will eventually corrode any other metal.

FOR CONTROL OF LARGER ANNUALS AND BURNDOWN OF PERENNIAL WEED GROWTH: Larger annual weeds and perennials are more difficult to control and may nes require retreatment. Treat initially as recommended above and repeat if new growth o leaves appears. Thorough coverage of all weed foliage is necessary to achieve desirable control. To reduce the amount of product required to control well-established weed stands, mow to a height of 4-6 inches and remove cut vegetation before applying this

TIMING OF APPLICATIONS: Best results are achieved from applications to actively growing young weeds. Weeds that are mature, dormant, or hardened due to moisture stress are more tolerant of herbicide treatments. For best results spray in full sunshine at temperatures above 65°F. Only contacted vegetation will be affected

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM. NOTE: ALL CONTACTED VEGETATION WILL BE AFFECTED DO NOT ALLOW THIS PRODUCT TO COME IN CONTACT WITH DESIRABLE PLANTS, OVER SPRAY OR DRIF WILL INJURE OR KILL CONTACTED VEGETATION INCLUDING ANY CROP PLANTS.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage and disposal. PESTICIDE STORAGE: Keep from freezing. Store only in original tightly sealed container and out of reach of children. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, Clean container promptly after emptying, Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closure. Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

WARRANTY STATEMENT: To the extent required by law, Medina Agriculture Products Co., Inc. warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with the label directions. Medina Agriculture Products Co., Inc. makes no other express or implied warranties either of the merchantability of fitness for a particular use and shall not be liable for misuse or conditions beyond its control, such as handling or storage.

> Medina Agriculture Products Co., Inc. P.O. Box 309 • Hondo, TX 78861

> medinaag.com • 830-426-3011 EPA Registration No: 85208-1-90366

