

Monterey Consan[®] 20

Disinfectant • Deodorizer • Fungicide

ACTIVE INGREDIENTS:

Alkyl (C₁₄ 60%, C₁₆ 30%, C₁₂ 5%, C₁₈ 5%) dimethyl benzyl ammonium chloride 10.0%

Alkyl (C₁₂ 68%, C₁₄ 32%) dimethyl ethylbenzyl ammonium chloride..... 10.0%

INERT INGREDIENTS:.....80.0%

TOTAL:100.0%

EPA Reg. No.: 10324-94-54705

EPA Est. No.: 48498-CA-1

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

PELIGRO: SI NO PUEDE LEER EN INGLES, PREGUNTE A SU SUPERVISOR SOBRE LAS INSTRUCCIONES DE USO APROPIADAS ANTES DE TRABAJAR CON ESTE PRODUCTO.

DANGER: IF YOU CANNOT READ ENGLISH, ASK YOUR SUPERVISOR TO EXPLAIN THE APPROPRIATE DIRECTIONS FOR USE BEFORE WORKING WITH THIS PRODUCT.

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Keep Out of Reach of Children. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or clothing. Wear goggles or face shield and rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT: Applicators and other handlers must wear coveralls worn over long-sleeved shirt and long pants, waterproof gloves, chemical-resistant footwear and socks, protective eyewear, chemical-resistant headgear when using this product for algae control in overhead watering system and chemical-resistant apron when mixing, loading or cleaning equipment. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARD

This product is toxic to fish.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Do not mix with oxidizers, anionic soaps and detergents.

Net Contents: ☐ 1 Pint ☐ 1 Quart ☐ 1 Gallon

DISTRIBUTED BY:

Lawn and Garden Products, Inc.

PO Box 35000 • Fresno, CA 93745

(559) 499-2100 — info@montereylawnngarden.com

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Treatment is restricted to indoor uses or uses in which controlled conditions to prevent runoff and exposure to the environment. In the case of diseased trees, the spot treatment of trees must be at least 100 feet from any pond, lake, stream, or river to prevent possible runoff of the product into the waterway. **DO NOT APPLY BY AERIAL SPRAY.**

GENERAL DISINFECTION PERFORMANCE IN NON-MEDICAL FACILITIES: At 1.5 ounces of this product per 5 gallons of water use level (or equivalent use dilution) (469 ppm active) and with a 10-minute contact time, this product is bactericidal according to the AOAC Use Dilution Test method on hard, non-porous surfaces modified in the presence of 5% organic serum against: *Campylobacter jejuni*, *Escherichia coli* 0157:H7, *Listeria monocytogenes*, *Proteus mirabilis*, *Salmonella enterica*, *Staphylococcus aureus*, *Staphylococcus aureus* Methicillin resistant, *Yersinia enterocolitica*.

FUNGICIDAL PERFORMANCE: At 2 ounces of this product per 5 gallons of water use level (or equivalent use dilution) (625 ppm active) and with a 10-minute contact time, this product is effective in the presence of 5% blood serum against the following on hard non-porous surfaces: *Trichophyton mentagrophytes*, *Aspergillus niger*.

HORTICULTURAL AND ORNAMENTAL PLANT

MONTEREY CONSAN 20 is a general disinfectant. It will help in the control of fungal, bacterial and viral plant pathogens, bacterial and fungal leaf rots, slime forming fungi, and odor causing bacteria and algae. It is designed specifically for use where disease cross contamination may occur between plants. Pre-clean all surfaces prior to application of the disinfectant solution. All inanimate surfaces must be thoroughly wetted and remain wet for 10 minutes for disinfection. Then wipe up excess or allow to air dry. Fresh solution must be prepared daily and when use solution becomes visibly dirty. To apply solution for sprayer applications, use as a coarse spray only.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE). The requirements in this section only apply to uses of this product that are covered by the Worker Protection Standard.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard must wear: coveralls worn over long-sleeved shirt and long pants, waterproof gloves, chemical-resistant footwear and socks, protective eyewear, chemical-resistant headgear when using this product for algae control in overhead watering systems and chemical-resistant apron when mixing, loading or cleaning equipment.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Users must: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours. Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls worn over long sleeved shirt and long pants, waterproof gloves, chemical-resistant shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this section (box) apply to uses of Monterey Consan 20 that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40CFR170). The WPS applies when Monterey Consan 20 is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Applicators and other handlers must wear long sleeved shirt and long pants, shoes and socks, eye protection and rubber gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

CHEMIGATION: Do not apply Monterey Consan 20 through any type of irrigation system except when used to control algae in nozzles as directed on this label.

DISINFECTING

The requirements in this section (box) apply to uses of Monterey Consan 20 that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40CFR170). The WPS applies when Monterey Consan 20 is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

WORK AREAS AND BENCHES: Spray or swab hard nonporous working surface with a solution of 1.5 oz. of MONTEREY CONSAN 20 per 5 gallons of water (or equivalent use dilution) before each work period and again after each plant is completed to help control transfer of diseases such as *Botrytis*, crown rot, downy mildew, *Erwinia* and root rot. Allow surface to remain wet for 10 minutes. To apply solution as a sprayer applications, use as a coarse spray only and spray 6-8 inches from surface. Allow to air dry. Change cloth, sponge or towels frequently to avoid redeposition of soil. Prepare a fresh solution at least daily or more often if use solution becomes visibly dirty.

POTS, FLATS AND FLOWER BUCKETS: Brush or wash used pots and flats, then soak in a solution of 1.5 oz. of MONTEREY CONSAN 20 per 5 gallons of water (or equivalent use dilution) for 10 minutes to help control transfer of diseases such as anthracnose, blight, *Botrytis*, crown rot, *Erwinia*, mildew, root rot and rust. Allow to air dry. Prepare a fresh solution at least daily or more often if use solution becomes visibly dirty.

CUTTING TOOLS: Soak cutting edge of tool for 10 minutes in a solution of 1.5 oz. of MONTEREY CONSAN 20 per 5 gallons of water (or equivalent use dilution) before use to help control transfer of diseases such as *Botrytis*, *Erwinia*, root and stem rots, and Tobacco Mosaic Virus (TMV). Allow to air dry. Prepare a fresh solution at least daily or more often if use solution becomes visibly dirty. Dry and oil tools at the end of each workday.

ALGAE CONTROL (Not for use in CA)

When used as directed, MONTEREY CONSAN 20 is effective against algae. With a little experience, you will quickly recognize algae contamination as it starts. Algae must be killed immediately! Remember – algae can double or quadruple in growth every day. Algae are generally dead when it does not feel slick to the touch and the color has changed. However, algae do not always change color when killed.

GREENHOUSE GLASS: Spray or swab hard nonporous surfaces with a solution of 1.5 oz. of MONTEREY CONSAN 20 per 5 gallons of water (or equivalent use dilution). Allow to air dry. Wash off dead algae with water. Spray clean surface again with the solution. Do not rinse. For sprayer applications, use as a coarse spray only. Allow surfaces to remain wet for 10 minutes then air dry. Prepare a fresh solution at least daily or sooner if use solution becomes visibly dirty.

HOUSEPLANT AND GARDENING CONTAINERS AND EQUIPMENT: To control algae and cross contamination in terrariums, bubbles, and other hard nonporous gardening surfaces, wipe surface with a solution of 1.5 oz of MONTEREY CONSAN 20 per 5 gallons of water (or equivalent use dilution). Lightly spray plants with same solution. Allow surfaces to remain wet for 10 minutes then air dry. Prepare a fresh solution at least daily or sooner if use solution becomes visibly dirty.

WALKWAYS: For heavy infestations, spray or swab surface with a solution of 1.5 oz. of MONTEREY CONSAN 20 per 5 gallons of water (or equivalent use dilution). Let stand for an hour or more then brush and wash away dead algae. Soak area again with the solution. Do not rinse. This product will inhibit the growth of algae on hard nonporous walkways. Allow to dry on the surface and repeat application when algae growth returns. For sprayer applications, use as a coarse spray only. Allow surfaces to remain wet for 10 minutes then air dry. Prepare a fresh solution at least daily or sooner if use solution becomes visibly dirty.

BIRD BATHS: DO NOT use when fish are present. Clean to remove algae growth prior to filling birdbath and spray the exposed surface with a solution of 0.25 oz. (1 tbsp.) of this product per gallon of water. Allow to air dry and brush off dead algae.

FOUNTAINS, WATER DISPLAYS, DECORATIVE POOLS/PONDS AND STANDING WATER: DO NOT use when fish are present. Remove floating type algae by raking, dragging with cable or chain, skimming, or any other suitable method. It is more economical to remove floating type algae before the water is treated. Treat water by “dribbling” this product 1 part of MONTEREY CONSAN 20 to 4 parts water around the edges where visible algae growth is evident. Apply 12 ounces of use solution per 1000 gallons of water. The use solution is often dribbled liberally along the windward side to float across the water and along the leeward side where contamination gathers. Where a spray rig is available, it may be used to apply the use solution instead of the hand “dribbling” method.

Note: Algae does not usually grow in water that is 5 feet or more in depth. However, floating type algae may grow on pond weeds that grow on the surface of the water.

DECORATIVE POOL APPLICATIONS

- Initial Dosage:** 0.17 oz. per 52 gallons of water (5 ppm*)
1 oz. per 312 gallons of water (5 ppm)
1 pint per 5,000 gallons of water (5 ppm)
1/2 quart per 10,000 gallons of water (5 ppm)
- Weekly Maintenance:** Add not more than one-half the initial dosage. When pool is to remain empty for several weeks or longer, spray exposed surfaces with a solution of 0.17 oz. of MONTEREY CONSAN 20 per gallon of water.

OPERATING FOUNTAINS APPLICATIONS

- Initial Dosage:** Add a 0.03 oz. of this product per 52 gallons of water at a time. Repeat until slight patches of foam appear. Then discontinue. Overdose reduces lighting efficiency.
- Weekly Maintenance:** Add not more than 1/2 the initial dosage until slight foam reappears.
- Monthly Maintenance:** Drain and clean bowl. Refill with fresh water and repeat initial treatment. Draining also removes airborne dirt, dust, contamination and alkali buildup.

TREATMENT OF LAWNS *(Use against dollar spot Scierotinia, fusarium blight Fusarium, Southern blight Scierotium rolfsii, rust, toad stools and fairy rings not allowed in CA)*

To control fungal diseases including dollar spot *Scierotinia*, fusarium blight *Fusarium*, leaf spot *Helminthosporium*, brown patch *Rhizoctonia*, fading out *Curvularia*, Southern blight *Scierotium rolfsii*, pythium blight, rust, toad stools, fairy rings and algal build-up in turf grasses, including lawns of GRASS or DICHONDRA, spray a solution of two tablespoons of this product per gallon of water. Treat at the rate of one gallon of solution to 50 square feet (5 ft. x 10 ft.) of lawn. Treat entire lawn to prevent spread of infection at least one foot beyond the edges of isolated diseased areas. Saturate the entire treatment area, wetting all plant and soil surfaces. Repeat spray applications every 10 to 14 days, as needed to control new or established disease conditions. Treatment is required during the warm growing season and must be made when the lawn is dry or damp, but not during wet rainy weather.

Do not graze treated areas or feed clippings to livestock. Do not exceed treatment of 10 acres (2000 gallons) with retreatment at 10 day intervals not to exceed 6 treatments per year.

MOSQUITO CONTROL *(Not for use in CA)*

MONTEREY CONSAN 20 is not intended to be used for broadcast mosquitocide application but rather is limited to small residential ponds, decorative ponds, and similar areas. This product is not to be used in open waterways connected to larger watersheds or in waters that serve as natural habitats for aquatic and amphibious organisms. MONTEREY CONSAN 20 controls mosquitoes where they breed fountains, water displays, decorative pools, decorative ponds, sewage treatment systems, spas, hot tubs, swimming pools and standing water in old tires, empty tin cans, barrels, puddles, and water drains around buildings. Do not exceed 10 applications at a minimum of 10-day intervals per year. Note: Only out-of-season, not-in service, or inactive spas, hot tubs, swimming pools require treatment for mosquito control. Do not treat during the swimming season. Spray from fountains treated with this product will not harm poolside plantings. DO NOT use when fish or other wildlife (for example, amphibians) are present.

When used as directed, MONTEREY CONSAN 20 is effective as an ovicidal, larvacidal, pupacidal control of *aedes aegypti* (Yellow fever carrier) and *Culex nigripalpus* carrier of various diseases of man and animal, such as heartworm and Hepatitis mosquitoes. This product will also reduce larval populations of *Anopheles quadrimaculatus* (Malaria carrier) mosquitoes.

MONTEREY CONSAN 20 controls mosquitoes where they breed fountains, water displays, decorative ponds, spas, hot tubs, swimming pools and standing water in such places as old tires, empty tin cans, barrels, puddles, and water drains around buildings. NOTE: Only out-of-season, not-in-service, or inactive spas, hot tubs, swimming pools require treatment for mosquito control. Do not treat during the swimming season.

Spray from fountains treated with MONTEREY CONSAN 20 will not harm poolside plantings. DO NOT use when fish are present.

INITIAL DOSAGE				
This product		Per	Unit water for 200 ppm*	
Metric	US		Metric	US
3.8 cc	0.13 oz.		3.78 L	1 gal.
38.0 cc	1.28 oz.		37.85 L	10 gal.
380 cc	12.8 oz.		378.5 L	100 gal.
3785.0 cc	1.0 gal.		3.75 KL	1,000 gal.
Treat Water by "dribbling" this product solution around the edges of the site to be treated. Where a spray rig is available – it may be used to apply the use solution – instead of the hand dribbling method.				

WEEKLY MAINTENANCE				
This product		Per	Unit water for 100 ppm*	
Metric	US		Metric	US
1.9 cc	0.065 oz.		3.78 L	1 gal.
19.0 cc	0.62 oz.		37.85 L	10 gal.
190 cc	6.2 oz.		378.5 L	100 gal.
1892.5 cc	62 oz.		3.75 KL	1,000 gal.
Each week repeat maintenance dosage or add this product using a test kit to maintain the proper concentration.				

* ppm - parts per million active ingredient.

OTHER USES

EVAPORATIVE COOLER (Not for use in CA): Controls algae, slime forming fungi on cooler pads and certain fungal plant pathogens that may carry over in the cooler as well as bacterial odors. Use 1 teaspoon of MONTEREY CONSAN 20 per 15 gallons of water every other week.

Note: Not all varieties or strains of plants listed have been tested. Unintentional consequences such as crop injury or ineffectiveness may result because of certain environmental or growing conditions, manner of use or applications, and insect resistance. Therefore, before treating a large number of plants, spray a few plants and observe for plant damage and for performance prior to full-scale application. All risks shall be assumed by the user.

ORCHIDS

Species	Diseases Controlled	Rate/Gal. of Water	Additional Information
Seedlings	Damping-off (caused by Pythium, Fusarium or Rhizopus)	0.2 oz.	From flask, soak seedling in solution for 2-5 minutes before potting to remove agar. Spray seedlings once a month as preventative maintenance. Spray any time disease is evident with solution.
Dividing plants		0.5 oz.	Remove compost then soak plant for 2-5 minutes in solution. Operate with wet hands and cutting tools (see Disinfecting). After division and root trim, soak divisions in solution for 2 to 5 minutes. Pot in compost drenched with solution.
Adult Plants		0.25 oz.	Spray any time disease, other than rot (see Adult Orchid Rots), is evident with solution. Spray once per month as preventative maintenance. Drench compost when fungus is evident.
	Botrytis petal blight	0.02 oz.	During cool weather, spray plants, flowers and under benches at first sign of spotting on flowers. Normally, weekly sprayings will greatly reduce or eliminate the botrytis spotting.
	Rots (such as <i>Erwinia</i> , <i>et al.</i>)	0.5 oz.	Remove compost from plant, then soak entire plant 10 minutes or more in solution. Cut away all rot with treated tool (see Disinfecting). Soak plant wounds again for 10 minutes in solution. Repot in drenched compost.
	Crown rot	0.25 oz.	Spray plants weekly with solution, assuring coverage around and in base of leaves.

HORTICULTURAL USAGES

Species	Diseases Controlled	Rate/Gal. of Water	Additional Information
Bulbs such as Daffodils, Easter lilies, Gladioli, Hyacinth, Iris, Narcissus, Tulips	Fusarium basal rot	(1 tbsp.)	Prestorage: Treat the corms and bulbs immediately after they are "cleaned" by soaking them for 10 minutes in the solution. Preplant: Treat again immediately before planting.
Cut Flowers in containers, tubes and decorative vases	Bacterial stem plugging (restricts water uptake) Fungi (can create ethylene gas, a case of petal drop)	0.13 oz.	Flower spikes should be hardened in solution during harvesting, storage and shipping to help control the transfer of diseases.
Plant Cuttings such as Begonias, Carnations, Chrysanthemums, Fuchsias, Geraniums)	Carnation rust, Crown gall, Gray mold leaf spots and Powdery mildew	0.25 oz.	Soak cuttings 2-5 minutes in the solution. Plant in compost drenched with the solution.
Seedlings	Damping off	0.25 oz.	Treat seeds and cuttings before planting by spraying with solution to completely wet all surfaces. Use inert material for planting seeds and cuttings, such as sphagnum moss, vermiculite, perlite, pumice, sterilized commercial mix rather than garden soil. If garden soil must be used, thoroughly saturate with the solution. After planting, water with solution (unless using soil saturated as above) and again after seedlings emerge. Spray weekly as needed.
	Downy mildew	0.25 oz.	Spray with solution at 5 day intervals as needed.

ORNAMENTAL USAGES

Species	Diseases Controlled	Rate/Gal. of Water	Additional Information
Aucuba, Dahlia, Fuchsia, Geranium, Gladiolus, Peony, Piersis	Botrytis	0.25 oz.	Spray buds and blooms lightly at first sign of disease, and repeat 2-3 times per week as needed. Before shipping, dip spikes in solution.
	Downy mildew, Gray mold	0.25 oz.	Spray at first sign of disease repeating at 7-14 day intervals. Before shipping, dip spikes in solution.
	Rust	0.25 oz.	Spray at 7-14 day intervals during the wet periods starting in early Summer. Before shipping, dip spikes in solution.
Cactus	Bacterial leaf spot, Root rot, Top rot	0.25 oz.	Sterilize seeds and disease free cuttings by dipping cuttings in the solution for 5 minutes before planting. Then spray at 10-14 day intervals.
	Slimy collar rot	0.25 oz.	Just before planting, dip entire plant in the solution.
Calendulas, China Asters, Chrysanthemum, Dahlias, Lilac, etc.	Powdery mildew	0.25 oz.	Spray upper and lower surface of leaves and branches with solution until thoroughly wet. Spray when buds burst in Spring and repeat in 2 weeks. Spray every 2 weeks during the mildew season.

ORNAMENTAL USAGES (Contd.)

Species	Diseases Controlled	Rate/Gal. of Water	Additional Information
Carnations	Canker rot, Carnation rust, Crown rot, Leaf spot, Root rots, Stem rot	0.25 oz.	Spray at first sign of disease and repeat at 7-10 day intervals.
	Gray mold rot	0.25 oz.	Spray 1-2 times just before bloom and again as flowers are opening.
Crassula, Kalanchoe, Sedum	Crown gall, Crown rot, Root rot	0.25 oz.	Destroy infected plant parts and drench the remainder. Spray plants weekly as needed.
Dogwood	Anthrachnose, Leaf blight, Leaf spots	0.25 oz.	Spray 4 times: when flower buds begin to open, as soon as flower bracts have fallen, 4 weeks later, and in late summer after flower buds are well formed. If possible, spray just before rainy period.
Fuchsia, Gardenia, Geranium, Heliotrope	Bud rot, Gray mold, Stem rot	0.25 oz.	Spray at first sign of disease and repeat weekly as needed.
Hawthorn	Rust	0.25 oz.	Begin protective spraying with new leaf formation after winter dormancy but before onset of rainy season. Thoroughly spray undersides of infected leaves.
Hawthorn (Crataegus)	Leaf blights	0.25 oz.	Make 3 applications: begin when leaves emerge from buds, when leaves are half grown, and again when fully grown.
Hollyhock	Rust	0.25 oz.	Apply at weekly intervals as soon as growth begins. Repeat applications in late Summer.
Iris	Botrytis, Crown rot, Gray mold rot	0.25 oz.	Treat all new rhizomes before planting. Submerge rhizomes and 3-4" of leaves in the solution for 30 minutes. Plant at once while still wet with solution.
	Leaf spots	0.25 oz.	Spray weekly with solution while leaves are growing.
Ornamental trees (such as Sycamore, Ash, Crape Myrtle, Live Oak, etc.)	Anthrachnose	0.25 oz.	Spray upper and under sides of leaves and twigs until thoroughly wet. A large tree requires 50-60 gallons of spray per treatment for adequate coverage. Make 3 applications: when buds start to swell in the spring, as leaves start to expand, and when leaves are about 3/4 grown.
	Powdery mildew	0.25 oz.	Spray upper and lower surface of leaves and branches with solution until thoroughly wet at bud break and again in April. Apply 50-60 gallons of spray per tree for adequate coverage.
Palms	Heart rot, Penicillium, leaf base rot	0.25 oz.	At first sign of rot, completely fill cup or "heart" formed by the palm branches at top center of the plant. Repeat every 7 days as needed.
Roses	Black spots, Powdery mildew, Rusts, Stem gall	0.25 oz.	When symptoms appear, spray upper and lower surface of leaves and branches with solution until thoroughly wet. Apply at 7-10 day intervals as needed.
	Brown canker	0.25 oz.	Begin spray applications in early spring when new growth starts and repeat weekly until plant is dormant.
Saintpaulias (African Violets)	Bud rot, Crown rot, Downy mildew, Gray mold rot Botrytis, Powdery mildew, Root rot	0.09 oz.	Dip entire plant and pot in the solution for 5 minutes. Thereafter, spray every 10 days as needed.
Snapdragons	Anthrachnose, Downy mildew, <i>Phyllosticta</i> blight, Powdery mildew, Stem rot, Wilt	0.25 oz.	When symptoms appear, spray upper and lower sides of leaves until thoroughly wet. Repeat every 7-10 days as needed. Collect and burn tops in the fall.
	Rust	0.25 oz.	Apply at 7-10 day intervals as soon as plants are set out. Collect and burn tops in the Fall.

WATER AND SMOKE DAMAGE RESTORATION

Effective against odors caused by smoke and water damage for home use. MONTEREY CONSAN 20 is particularly suitable for use in water damage restoration situations. Saturate affected materials with enough product to remain wet for at least 10 minutes. Use proper ventilation.

SPECIAL INSTRUCTIONS FOR CLEANING CARPETS: This product can be used in residential areas. For use on wet cleanable synthetic fibers. Do not use on wool. Vacuum carpet thoroughly prior to cleaning. Test fabric for color fastness.

For Portable Extraction Units: Mix 1/2 ounce of MONTEREY CONSAN 20 per two gallons of water (or equivalent use dilution).

For Rotary Floor Machines: Mix 1/2 ounce of MONTEREY CONSAN 20 per two gallons of water (or equivalent dilution) and apply at the rate of 300-500 sq. ft. per one gallon.

Do not mix MONTEREY CONSAN 20 with other cleaning products. Follow the cleaning procedures specified by the manufacturer of the cleaning equipment. After using MONTEREY CONSAN 20 set the carpet pile and protect the carpet from furniture legs and bases while drying. Do not over wet. If applied to stain resistant nylon carpet, apply a fabric protector according to the carpet manufacturer's directions.

GENERAL DEODORIZATION: *Use as a deodorizer is not allowed in CA.* To deodorize, add 1.5 ounces of MONTEREY CONSAN 20 to 5 gallons of water (or equivalent use dilution). Wipe up excess liquid.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

CONTAINER DISPOSAL/PESTICIDE STORAGE: Non-refillable container. Store only in original container. Do not reuse empty container. Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep this under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use. Triple rinse as follows: Fill container 1/4 full with water and reclose the container. Agitate vigorously and dispose of rinsate. Repeat two more times. Wrap container and put in trash or offer for recycling if available.

RESIDUE REMOVAL INSTRUCTIONS: Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Fill the container 1/4 full with water and recap. Shake for 10 seconds. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

BATCH NO. _