



Maxforce[®] FC MAGNUM

ROACH KILLER BAIT GEL

79432135
80215495P 200521AV1

FOR USE IN COMMERCIAL, INDUSTRIAL AND RESIDENTIAL AREAS

ACTIVE INGREDIENT: Fipronil 1, [5-Amino-1-(2, 6-dichloro-4-fluorophenyl)-4-(1, R, S)-
(-)-1-fluoro-2-propyl-1H-pyrazole-3-carbonitrile] 0.05%
OTHER INGREDIENTS: 99.95%
TOTAL: 100.00%
CAS NO. 120068-37-3
EPA Reg. No. 402-4-400

EPA Est. No. 071 906 -GA-003

**STOP - READ THE LABEL
BEFORE USE.
KEEP OUT OF REACH OF
CHILDREN.
CAUTION**

See Panel for First Aid Instructions and Label for Complete
Precautionary Statements and Directions for Use.

Product may stain fabrics and clothing.

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue to rinse. Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-354-7577 for emergency medical treatment information.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PHYSICAL AND CHEMICAL HAZARDS: Do not use this product in or on electrical equipment where a possibility of shock hazard exists.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its label. Use MAXFORCE[®] FC MAGNUM ROACH KILLER BAIT GEL as a spot or crack and crevice treatment for the interior and exterior (adjacent to homes & structures) control of cockroaches in residential areas and the non-food/non-feed areas of institutional, warehousing and commercial establishments, warehouses, restaurants, food processing plants, supermarkets, hospitals, nursing homes, hotels, hotels, apartment buildings, bakeries, beverage plants, breweries, bottling facilities, candy plants, canneries, cereal processing and manufacturing plants, dairies and dairy product processing plants, frozen food plants, schools, laboratories, computer facilities, sewers, "aircraft," boats, boatslips, buses, homes, houses, industrial buildings, office buildings, kennels,

kitchens, manufacturing facilities, spice plants, stores, wineries and similar structures, mass deans, meat processing and packaging plants, meat and vegetable canneries, pet shops, and zoos. Non-food/non-feed areas include bedrooms, living rooms, bathrooms, dining, music rooms, garbage rooms, lawns, porches, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling). This product may also be used in food/feed areas of food/feed handling establishments as outlined below.

USE RESTRICTIONS

- Keep gel away from foods and food contact or preparation surfaces.
- Do not apply to areas where food/feed, utensils, or processing surfaces may become contaminated.
- USE ONLY IN AREAS NOT EASILY ACCESSIBLE TO CHILDREN, PETS, AND OTHER NON-TARGET ORGANISMS.
- In pet shops and zoos place bait out of reach of pets and zoo animals.
- Do not apply bait to surfaces where the temperature exceeds 130°F as bait may run.
- Do not deposit gel into exposed surfaces.
- Do not apply bait gel to areas which have been recently sprayed with insecticide.
- Do not spray insecticide over gel as it may cause the bait to become repellent.
- Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food prep and prepared food held surfaces. Bait may be removed by washing.

USE DIRECTIONS

Place bait into cracks and crevices or openings leading into voids, or make spot applications near harborage, aggregation, foraging / feeding areas.

Locations for bait placement: between different elements of construction; the hollow spaces in floors, walls, ceilings, or in equipment legs and bases; cracks and crevices around pipes, doors or windows, refrigerators, stoves, water heaters, dishwashers, and other equipment; behind or under baseboards, cupboards, cabinets, countertops, sinks; and wherever cockroaches may be found hiding. Numerous smaller placements will provide better control than fewer larger spots, especially for German cockroach control.

Application rates for all cockroaches: Application rate is dependent on level of infestation and species to be controlled. Apply gel as round spots or thin beads into cracks and crevices or as a spot treatment.

- For use in **resistance programs** where regular (monthly, quarterly) applications are vital for continuous control of cockroach population, use 0.75 grams of bait per square yard.
- For **light to moderate infestations** (per retreatments), use up to 1.5 grams of bait per square yard.
- In **do not enter or heavy infestation**, use up to 3.0 grams of bait per square yard.

BAIT QUANTITY REFERENCE TABLE

Approximate weight per spot diameter	Approximate weight per bead length
0.1 gram = Pin size spot	1/3 gram = 1.5 inch bead
1.0 gram = Dime size spot	2.5 gram = 3.0 inch bead
1.5 gram = Quarter size spot	1.5 gram = 6 inch bead

continued on back