

LESCO®

LESCO®

#681924

ST. AUGUSTINE GRASS WEED AND FEED

WITH PENOXsulAM  Made in USA

Combinación para deshierbar y abono



ST. AUGUSTINE GRASS WEED AND FEED 17-0-7 WITH PENOXsulAM  Made in USA Combinación para deshierbar y abono

17-0-7

N

NITROGEN
For vigorous blade, leaf and stem growth and dark green color

NITRÓGENO
Para que el tallo y hojas tengan un crecimiento vigoroso y un color verde oscuro

P

PHOSPHATE
Contains no phosphorus

FÓSFORO
Contiene ningún fósforo

K

POTASH
Promotes strong roots and improves turf hardness

POTASA
Promueve raíces fuertes y mejora el vigor del césped



Controls Dollarweed and other tough weeds

Controla la Centella y otras malezas difíciles



Recommended for use on St. Augustine grass lawns

Se recomienda su uso en césped de San Agustín

ACTIVE INGREDIENTS


Penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy [1,2,4] triazolo [4,5-c]pyrimidin-2-yl)-6-(trifluoromethyl) benzenesulfonamide.....	0.03%
OTHER INGREDIENTS.....	99.97%
TOTAL.....	100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION | PRECAUCION

If you do not understand this label, find someone to explain it to you in detail. Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
Read the entire label before using this product. Use only according to label instructions.

ST. AUGUSTINE GRASS WEED AND FEED 17-0-7 WITH PENOXsulAM  Made in USA Combinación para deshierbar y abono



#681924



Covers up to **14,500** Sq. Ft.

#090999 NET. WT. 50 LB (22.7 KG)

LESCO®

ST. AUGUSTINE GRASS WEED AND FEED 17-0-7 WITH PENOXsulAM  Made in USA Combinación para deshierbar y abono

LESCO®

LESCO®

St. Augustinegrass Weed & Feed with Penoxsulam



17-0-7

HOW THIS PRODUCTS WORKS

LESCO® St. Augustine Weed & Feed with Penoxsulam herbicide provides post-emergence control of annual, biennial and perennial broadleaf weeds in established home lawns and other recreational turfgrass areas.

GUARANTEED ANALYSIS

Total Nitrogen (N).....	17.00%
17.00% Urea Nitrogen*	
Soluble Potash (K ₂ O).....	7.00%
Iron (Fe).....	2.00%
0.02% Water Soluble Iron	
Manganese (Mn).....	1.00%
0.01% Water Soluble Manganese	
Derived From: Iron Sulfate, Manganese Sulfate, Muriate of Potash, Polymer Coated Urea, Urea	
*4.59% Slowly Available Nitrogen from Polymer Coated Urea	F1560

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Application Techniques for Applying LESCO® St. Augustine Weed & Feed with Penoxsulam.

- When using a drop-type spreader, a splash board mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treating smaller areas, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended for hand held units, walk and turn the crank at a constant rate of speed.
- A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully, measure the amount of product needed to avoid over application.

HOW MUCH TO APPLY

Apply LESCO® St. Augustine Weed & Feed with Penoxsulam to established turfgrass. Possible retreatment in approximately 30 days may be necessary if turfgrass is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed specified rates because damage to turfgrass may occur. Do not make more than two applications per season. Note: When applying to warm season turfgrass under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended to reduce application rates by 50% during hot, dry conditions to avoid possible dehydration or browning of turfgrass.

WHEN TO APPLY

Best weed control results are obtained when LESCO® St. Augustine Weed & Feed with Penoxsulam is applied to small, actively growing weeds in the spring or early fall. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control. Broadleaf weed species germinate at different times. Florida Applications: It is recommended to follow the Florida Green Industries BMP's at http://www.dep.state.fl.us/water/nonpoint/BMP_Book_final.pdf.

HOW TO APPLY

Apply product using a drop or rotary-type spreader designed to apply granules. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. To optimize product performance mow the turfgrass 2 to 3 days before applying LESCO® St. Augustine Weed & Feed with Penoxsulam. If the turfgrass is under moisture stress, a thorough sprinkler irrigation of a least ½ inch may be needed at least one to two days before application. APPLY LESCO® St. Augustine Weed & Feed with Penoxsulam IN THE MORNING WHEN DEW IS PRESENT. IF NECESSARY, SPRINKLER IRRIGATE WITH LESS THAN ¼ INCH OF WATER JUST PRIOR TO APPLICATION TO MAXIMIZE PARTICLE ADHERENCE TO LEAF SURFACES. Avoid watering of treated area within 24 hours of application. Avoid making application if rainfall expected within 24 hours after treatment.

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, and shoes plus socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS: To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bottles or drainage systems.

**KEEP OUT OF REACH
OF CHILDREN
CAUTION/PRECAUCIÓN**

If you do not understand this label,
find someone to explain it to you in detail.
Si usted no entiende la etiqueta, busque a alguien
para que se la explique a usted en detalle.

EPA Reg. No. 62719-585-10404 | EPA Est. No. 59144-GA-1(A), 32802-MO-1(B)
EPA Est. No. 9611-PA-011(L), 9611-LO-01(D), 37361-CO-1(E) | EPA Est. No. 8375-N-1(A),
35497-OR-1(H), 49991-TX-2(Q) | EPA Est. No. 82757-OH-001(M), 82757-FL-001(S)
EPA Est. No. 82757-MA-002(H)
First letter in lot code indicates manufacturing site.

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label when calling a poison control center or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.

Spreader Settings as a Guide for Calibration: NOTE: These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by LESCO, Inc. or the spreader manufacturer.

ROTARY SPREADER	3.4 lb./1000 sq. ft.	4.6 lb./1000 sq. ft.
LESCO® Commercial Calibration Gauge (80 pound hopper and larger)	14	16
Scotts® Retail Rotary Spreaders	5	6
Scotts® EasyGreen®	25	26
Vigoro® 4300 Walk-Behind Rotary Spreader	3	4

AN EXPANDED SPREADER SETTING LISTING CAN BE FOUND AT: www.LESCO.com

GENERAL USE PRECAUTIONS AND RESTRICTIONS: DO NOT apply LESCO® St. Augustine Weed & Feed with Penoxsulam on dichondra, carpetgrass, or turfgrass containing desirable clovers. DO NOT apply LESCO® St. Augustine Weed & Feed with Penoxsulam directly to, or otherwise permit LESCO® St. Augustine Weed & Feed with Penoxsulam to come into contact with flowers, ornamental shrubs, or trees, or other desirable broadleaf plants, as serious plant injury may occur. DO NOT apply to centipede grass during the period of winter dormancy. Grass should be green and actively growing at the time of application. The active ingredient in this product may leave residues in treated grass clippings which could be harmful to some plants and trees. DO NOT collect treated grass clippings for use as mulch around plants. The applicator must notify in writing the appropriate residences/individuals of this precaution when this product is applied by a lawn service/commercial applicator. To minimize the potential for turfgrass injury, allow four weeks between applications. Avoid unnecessary disturbances, including watering and mowing, for approximately 24 hours after application. Apply LESCO® St. Augustine Weed & Feed with Penoxsulam to newly seeded or sprigged turfgrass only after the third mowing or after the turfgrass seedlings have developed an adequate secondary root system. Reseeding is not recommended for at least three to four weeks after application. Use LESCO® St. Augustine Weed & Feed with Penoxsulam on solid (monoculture) stands of perennial ryegrass or tall fescue at a maximum rate of 1.7 lb. per 1000 sq. ft. Some yellowing of these turfgrasses may occur within 2 weeks of application. Any discoloration is transitory and will dissipate under adequate growing conditions. Do not apply more than 4.6 lb. of LESCO® St. Augustine Weed & Feed with Penoxsulam per 1000 sq. ft. per application. Do not apply more than 6.9 lb. of LESCO® St. Augustine Weed & Feed with Penoxsulam per 1000 sq. ft. per year.

1. Treatment on Turfgrass Species Not Listed on the Label for LESCO® St. Augustine Weed & Feed with Penoxsulam. Users who wish to use LESCO® St. Augustine Weed & Feed with Penoxsulam on turfgrass species not listed on this label may determine the suitability for use by treating a small area at a specified rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the target turfgrass species is tolerant. Applications should not be made to turfgrass under stress. The user assumes responsibility for any plant damage or other liability resulting from use of LESCO® St. Augustine Weed & Feed with Penoxsulam on turfgrass species not listed on this label. LESCO® St. Augustine Weed & Feed with Penoxsulam is recommended for use on the following established turfgrass species: Established Warm Season Turfgrass (3) | Bermuda grass (common or hybrid) | Centipede grass (4) | Fescue, tall (growing in warm season areas) (2) | Kikuyugrass | St. Augustinegrass | Zoysiagrass
2. Use LESCO® St. Augustine Weed & Feed with Penoxsulam on these species at a maximum rate of 1.7 lb. per 1000 sq. ft. per application and only if some injury can be tolerated. A slight yellowing of turfgrass may occur within two weeks of application. The discoloration is temporary and is generally gone within four weeks of treatment under adequate growing conditions.
3. Use no more than 4.6 lb. of LESCO® St. Augustine Weed & Feed with Penoxsulam per 1000 sq. ft. per application.
4. Do not apply to centipede grass during the period of winter dormancy. Grass should be green and actively growing at the time of application.

Weeds Controlled and Suppressed: Generally, application rates at the lower end of the rate range will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands and/or larger weeds), the higher rate within the rate range or repeat applications will be needed. Weeds growing in the absence of competition from other vegetation generally require higher rates or repeat applications to obtain satisfactory control or suppression.

Weeds Controlled or Suppressed	Rate
chickweed, common chickweed, mouseear clover, hop clover, red clover, white dandelion, common dollarweed English lawn daisy false dandelion	Florida, betony ivy, ground Lespedeza, common Medic, black Parley-piert Plantain, broadleaf Plantain, buckhorn Virginia buttonweed ¹
	(3.4 - 4.6) lb./1000 sq.ft.
Kyllinga, annual and green (suppression only) ¹	Nutsedge, purple and yellow (suppression only) ¹

¹ Re-treatment may be necessary

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. **Pesticide Storage:** Store in a cool, dry place in original container. **Container Handling:** Non-Refillable container. Do not reuse or refill this container. If empty, Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

IMPORTANT: READ BEFORE USE

Terms and Conditions of Use: If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use: It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies: To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
 2. Replacement of amount of product used.
- To the extent permitted by law, Seller shall not be liable for losses or damages resulting from handling or use of this product, unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner. Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aspfco.org/metals.htm>. Information concerning the raw material composing this product can be obtained by writing to: LESCO, Inc Attn: RA Dept 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag. LESCO® and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. SCOTTS, SpeedyGreen® And EasyGreen® are registered trademarks of SCOTT Company; Vigoro is a registered trademark of the United Industries Corporation.

#090999

NET. WT. 50 LB (22.7 KG)

Covers up to
14,500
Sq. Ft.

FOR PRODUCT ASSISTANCE, CALL LESCO

1-800-321-5325

OR VISIT US AT WWW.LESCO.COM

FREIGHT SHIPPING BAG
Meeting Requirements of
APPLICABLE FREIGHT CLASSIFICATION
Guaranteed by
TRINITY PACKAGING CORPORATION

F1560 | Rev. Fall 2015
Manufactured for LESCO, Inc.
1385 East 36th St.
Cleveland, OH 44114-4114

18 X 29