

**GROUP 4 HERBICIDE**

# Change Up™

## Selective Herbicide

**FOR SELECTIVE BROADLEAF WEED CONTROL  
IN ORNAMENTAL LAWNS AND TURF GRASSES.**

**CONTAINS MCPA, FLUROXYPYR AND DICAMBA**

**CONTROLS:** Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

**ACTIVE INGREDIENTS:**

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	51.05%
1-Methylheptyl Ester of Fluroxypyr:[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic Acid, 1-methylheptyl ester**	6.00%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	4.17%

**OTHER INGREDIENTS** ..... 38.78%

**TOTAL:** ..... 100.00%

Isomer Specific Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	41.68%, 4.0 lbs./gal.
**[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic acid	4.17%, 0.4 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	4.17%, 0.4 lbs./gal.

**DO NOT SELL, DISTRIBUTE OR USE THE PRODUCT IN NASSAU AND SUFFOLK COUNTY IN NEW YORK**

### **KEEP OUT OF REACH OF CHILDREN WARNING / AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND  
ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire,  
or Exposure, Call CHEMTREC  
(800) 424-9300  
For Medical Emergencies Only,  
Call (877) 325-1840

EPA Reg. No. 228-445

Manufactured for  
Nufarm Americas Inc.  
11901 S. Austin Ave.,  
Alsip, IL 60803



# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING/AVISO

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing.

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistant selection chart.

### Mixers, loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants
- shoes plus socks
- chemical-resistant gloves when mixing, loading, or using any hand-held equipment
- protective eyewear (goggles, face shield, or safety glasses)

## User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## USER SAFETY RECOMMENDATIONS

### Users Should:

- Wash hands, face and arms with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## FIRST AID

<b>IF IN EYES</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

## HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

## ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish, aquatic invertebrates and aquatic plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS.

Do not apply this product in a way that will contact any person, or pet, either directly or through drift. Keep people and pets out of the area during application. Aerial application is prohibited.

This pesticide must be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposure. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Under some conditions this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing; these methods also reduce pesticide run-off. Where feasible, use application techniques such as T-banding and in-furrow techniques which incorporate the pesticide into the soil. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where run-off could occur will minimize water run-off is recommended.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore, the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls worn over short-sleeved shirt and short pants
- chemical-resistant footwear plus socks
- chemical-resistant gloves made of any waterproof material
- chemical-resistant headgear for overhead exposure, and
- protective eyewear

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For Turf use, the maximum number of broadcast applications per treatment site is 2 per year.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter or allow others to enter the treated area until sprays have dried.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees\* and Roughs); also for use on Sod Farms.

\*Excluding Bentgrass Tees.

### **SPRAY DRIFT MANAGEMENT**

Preventing spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2–10 mph at the application site.

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not use on Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment. Use the higher rate for hard-to-control weeds. Avoid broadcast applications when air temperature exceeds 90°F. When using in small, spot treatment applications in temperatures over 90°F, turf injury may occur.

When treating Carpetgrass and St. Augustinegrass, avoid broadcast applications when air temperature exceeds 80°F. When air temperature exceeds 80°F limit application to spot treatment only.

### **APPLICATION RESTRICTIONS**

- Do not apply more than 3 pints of this product per application (1.5 lbs MCPA ae, 0.15 lbs Fluroxypyr ae, 0.15 lbs Dicamba ae).
- Do not apply more than 2 applications of this product to the same treatment site per year.
- The minimum retreatment interval for this product is 21 days.
- Do not apply this product to improved varieties of St. Augustinegrass including Bitterblue, Floratam and Seville.
- Do not apply this product to lawn or turf during spring transition (green-up).
- Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species.
- Do not apply to newly seeded grasses until well established.
- Do not apply by air.

## ACTIVE INGREDIENT APPLICATION RESTRICTIONS

- Application of MCPA is limited to 3.0 lbs ae per acre per year, no more than 1.5 lbs ae per application, and no more than 2 applications per year.
- Application of Fluroxypyr is limited to 0.5 lbs ae per acre per year
- Application of Dicamba is limited to 2.0 lbs ae per acre per year, no more than 1.0 lbs ae per application, and no more than 2 applications per year.

The suitable use of this product on non-labeled turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

**For optimum results:** (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

## WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

Alder	Cockle	Hedge bindweed	Pigweed	Spiny cocklebur	Wild marigold
Annual yellow	Cocklebur	Hedge mustard	Pineywoods bedstraw	Spotted catsear	Wild onion
sweet clover	Coffeebean	Hemp	Plains coreopsis	Spotted knapweed	Wild parsnip
Artichoke	Coffeeweed	Henbit	Plantain	Spotted spurge	Wild radish
Aster	Common	Hoary cress	Poison hemlock	Spurge	Wild rape
Austrian fieldcress	chickweed	Hoary plantain	Poison ivy	Spurweed	Wild strawberry
Bedstraw	Common mullein	Hoary vervain	Poison oak	Stinging nettle	Wild sweet potato
Beggartick	Common	Honeysuckle	Pokeweed	Stinkweed	Wild vetch
Biden	sowthistle	Hop clover	Poorjoe	Stitchwort	Willow
Bindweed	Corn Chamomile	Horsenettle	Povertyweed	Strawberry clover	Witchweed
Bird vetch	Creeping jenny	Horsetail	Prickly lettuce	Sumac	Woolly morningglory
Bitterweed	Crimson clover	Indiana mallow	Prickly sida	Sunflower	Woodsorrel
Bitter wintercress	Croton	Ironweed	Primrose	Sweet clover	Woolly croton
Black-eyed Susan	Cudweed	Jewelweed	Prostrate knotweed	Tall nettle	Woolly plantain
Black medic	Curly dock	Jimsonweed	Prostrate pigweed	Tall vervain	Wormseed
Black mustard	Curly indigo	Kochia	Prostrate spurge	Tansy mustard	Yarrow
Black-seed plantain	Dandelion	Knawel	Prostrate vervain	Tansy ragwort	Yellow rocket
Blessed thistle	Dead nettle	Knotweed	Puncture vine	Tanweed	Yellowflower
Blue lettuce	Dock	Lambsquarter	Purslane	Tarweed	pepperweed
Blue vervain	Dogbane	Lespedeza	Ragweed	Thistle	and other broadleaf
Box elder	Dogfennel	Locoweed	Red clover	Tick trefoil	weeds.
Bracted plantain	Elderberry	Lupine	Redroot pigweed	Toadflax	
Brassbuttons	English daisy	Mallow	Red sorrel	Trailing crownvetch	
Bristly oxtongue	Fall dandelion	Marshelder	Redstem filaree	Tumble mustard	
Broadleaf dock	False dandelion	Matchweed	Rough cinquefoil	Tumble pigweed	
Broadleaf plantain	False flax	Mexicanweed	Rough fleabane	Tumbleweed	
Broomweed	False sunflower	Milk vetch	Roundleafed marigold	Velvet leaf	
Buckhorn	Fiddleneck	Milkweed bloodflower	Rush	Venice mallow	
Buckhorn plantain	Field bindweed	Mugwort	Russian pigweed	Veronica	
Bulbous buttercup	Field pansy	Morningglory	Russian thistle	Vervain	
Bull nettle	Fleabane (daisy)	Mousear chickweed	St. Johnswort	Vetch	
Bull thistle	Flixweed	Musk thistle	Scarlet pimpernel	Virginia buttonweed	
Burdock	Florida betony	Mustard	Scotch thistle	Virginia creeper	
Burning nettle	Florida pusley	Narrowleaf plantain	Sheep sorrel	Virginia pepperweed	
Bur ragweed	Frenchweed	Narrowleaf vetch	Shepherdspurse	Wavyleaf bullthistle	
Burweed	Galinsoga	Nettle	Slender plantain	Western clematis	
Buttercup	Garlic mustard	Nutgrass	Smallflower galinsoga	Western salsify	
Canada thistle	Goathead	Orange hawkweed	Smartweed	White clover	
Carolina geranium	Goatsbeard	Oxalis	Smooth dock	White mustard	
Carpetweed	Goldenrod	Oxeye daisy	Smooth pigweed	Wild mustard	
Catchweed bedstraw	Ground ivy	Parsley-piert	Sneezeweed	Wild aster	
Catsear	Gumweed	Parsnip	Southern wild rose	Wild buckwheat	
Catnip	Hairy bittercress	Pearlwort	Sowthistle	Wild carrot	
Chickweed	Hairy fleabane	Pennycress	Spanishneedle	Wild four-o'-clock	
Chicory	Hawkweed	Pennywort	Spatterdock	Wild garlic	
Cinquefoil	Healall	Peppergrass	Speedwell	Wild geranium	
Clover	Heartleaf drymary	Pepperweed	Spiny Amaranth	Wild lettuce	

## MIXING INSTRUCTIONS

Fill spray tank with water. Start agitation and slowly add the specified amount of concentrate. Maintain continuous agitation after mixing and during application. If this product is allowed to stand in tank for extended periods of time, some separation may occur. Reagitate before use.

## ORNAMENTAL LAWNS AND TURF

This product may be applied by broadcast or spot treatment application methods. Spot treatment is defined as a treatment area no greater than 1,000 ft<sup>2</sup> per acre.

### Rates for Ornamental Lawns and Turf

For Bluegrass, Ryegrass and Fescue, apply 2 to 3 pints per acre. For Creeping Bentgrass use 1.25 pints per acre. For Common and Hybrid Bermudagrass, Bahiagrass, Carpetgrass, Centipedegrass and Zoysiagrass apply 1.5 to 2.0 pints per acre.

Make broadcast applications of 1.25 to 3 pints per acre in 20 to 240 gallons of water per acre (or spot treatment application of 0.46 to 1.10 fl. oz. in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf grasses.

**Lower Volume Equipment:** Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

**NOTE:** For all grasses (1) avoid double coverage resulting from overlap unless temporary turf injury can be tolerated and (2) use reduced rates if grass is stressed from heat, drought, etc.

## SOD FARMS

This product is also intended for use on sod farms. This herbicide provides selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

### Timing

Apply this herbicide to broadleaf weeds that are actively growing. Follow-up applications may be required for dense infestations of perennial and biennial weeds.

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations. Apply this herbicide to grass seedlings after the second mowing. Also, applications to dormant Bermudagrass, dormant Zoysiagrass and dormant Bahiagrass are suggested.

### Rates for Sod Farms

For Bluegrass, Ryegrass and Fescue, apply 2 to 3 pints per acre. For Creeping Bentgrass use 1.25 pints per acre. For Common and Hybrid Bermudagrass, Bahiagrass, Carpetgrass, Centipedegrass and Zoysiagrass apply 1.5 to 2.0 pints per acre. Spray volume for Sod Farm application is 5 to 175 gallons per acre.

Generally, the lower application rates will provide satisfactory control of sensitive weeds. The higher application rates will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

## OTHER NON-CROP AREAS

**(fencerows, hedgerows, roadsides (including aprons and guard rails), ditches, rights-of-way, utility, pipelines, powerlines, railroads, airports, commercial plants, storage and lumber yards, barrier strips and firebreaks, equipment areas, nurseries and ornamental plantings, fuel tank farms, pumping stations and other industrial sites)**

For broadcast application to control broadleaf weeds, mix at a rate of 2 to 3 pints per acre. Spray volume for these applications is 50 to 300 gallons of water per acre. Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (Spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume.

For small (spot) applications with small tank sprayers, apply at the rate of 0.75 to 2.0 ounces of this product per gallon of water and spray to thoroughly wet all foliage.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Certain species are more resistant to this herbicide and may display regrowth. Mix thoroughly before spraying.

## COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.



1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product.)
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 25°F. Protect product from freezing. If allowed to freeze, remix well before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### **CONTAINER DISPOSAL AND HANDLING:**

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## **WARRANTY DISCLAIMER**

**The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.** UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

## **LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations of liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(RV110614)

Change Up is a Trademark of Nufarm Americas Inc.  
All other trademarks are the property of their respective owners.

SPECIMEN