



Product Bulletin

Aquasweep®

EPA Reg. No. 228-316

2(ee) Recommendation[†]

For Distribution and Use Only in the State of Missouri Control of Salt Cedar (*Tamarix* spp.)

[†]This recommendation is made as permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA.

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR THIS PRODUCT BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. "Label" as used in this supplemental labeling refers to the label booklet for this product and this supplemental.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING.

This label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS IN THE PRODUCT LABEL BOOKLET MUST BE FOLLOWED, INCLUDING STATEMENTS PERTAINING TO THE WORKER PROTECTION STANDARDS, ON THE EPA REGISTERED LABEL FOR AQUASWEEP (EPA REG. NO. 228-316).

Aquasweep may be used on pasture, rangeland, aquatic and other non-crop sites, including fencerows, rights-of-way, and roadsides for the control of Salt Cedar (*Tamarix* spp.). Use a higher rate in the rate range for dense stands or larger plants.

Optimum control can be achieved when applications are made between July and October (prior to frost). Complete control will likely require follow-up applications in the years following the initial application.

Apply at the specified rate (see below) plus a surfactant (typically 1/4 to 1% v/v), such as a non-ionic surfactant containing 80% or greater active ingredient. Use of this product without a surfactant will result in reduced herbicide performance. When applying as a foliar spray, the addition of bark oil at 2% will result in increased performance. Refer to the Mixing and Application Instructions sections of the product label and the surfactant manufacturer label for more information.

Method of Application	Application Rate	Spray Volume
Aerial	4 pints/acre	10 to 15 GPA
Handgun foliar plus bark spray	4 - 5.5 pints	Spray to wet
Ground High Volume Applications	1.5% Solution	High Volume Conventional Basal or Dormant Stem Applications Recommended

RESTRICTIONS

Do not apply more than 5-1/2 pints (2.0 lb of 2,4-D acid equivalent) per acre per application.
Limited to 1 application per growing season.

NOTICE

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

**Manufactured For Nufarm Americas Inc.
11901 S. Austin Ave. | Alsip, IL 60803**

Aquasweep is a registered trademark of Nufarm Americas Inc.

000228-00316.20141223.2(ee) Salt Cedar.Missouri