



Change Up™

SELECTIVE HERBICIDE

STRIKE OUT THE WEED CONTROL GAP FOR WINNING TURFGRASS.

Change Up™ premium selective herbicide combines MCPA, fluroxypyr and dicamba to deliver powerful weed control results. An excellent herbicide for warm- and cool-season turf, Change Up controls more than 200 broadleaf weeds and it starts working quickly to strike out weeds and help reduce callbacks. Plus, Change Up is ideal for any accounts that may be sensitive to the use of 2,4-D on the property.







COMPETITIVE ADVANTAGES

MAXIMUM POWER. CONVENIENT FORMULA.

- Transition zone convenience of spraying both warm- and cool-season turfgrass without having to change herbicides
- Offers the power of three proven herbicides in one convenient pre-mixed formulation
- Contains MCPA – ideal for accounts requesting a 2,4-D alternative
- Labeled for warm- and cool-season turfgrass including Bahiagrass, Creeping bentgrass, Bermudagrass, Kentucky bluegrass, Ryegrass, Fescue, Carpetgrass, Centipedegrass, Zoysiagrass and St. Augustinegrass
- Reduced callbacks following a quick visual response and excellent weed control
- Contains the pyridine herbicide fluroxypyr for unparalleled clover control
- Broad spectrum – excellent control of more than 200 broadleaf weeds including dandelion, plantain, oxalis, chickweed, thistles and other hard-to-control species
- Convenient to use – tank mix compatible with a number of liquid fertilizers and iron formulations

AT-A-GLANCE INFO

EPA REG. NO.	228-445
ACTIVE INGR.	MCPA (41.68%), fluroxypyr (4.17%), dicamba (4.17%)
FORMULATION	liquid
CHEM. FAMILY	phoxycarboxylic acid, pyridines and benzoic acids
HRAC NO.	4
MODE OF ACT.	auxin growth regulators
SIGNAL WORD	warning
RESTRICTED	no
PACKAGE SIZE	2.5 gal (2 per case), 30 gal

-  RESIDENTIAL
-  COMMERCIAL
-  TURF
-  GOLF TURF
-  POST-EMERGE
-  SOD FARMS

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

Celero
HERBICIDE

Escalade²

SureGuard^{SC}
HERBICIDE



Nufarm
Grow a better tomorrow



STRIKE OUT THE WEED CONTROL GAP FOR WINNING TURFGRASS.

KEY USES

Golf course turf (fairways, aprons, roughs and tees, excluding bentgrass tees)
Ornamental turf lawns (residential, institutional and industrial)
Parks, cemeteries and athletic fields
Roadsides, rights-of-way, and other similar non-crop areas
Sod farms

KEY WEEDS CONTROLLED

(See product label for complete list)

Alder
Annual yellow sweet clover
Bitter wintercress
Bracted plantain
Broadleaf plantain
Chickweed
Chicory
Dandelion
Dock
False flax
Florida betony
Ground ivy
Gumweed
Hairy bittercress
Hairy fleabane
Indiana mallow
Ironweed
Jewelweed
Jimsonweed
Kochia
Knawel
Knotweed
Lambsquarters
Mallow
Morningglory
Nettle
Oxeye daisy
Parsley-piert
Plantain
Prostrate spurge
Prostrate vervain
Ragweed
Red clover
Sweet clover
Thistle
Tumbleweed
Velvet leaf
Western salsify
White clovers

USE SUMMARY

ORNAMENTAL LAWNS, GOLF COURSE AND ATHLETIC FIELD TURF

For Kentucky bluegrass, Ryegrass and Fescue, apply 2 to 3 pts/A. For Creeping bentgrass use 1.25 pts/A. For Common and Hybrid Bermudagrass, Bahiagrass, Carpetgrass, Centipedegrass and Zoysiagrass apply 1.5 to 2 pts/A. For St. Augustinegrass apply 1 pt/A.

Do not apply more than 3 pts per acre per application. Do not apply more than two broadcast applications per year, excluding spot treatments, with a minimum retreatment interval of 21 days.

Maximum control of weeds will occur from spring or early fall applications when weeds are actively growing. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during and after the time of treatment. Use the higher rate for hard to control weeds.

For optimum results: (1) avoid applying during excessively hot or dry periods unless irrigation is used, (2) turf should not be mowed 1 to 2 days before and following application, (3) reseed no sooner than 3 to 4 weeks after application of this product.

Adding oil, wetting agent or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage.

Do not exceed specified dosages for any area. Be particularly careful within the drip line of trees or other ornamental species. Do not apply to newly seeded grasses until well established. Avoid broadcast applications when air temperature exceeds 90°F. When using in small, spot treatment applications in temperatures over 90°F, turf injury may occur.

When treating Carpetgrass and St. Augustinegrass, avoid broadcast applications when air temperature exceeds 80°F. When air temperature exceeds 80°F limit these applications to spot treatment only.

Note: For all grasses (1) avoid double coverage resulting from overlap unless temporary turf injury can be tolerated and (2) use reduced rates if grass is stressed from heat, drought, etc.

Lower volume applications: Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

SOD FARMS

For Bluegrass, Ryegrass and Fescue, apply 2 to 3 pts/A. For Creeping bentgrass use 1.25 pts/A. For Common and Hybrid Bermudagrass, Bahia and Zoysia apply 1.5 to 2 pts/A. For St. Augustinegrass apply 1 pt/A. Spray volume for sod farm application is 5 to 175 gals/A. For sod farm use, do not apply more than 3 pts per acre per application. Do not apply more than 2 broadcast applications per year, excluding spot treatments, with a minimum retreatment interval of 21 days.

See product label for complete directions for use.



nufarm.com/usturf

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/usturf to download a full label.

©2019 Nufarm. Important: Always read and follow label instructions. Change Up™ and Escalade® are trademarks of Nufarm. SureGuard® is a trademark of Valent U.S.A. LLC. Celero® is a trademark of Sumitomo Chemical Company, Ltd.



15-TO-0011-D