

RESTRICTED USE PESTICIDE

Due To Potential Human Health Risks of Concern (Cancer) From Chronic Dietary or Dermal Exposure When Used in Accordance with Label Directions.
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.



Nutrite Professional Turf Fertilizer with 1.0% Oxadiazon Herbicide

FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF MANAGEMENT & LANDSCAPE PERSONNEL. NOT FOR USE ON HOME LAWNS. FOR USE ON GOLF COURSES, NURSERIES, PARKS, SOD FARMS, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS.

A Turf Fertilizer plus pre-emergent weed control product. For use on varieties of Perennial Bluegrass, Bentgrass, Bermuda grass, Buffalo grass, Perennial Ryegrass, St. Augustine grass, Seashore Paspalum, Tall Fescue and Zoysiagrass. Also for use around Ornamental Shrubs, Vines, Trees, and Ground Covers, as listed on this label. Controls Crabgrass, Goosegrass, Poa annua, Annual Sedge and other weeds as listed, on Golf Courses, Parks and recreational and Ornamental turf.

0-0-7

Mini
SGN 145

GUARANTEED ANALYSIS

Soluble Potash (K₂O)..... 7.0%
Sulfur (S)..... 2.5%
 2.5% Combined Sulfur (S)

Derived from : Potassium sulfate

ACTIVE INGREDIENT:	
Oxadiazon*	1.0%
OTHER INGREDIENTS:	
	99.0%
TOTAL:	100.0%

CAS* [19666-30-9]

EPA REG. NO. 53883-180-90561
 EPA EST. NO. 87245-CAN-001 (C)
 EPA EST. NO. 089135-AZ-1 (A)
 EPA EST. NO. 93243-CT-1 (CT)
 EPA EST. NO. 90561-NC-1 (N)
 EPA EST. NO. 93243-TX-1 (TX)

First Letters of Lot Number indicate EPA Establishment where product was manufactured.

Product Code: P160367N

Lot Number XX-XXXX

Net Contents: 1000 lb (453.59 kg)

MANUFACTURED, GUARANTEED AND DISTRIBUTED BY:
 Ferti Technologies
 111 McNess Road, Statesville, NC 28677

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Wear protective eyewear (goggles, face shield or safety glasses). Causes skin irritation. Do not get in eyes, on skin or clothing. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid breathing dust, or contact with skin, eyes or clothing. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

Personal Protective Equipment (PPE): Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; waterproof gloves, chemical resistant footwear plus socks; protective eyewear, headgear for overhead exposure; chemical resistant apron when cleaning equipment. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations: Applicators should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF SWALLOWED:	Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
IF ON SKIN:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This pesticide is toxic to oysters and shrimp.

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water.

This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of oxadiazon from runoff water and sediment.

OXADIAZON	GROUP	14	HERBICIDE
------------------	--------------	-----------	------------------

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

DIRECTIONS FOR USE Restricted Use Pesticide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system. Apply only as specified on this label.

After application, user must thoroughly irrigate the site as soon as possible on the same day of application.

USE RESTRICTIONS FOR ALL USE SITES:

- The minimum retreatment interval (RTI) between applications is 60 days.
- Total applications of oxadiazon of all physical forms must not exceed 6.0 lbs a.i./A/year if this product is used on the same site with liquid formulations.

WEED RESISTANCE MANAGEMENT:

For resistance management, this product is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - (2) a spreading patch of non-controlled plants of a particular weed species;
 - (3) surviving plants mixed with controlled individuals of the same species.
 If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local sales rep.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry Interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls; waterproof gloves, chemical resistant footwear plus socks, protective eyewear, and chemical resistant headgear for overhead exposure.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, and nurseries. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and waterproof gloves when handling Laundry exposed clothing before reuse. Do not allow entry to areas treated until dusts have settled or if watering in is required, until treated area and grass is dry.

WEEDS CONTROLLED:

Annual Bluegrass	*Jungle Ricegrass
Annual Sedge	Lambsquarter
*Barnyardgrass	*Liverwort
Bittercress	Oxalis
Carpetweed	*Panama Paspalum
Carpetgrass	Pennsylvania Smartweed
Common Groundsel	Redroot Pigweed
Common Purslane	Sensitive plant
Crabgrass	Shepardspurse
*Evening Primrose	Sow Thistle
Fall Panicum	Speedwell-veronica
Field Sandbur	Spotted Catsear
Fireweed	Spurge; Garden,
Foxtail; Green, Bristly	Petty, *Prostrate
Florida Pusley	Stinging Nettle
*Galinsoga	*Swinecress
Golden Ragwort	Virginia Buttonweed
*Goosegrass	Yellow Woodsorrel.

*For these weeds use 3 lbs. a.i. per acre (equal to 300 lbs. of this product per acre or 6.9 lbs. of this product per 1,000 sq. ft.).

WHERE, HOW MUCH AND WHEN TO USE ON TURFGRASS: This product may be used at full rate or as a split application (see SPRING AND SPLIT APPLICATIONS ON GOLF COURSES section for instructions on how to apply the 3 lbs. or 4 lbs. a.i. per acre annual rate as a split application) in established Perennial Bluegrass, Bentgrass (see restrictions), Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue, and Zoysiagrass turf, at apartment/condominium complexes, corporate and public parks, golf courses, sod farms, athletic fields, playgrounds, cemeteries and other ornamental and recreational turfgrass areas. Not for use on home lawns. The turf should be free of leaves and thatch and in condition to allow the granules to reach the soil surface. Know the square footage of the area to be treated and uniformly apply with a granule spreader. Do not apply to turf when it is wet. When using at full rate do not overlap spreader applications, especially on St. Augustinegrass since discoloration "streaking" damage may temporarily occur from the "doubled-up rate." See section on this label entitled PRECAUTIONS IN USING ON TURFGRASS for the maximum rate for St. Augustinegrass and Bentgrass of 3 lbs. a.i. per acre which is equiv. to 6.9 lbs. of this product per 1,000 sq. ft. After applications, sprinkler irrigate the treated turf with up to a 1/4 inch of water to release the herbicide for maximum weed control and quicker response from the fertilizer nutrients. This product is a pre-emergence herbicide and must be applied prior to weed seed germination. It will not control growing weeds.

TURF USE RESTRICTIONS:

- A maximum single application rate of 3.0 lbs a.i./A, and a maximum annual application rate of 6.0 lbs a.i./A."
- For applications on non-agricultural turf (e.g., golf courses, parks, athletic fields) that follow a liquid application, this product must be applied 60 days or later after the liquid application. After applying this product, no liquid applications of oxadiazon are allowed within the same calendar year on the same area of turf.
- For applications to non-agricultural turf (e.g., golf courses, parks, athletic fields), this product may not be applied prior to any liquid applications of oxadiazon within the same calendar year on the same area of turf.
- On golf course turf only: In areas of heavy weed infestation only, the maximum single application rate is 4.0 pounds of active ingredient per acre, and the maximum annual application rate is 8.0 pounds of active ingredient per acre. No liquid applications may be applied on areas of turf treated at this maximum application rate in the same year.
- On golf course turf only: total applications of oxadiazon of all physical forms are limited to 30% of the total managed turf area of a golf course per year. Applications may be made to any turf surface type, including fairways, tees,

greens, roughs, non-playable turf surfaces, or other managed turf areas. This 30% limit may be applied to the same or different turf areas from year to year.

- On golf course turf and sod farms: a minimum 10-foot-wide untreated vegetative buffer must be maintained between the treated area and any surface water features.

PRECAUTIONS IN USING ON TURFGRASS: For St. Augustinegrass and Bentgrass turf, DO NOT exceed the rate of 3 lbs. a.i. per acre, equal to 300 lbs. of this product or 6.9 lbs. of this product per 1,000 sq. ft. DO NOT apply to **Bentgrass** turf mowed less than 3/8 inch. This product may cause discoloration of **Bentgrass, Bermudagrass** and **St. Augustinegrass** but this discoloration is temporary. Avoid contact with flowers and shrubs except as recommended elsewhere for ornamental plants on this label. DO NOT apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. DO NOT apply to **Dichondra** or **Centipedegrass**. DO NOT use on golf course putting greens or tees. NOTE: For newly seeded lawns DO NOT apply this product until after the second cutting. For best turf management, place new grass seed in the fall. Spring pre-emergence weed control can then be applied.

SPRING AND SPLIT APPLICATIONS ON GOLF COURSES: The 3 LBS. & 4 LBS. active ingredient, annual per acre rates may be applied as a split application. Weed control and better turf management may be programmed by splitting the annual per acre rate into "two" separate applications. For example, to accomplish this in region #5 (see map of US) apply 1.5 lbs a.i. (for a required 3 lbs. a.i./acre annual rate) or 2 lbs. a.i. (for a required 4 lbs. a.i./acre annual rate) as the first application 1 to 2 weeks before estimated weed seed germination or in late December or early January. Follow with the second "1/2 rate" in 60 days or the first of March. The timing for these applications should be closely programmed to the geographic region (see map of US) and adjusted to an "early" or "late" spring. Do not split the 2 lbs. a.i./acre rate. The 3 lbs. a.i./acre rate should not be split if applied to new turf areas first coming under turf management where weed pressure is expected to be heavy.

PRE-EMERGENCE WEED CONTROL (Refer to map of US)

Weeds Controlled	Pounds of product per acre	Pounds of product per 1,000 sq. ft.	Application Instructions
Crabgrass (<i>Digitaria spp.</i>) Goosegrass (<i>Elyusine indica</i>) Field Sandbur (<i>Cynchus insertus</i>) Annual Sedge (<i>Cyperus compressus</i>)	200 to 300 lbs. per acre (2 to 3 lbs. a.i. per acre)	4.6 to 6.9 lbs. per 1,000 sq. ft.	Apply in late winter or early spring prior to weed seed germination. See map. In areas 4 & 5 application made after April or May not be fully effective.
Annual Bluegrass (<i>Poa annua</i>)	200 to 300 lbs. per acre (2 to 3 lbs. a.i. per acre)	4.6 to 6.9 lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to weed seed germination.
Virginia Buttonweed (<i>Diodia</i>) Pigweed (<i>Amaranthus spp.</i>) Florida Pusley (<i>Richardia scabra</i>) Oxalis (<i>Oxalis spp.</i>) Stinging Nettle (<i>Urtica dioica</i>) Carpetgrass (<i>Axonopus officinis</i>)	300 lbs. per acre (3 lbs. a.i. per acre)	6.9 lbs. per 1,000 sq. ft.	Apply in early spring prior to weed seed germination.

FALL APPLICATION FOR SPRING WEED CONTROL IN COOL SEASON TURF

Weeds Controlled	Pounds of product per acre	Pounds of product per 1,000 sq. ft.	Application Instructions

Crabgrass, Goosegrass and other annual weeds as listed	300 lbs. per acre (3 lbs. a.i. per acre)	6.9 lbs. per 1,000 sq. ft.	For use in Bluegrass, Bentgrass, perennial Ryegrass and Tall Fescue. Make application in the fall after turf has become dormant. For late fall applications, seeding into treated areas should be delayed until 4 months after greenup.
--	---	----------------------------	---

ANNUAL BLUEGRASS CONTROL IN BERMUDAGRASS FAIRWAYS OVERSEEDED WITH PERENNIAL RYEGRASS ONLY

Weeds Controlled	Pounds of product per acre	Pounds of product per 1,000 sq. ft.	Application Instructions
Annual Bluegrass (<i>Poa annua spp.</i>)	200 lbs. per acre (2 lbs. a.i. per acre)	4.6 lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to <i>Poa annua spp.</i> seed germination. Any <i>Poa annua spp.</i> established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where <i>Poa trivialis</i> is present in the overseed mixture. Apply the full 2 lb a.i. rate in one application. Note: In many areas annual bluegrass exists as at least two biotypes or subspecies of <i>Poa annua</i> . The true annual biotype, " <i>Poa annua spp. annua</i> " which germinates from seed every year is controlled while the biotype " <i>Poa annua spp. raptans</i> " which behaves like a perennial and survives through the summer, as an established plant, cannot be controlled. Consult the extension service or University weed specialist in your area for more information concerning your locale.

ANNUAL WEED CONTROL DURING ESTABLISHMENT FROM "SPRIGS" OF BERMUDAGRASS, ZOYSIAGRASS, AND SEASHORE PASPALUM

Weeds Controlled	Pounds of product per acre	Pounds of product per 1,000 sq. ft.	Application Instructions
Crabgrass, Goosegrass, Annual Sedge and other annual weeds	200 to 300 lbs per acre (2 to 3 lbs. a.i. per acre)	4.6 to 6.9 lbs. per 1,000 sq. ft.	Refer to Map of U.S. on this label. For Bermudagrass make application immediately prior to, or, immediately after sprigging. For Seashore Paspalum and Zoysia make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable injury. Disturbance of the soil surface after application may disrupt the herbicide barrier and reduce effectiveness. For best results sprinkler irrigate the "sprigged" areas with up to 1/4 inch water after application.

WHERE, HOW MUCH, AND WHEN TO USE, AROUND ORNAMENTS Can be applied anytime of the year prior to weed seed germination to both newly transplanted and established actively growing or dormant ornamental plants where both weed control and fertilization is desired. Rainfall or overhead irrigation will improve weed control activity. Do not mix into soil. Disturbing or cultivating the soil after treatment may reduce weed control. Do not apply granules to wet foliage or under conditions in which granules will collect on leaves, if so, sprinkler-irrigate the plants to move granules off foliage and down to soil level. Do not apply in greenhouses as plant injury may likely result. Use the higher application rates for areas that have a history of heavy weed pressure. For best weed control around ornamentals first remove any existing weed growth and if possible rake, cultivate and smooth the soil surface by breaking-up clumps. Broadcast the granules evenly over the prepared soil by applying the 2 to 3 lbs a.i. per acre, equal to 200 to 300 lbs. of this product per acre or 4.6 to 6.9 lbs. per 1,000 sq. ft.

ORNAMENTAL USE RESTRICTIONS:

- A maximum single application rate of 3.0 lbs a.i./A, and a maximum annual application rate of 6.0 lbs a.i./A

CRABGRASS SEED GERMINATION DATES



- 1. Mid-north & northwestern After May 20-30
- 2. Upper/mid-states & northeast After May 10
- 3. Lower mid-states After April 20
- 4. Upper southern states After Mar 20
- 5. Warm seasons states Jan 1 to Mar 20

SPREADER SETTINGS FOR POUNDS OF PRODUCT PER ACRE: The following spreader settings are for new equipment. Used, or heavily worn equipment may require adjustments. For proper setting calibration refer to the spreader manufacturer's Maintenance Manual. Do not apply this product until equipment is properly calibrated to accurately apply the desired pounds of product per acre, as recommended on this label.

SPREADER SETTINGS	for 1.5 lb. a.i./acre	for 2 lb. a.i./acre	for 3 lb. a.i./acre
Spyker 24-22 Rotary	36	4	4½
Lesco Rotary	D/10	G/15	I/17
Prizelawn CB Rotary	E/2½	J/4	L/4½
EarthWay 2400 Rotary	12	14	15
Cyclone Pro	3½	4	4½
Scotts R8A	I	K	L
Lely Broadcast	3II	4II	5II
Bigfoot	E	J	L

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE AND SPILL PROCEDURES: Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Do not rinse unless required for recycling. If recycling is not available: then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **If partly filled:** Call your local solid waste agency for disposal instructions. Never pour unused product down any indoor or outdoor drain.

READ ENTIRE LABEL BEFORE USING

Read the entire Directions for Use, Conditions, Disclaimer of Warranties, and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties, and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or

the manner of use of application, all of which are beyond the control of Control Solutions. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Control Solutions is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Control Solutions, Inc. disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATION OF WARRANTIES: The exclusive remedy of the of user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Control Solutions option, the replacement of the product.

F002367

Rev 250211