


FRONT LABEL



CONSTRUCTION

BOUND BOOKLET

.375" 2.312" .375"
Book Placement



GARDEN TECH
Sevin
INSECT KILLER
CONCENTRATE
TRUSTED BRAND FOR OVER 50 YEARS

QUICK FACTS	
PRODUCT FACTS 	Kills by contact Visible results in minutes Allow product to dry completely before people and pets return to the lawn
WHERE TO USE 	Fruit and Vegetable Gardens, Ornamental Plants and Shrubs, Flowers and Ground Covers, Around Structures, Lawns
FRUITS & VEGETABLES 	Protects over 250 listed fruits and vegetables Tomato, Pepper, Squash, Cucumber, Broccoli, Kale, Lettuce, Blackberry, Carrot
KILLS 	Imported Cabbageworm, Cutworms, Armyworms, Flea Beetles, Stink Bugs, Whiteflies, Aphids, Japanese Beetles and other listed lawn & garden insects
HOW MUCH TO USE 	4 fl. oz. per gallon of water Treats 1,000 sq. ft. of lawn

**Open for Precautionary Statements,
First Aid and Complete Directions for Use**

Distributed by:
TechPac, L.L.C.
1000 Parkwood Circle, Suite 700
Atlanta, GA, 30339

EPA Reg. No. 279-3347-71004
EPA Est. 67572-GA-1,
67572-GA-2,
Subscript used is first letter of
run code on container.

If you have a medical
emergency, please call
1-(800) 420-9347



6 13499 01014 8
A080216RT

5.59"


3.062"

FOR VISUAL REFERENCE ONLY

BASE LABEL

BASE TEXT

(Underneath Booklet)



BOOKLET OUTSIDE




The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)




The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)



The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)




The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)




The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)



The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)

1.812"			1.812"																							
5.09"			5.59"																							
<table><tr><th>Plants</th><th>Pests</th><th>Use Before Harvest</th></tr><tr><td colspan="3">VEGETABLES</td></tr><tr><td>Brassica Vegetables including: • Broccoli; • Chinese Broccoli (gai lan, white flowering broccoli); • Brussels Sprouts; • Cauliflower; • Cavalo Broccoli; • Kohlrabi; • Cabbage; • Chinese Cabbage (napa); • Leafy Brassica Greens including: Broccoli Raab (rapini); • Chinese Cabbage (bok choy); • Chinese Mustard Cabbage (gai choy); • Collards; • Kale; • Mizuna; • Mustard Greens; • Mustard Spinach; • Rape Greens</td><td>Corn Earworm, Cucumber Beetles, Cutworm, Diamondback Moth, Flea Beetles, Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Southern Cabbageworm, Tobacco Budworm Alfalfa Looper, Armyworms, Cabbage Looper, Cabbage Webworm, Ground Beetles, Leafminers (adults), Lygus Bugs, Onion Thrips, Stinkbugs, Wireworm (adults), Aphids, and Whiteflies</td><td>Spray up to 1 day before harvesting</td></tr><tr><td>Root & Tuber Vegetables • Arracacha; • Arrowroot; • Artichoke; • Garden Beet; • Edible Burdock; • Edible Canna; • Carrot; • Turnip-rooted Chervil; • Chicory; • Chufa; • Dasheen (Taro); • Ginger; • Ginseng; • Horseradish; • Leren; • Turnip-Rooted Parsley; • Parsnip; • Potato; • Oriental Radish (Daikon); • Rutabaga; • Salsify (Oyster Plant);</td><td>Cutworms, Cabbage Looper, Cucumber Beetle, European Corn Borer, Flea beetles, Leafhoppers, Southern Corn Rootworm (adult), Vegetable Weevil, Whitefringed Beetle (adult), Armyworm, Beet Armyworm, Yellowstriped Cabbage Maggot, Colorado Potato Beetle, Grasshoppers, Imported Cabbageworm, Potato Leafhopper, Potato Psyllid, Potato Tubeworm, Tarnished Plant Bug</td><td>Spray up to 1 day before harvesting</td></tr></table>			Plants	Pests	Use Before Harvest	VEGETABLES			Brassica Vegetables including: • Broccoli; • Chinese Broccoli (gai lan, white flowering broccoli); • Brussels Sprouts; • Cauliflower; • Cavalo Broccoli; • Kohlrabi; • Cabbage; • Chinese Cabbage (napa); • Leafy Brassica Greens including: Broccoli Raab (rapini); • Chinese Cabbage (bok choy); • Chinese Mustard Cabbage (gai choy); • Collards; • Kale; • Mizuna; • Mustard Greens; • Mustard Spinach; • Rape Greens	Corn Earworm, Cucumber Beetles, Cutworm, Diamondback Moth, Flea Beetles, Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Southern Cabbageworm, Tobacco Budworm Alfalfa Looper, Armyworms, Cabbage Looper, Cabbage Webworm, Ground Beetles, Leafminers (adults), Lygus Bugs, Onion Thrips, Stinkbugs, Wireworm (adults), Aphids, and Whiteflies	Spray up to 1 day before harvesting	Root & Tuber Vegetables • Arracacha; • Arrowroot; • Artichoke; • Garden Beet; • Edible Burdock; • Edible Canna; • Carrot; • Turnip-rooted Chervil; • Chicory; • Chufa; • Dasheen (Taro); • Ginger; • Ginseng; • Horseradish; • Leren; • Turnip-Rooted Parsley; • Parsnip; • Potato; • Oriental Radish (Daikon); • Rutabaga; • Salsify (Oyster Plant);	Cutworms, Cabbage Looper, Cucumber Beetle, European Corn Borer, Flea beetles, Leafhoppers, Southern Corn Rootworm (adult), Vegetable Weevil, Whitefringed Beetle (adult), Armyworm, Beet Armyworm, Yellowstriped Cabbage Maggot, Colorado Potato Beetle, Grasshoppers, Imported Cabbageworm, Potato Leafhopper, Potato Psyllid, Potato Tubeworm, Tarnished Plant Bug	Spray up to 1 day before harvesting	<table><tr><th>Plants</th><th>Pests</th><th>Use Before Harvest</th></tr><tr><td>• Black Salsify; • Spanish Salsify; • Skirret; • Sweet Potato; • Tanier (Cocoyam); • Turmeric; • Turnip; • Yam Bean; • Yam (True)</td><td></td><td></td></tr><tr><td>Cucurbit Vegetables Group including but not limited to: • Chayote (fruit); • Chinese Waxgourd (Chinese Preserving Melon); • Citron Melon; • Cucumber; • Gherkin; • Gourd (edible) (including hyotan, cucuzza, hechima, Chinese orkra); • Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); • Muskmelon (hybrids and/or cultivars of Cucumis melo) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); • Pumpkin; • Summer Squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow,</td><td>Cutworms, Cabbage Looper, Cucumber Beetles. (adult), Leafhoppers, Melonworm, Pickleworm, Rindworm, Squash Bug, Squash Vine Borer, Corn Earworm, Plant Bugs, Stinkbugs</td><td>Spray up to 1 day before harvesting</td></tr></table>			Plants	Pests	Use Before Harvest	• Black Salsify; • Spanish Salsify; • Skirret; • Sweet Potato; • Tanier (Cocoyam); • Turmeric; • Turnip; • Yam Bean; • Yam (True)			Cucurbit Vegetables Group including but not limited to: • Chayote (fruit); • Chinese Waxgourd (Chinese Preserving Melon); • Citron Melon; • Cucumber; • Gherkin; • Gourd (edible) (including hyotan, cucuzza, hechima, Chinese orkra); • Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); • Muskmelon (hybrids and/or cultivars of Cucumis melo) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); • Pumpkin; • Summer Squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow,	Cutworms, Cabbage Looper, Cucumber Beetles. (adult), Leafhoppers, Melonworm, Pickleworm, Rindworm, Squash Bug, Squash Vine Borer, Corn Earworm, Plant Bugs, Stinkbugs	Spray up to 1 day before harvesting
Plants	Pests	Use Before Harvest																								
VEGETABLES																										
Brassica Vegetables including: • Broccoli; • Chinese Broccoli (gai lan, white flowering broccoli); • Brussels Sprouts; • Cauliflower; • Cavalo Broccoli; • Kohlrabi; • Cabbage; • Chinese Cabbage (napa); • Leafy Brassica Greens including: Broccoli Raab (rapini); • Chinese Cabbage (bok choy); • Chinese Mustard Cabbage (gai choy); • Collards; • Kale; • Mizuna; • Mustard Greens; • Mustard Spinach; • Rape Greens	Corn Earworm, Cucumber Beetles, Cutworm, Diamondback Moth, Flea Beetles, Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Southern Cabbageworm, Tobacco Budworm Alfalfa Looper, Armyworms, Cabbage Looper, Cabbage Webworm, Ground Beetles, Leafminers (adults), Lygus Bugs, Onion Thrips, Stinkbugs, Wireworm (adults), Aphids, and Whiteflies	Spray up to 1 day before harvesting																								
Root & Tuber Vegetables • Arracacha; • Arrowroot; • Artichoke; • Garden Beet; • Edible Burdock; • Edible Canna; • Carrot; • Turnip-rooted Chervil; • Chicory; • Chufa; • Dasheen (Taro); • Ginger; • Ginseng; • Horseradish; • Leren; • Turnip-Rooted Parsley; • Parsnip; • Potato; • Oriental Radish (Daikon); • Rutabaga; • Salsify (Oyster Plant);	Cutworms, Cabbage Looper, Cucumber Beetle, European Corn Borer, Flea beetles, Leafhoppers, Southern Corn Rootworm (adult), Vegetable Weevil, Whitefringed Beetle (adult), Armyworm, Beet Armyworm, Yellowstriped Cabbage Maggot, Colorado Potato Beetle, Grasshoppers, Imported Cabbageworm, Potato Leafhopper, Potato Psyllid, Potato Tubeworm, Tarnished Plant Bug	Spray up to 1 day before harvesting																								
Plants	Pests	Use Before Harvest																								
• Black Salsify; • Spanish Salsify; • Skirret; • Sweet Potato; • Tanier (Cocoyam); • Turmeric; • Turnip; • Yam Bean; • Yam (True)																										
Cucurbit Vegetables Group including but not limited to: • Chayote (fruit); • Chinese Waxgourd (Chinese Preserving Melon); • Citron Melon; • Cucumber; • Gherkin; • Gourd (edible) (including hyotan, cucuzza, hechima, Chinese orkra); • Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); • Muskmelon (hybrids and/or cultivars of Cucumis melo) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); • Pumpkin; • Summer Squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow,	Cutworms, Cabbage Looper, Cucumber Beetles. (adult), Leafhoppers, Melonworm, Pickleworm, Rindworm, Squash Bug, Squash Vine Borer, Corn Earworm, Plant Bugs, Stinkbugs	Spray up to 1 day before harvesting																								
14			15																							
2.312"			2.312"																							

The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)

Plants	Pests	Use Before Harvest
• Currant; • Elderberry; • Gooseberry; • Huckleberry		
Stone Fruit Trees • Apricot; • Cherry (Sweet and Tart); • Nectarine; • Peach; • Plum (including Chickasaw Plum, Damson Plum, and Japanese Plum); • Plumcot; • Prune	American Plum Borer, Black Cherry Aphid, Cherry Fruit Fly, Green Fruitworm, Leafrollers, Leafhoppers, Lesser Peach Tree Borer, Peach Tree Borer, Peach Twig Borer, Plum Curculio, Oriental Fruit Moth, Rose Chafer, Stink Bugs, Tarnished Plant Bug, Tufted Apple Budmoth, Western Cherry Fruit Fly	Spray up to 14 days before harvesting
Pome-Fruit Trees • Apple; • Crabapple; • Loquat; • Mayhaw; • Pear; • Oriental Pear; • Quince	Apple Maggot, Codling Moth, European Apple Sawfly, Green Fruitworm, Japanese Beetle, Lesser Appleworm, Oblique Banded Leafroller, Oriental Fruit Moth, Pandemis Leafroller, Pear Psylla, Plum Curculio, Potato Leafhopper, Redbanded Leafroller, Rosy Apple Aphid, Spirea Aphid, Spotted Tentiform Leafminer, Stink Bugs, Tarnished Plant Bug, Tufted Apple Bud Moth, Variegated Leafroller, White Apple Leafhopper	Spray up to 14 days before harvesting
Grapes	Eastern Grape Leafhopper, Variegated Leafhopper, Western Grape Leafhopper	Spray up to 1 day before harvesting
Peanuts	Cutworm spp., Green Cloverworm, Velvetbean Caterpillar, Red-necked Peanut Worm, Bean Leaf Beetle, Leafhopper spp., Southern Corn Rootworm (adult), Vegetable Weevil, Whitefringed Beetle (adult), Aphid spp., Armyworm, Beet, Fall Armyworm, Corn Earworm, Grasshopper spp., Lesser Cornstalk Borer, Soybean Looper, Stink Bug spp., Tobacco Thrips	Spray up to 7 days before harvesting

18

Plants	Pests	Use Before Harvest
Nut Trees • Almonds • Beech nut; • Brazil nut; • Butternut; • Cashew; • Chestnut; • Chinquapin; • Filbert (hazelnut); • Hickory nut; • Macadamia nut; • Pecan; • Walnut (Black and English)	Black Pecan Aphid, Codling Moth, Filbert Worm, Hickory Shuckworm, Leaffooted Bugs, Navel Orangeworm, Oblique-banded Leafroller, Peach Twig Borer, Pecan Leaf Casebearer, Pecan Casebearer, Pecan Phylloxera, Pecan Weevil, Plant Bugs, Stink Bugs, Walnut Aphid, Walnut Husk Fly, Yellow Pecan Aphid	Spray up to 7 days before harvesting Do not apply more than 0.125lb ai/A per season
Tropical Fruits • Avocado • Black Sapote • Canistel • Mamey Sapote • Mango • Papaya • Sapodilla • Star Apple	Avocado Lace Bug, Avocado Leafhopper, Avocado Leafroller, Avocado Looper, Avocado Tree Girdler, Avocado Whitefly, Brown Soft Scale, Caterpillars, Mirids, Omnivorous Loopers, Orange Tortrix, Scale Crawlers, Spanworm, Thrips, Twig Borers	Spray up to 1 day before harvesting

19

NOTES

5.09"

5.59"

2.312"

2.312"


The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)




The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)




The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)



The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)




The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)




The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)




The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)




The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)



The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)

1.812"			1.812"																																
5.09"			5.59"																																
<table><tr><th>Plantas</th><th>Plagas</th><th>Uso antes de la cosecha</th></tr><tr><td>Vegetales de bulbos (especies de Allium) incluyendo: - Ajo; ajo elefante - Chalota verde o echalote verde - Cebolleta o cebollita - Puerro o ajo porro - Cebolla verde o cebollino, cebolla de bulbo seco - Cebolleta o cebolla larga - Chalotas de bulbo seco y verde - Cebolla de verdeo, cebollino o cebolleta </td><td>Gusano cogollero, gusano cortador, minador de hojas (adultos), gusano de la cebolla (adultos), chinche hedionda, áfido y trip de la cebolla</td><td>Rocielo hasta 7 días antes de la cosecha.</td></tr><tr><td>Girasol</td><td>Gusano cortador, gusano blanco y cascarudo o saltaperico</td><td>Rocielo hasta 30 días antes de la cosecha.</td></tr><tr><td>Maíz dulce</td><td>Chinche, gusano de la raíz del maíz (adulto), mosca del maíz (Euxesta stigmatias), gusano cortador, alticino, saltahoja, escarabajo japonés (adulto), escarabajo nitidulido (adultos), chinche de las plantas opaca, gusano cogollero, bórer del maíz, gusano cogollero del maíz, saltamonte y áfido</td><td>Rocielo hasta 3 días antes de la cosecha.</td></tr><tr><td>Legumbres — jugosas y secas Guisantes de vainas comestibles jugosos, guisantes sin cáscara jugosos y guisantes sin cáscara secos (variedades de Pisum) incluyendo: - Guisante enano - Guisante de vaina comestible - Guisante mollar - Tirabeque dulce - Gandul - Petipuas - Arveja - Chicharo verde - Lenteja </td><td>Gusano cortador, oruga de la dama pintada, gusano peludo, Silverspotted Skipper (Epargyreus clarus), oruga de la alfalfa, gusano cogollero, gusano cogollero del sur, escarabajo de la hoja del frijol de franjas amarillas, escarabajo de la papa de Colorado, bórer del maíz europeo, bórer del maíz sureño, escarabajo de la raíz del maíz (adulto), Curculio del frijol, escarabajo del pepino, alticino, gusano verde, escarabajo de tierra, gusano de la col importado, escarabajo japonés, devorador de</td><td>Rocielo hasta 1 día antes de la cosecha para los guisantes o frijoles jugosos sin cáscara o en vainas y agregue 21 días antes de la cosecha para los guisantes o frijoles sin cáscara secos</td></tr></table>			Plantas	Plagas	Uso antes de la cosecha	Vegetales de bulbos (especies de Allium) incluyendo: - Ajo; ajo elefante - Chalota verde o echalote verde - Cebolleta o cebollita - Puerro o ajo porro - Cebolla verde o cebollino, cebolla de bulbo seco - Cebolleta o cebolla larga - Chalotas de bulbo seco y verde - Cebolla de verdeo, cebollino o cebolleta	Gusano cogollero, gusano cortador, minador de hojas (adultos), gusano de la cebolla (adultos), chinche hedionda, áfido y trip de la cebolla	Rocielo hasta 7 días antes de la cosecha.	Girasol	Gusano cortador, gusano blanco y cascarudo o saltaperico	Rocielo hasta 30 días antes de la cosecha.	Maíz dulce	Chinche, gusano de la raíz del maíz (adulto), mosca del maíz (Euxesta stigmatias), gusano cortador, alticino, saltahoja, escarabajo japonés (adulto), escarabajo nitidulido (adultos), chinche de las plantas opaca, gusano cogollero, bórer del maíz, gusano cogollero del maíz, saltamonte y áfido	Rocielo hasta 3 días antes de la cosecha.	Legumbres — jugosas y secas Guisantes de vainas comestibles jugosos, guisantes sin cáscara jugosos y guisantes sin cáscara secos (variedades de Pisum) incluyendo: - Guisante enano - Guisante de vaina comestible - Guisante mollar - Tirabeque dulce - Gandul - Petipuas - Arveja - Chicharo verde - Lenteja	Gusano cortador, oruga de la dama pintada, gusano peludo, Silverspotted Skipper (Epargyreus clarus), oruga de la alfalfa, gusano cogollero, gusano cogollero del sur, escarabajo de la hoja del frijol de franjas amarillas, escarabajo de la papa de Colorado, bórer del maíz europeo, bórer del maíz sureño, escarabajo de la raíz del maíz (adulto), Curculio del frijol, escarabajo del pepino, alticino, gusano verde, escarabajo de tierra, gusano de la col importado, escarabajo japonés, devorador de	Rocielo hasta 1 día antes de la cosecha para los guisantes o frijoles jugosos sin cáscara o en vainas y agregue 21 días antes de la cosecha para los guisantes o frijoles sin cáscara secos	<table><tr><th>Plantas</th><th>Plagas</th><th>Uso antes de la cosecha</th></tr><tr><td>Frijoles de vainas comestibles jugosos, frijoles sin cáscara jugosos y frijoles sin cáscara secos incluyendo: - Ayocote - Habichuela verde - Alubia - Frijol espárrago - Frijol largo chino - Moth Bean (Vigna aconitifolia) - Frijol serpiente - Frijol machete - Frijol de soya (semilla inmadura) - Haba espada - Haba de Lima - Haba - Frijol de carita - Carilla - Lupino - Lupino dulce - Lupino blanco - Lupino blanco dulce - Haba - Frijol colorado - Judía - Frijol pinto - Frijol tépari - Frijol adzuki - Catjang (Vigna unguiculata cylindrica) - Frijol Castilla - Alubia Crowder - Moth Bean (Vigna aconitifolia) - Judía mungo o frijol chino - Rice Bean (Vigna umbellata) - Frijol negro o lenteja negra - Garbanzo - Guar - Frijol de Egipto </td><td>hojas, saltahoja, minador de las hojas (adulto), escarabajo del frijol mexicano, gorgojo del guisante, gorgojo de las hojas del guisante, polilla de las plantas, saltahoja de la papa, escarabajo del grano de maíz, gusano del grano del maíz (adulto), cercopídeo, Three-Cornered Alfalfa Hopper (Spissistilus festinus), gusano de la yema del tabaco, oruga de las leguminosas, gusano tejedor y oruga de la polilla tigre o polilla lanuda</td><td></td></tr><tr><td colspan="3">FRUTAS Y NUECES</td></tr><tr><td>Grupo de las bayas incluyendo: - Mora - Mora roja - Frambuesa roja y negra - Arándano azul (arbusto alto y arbusto bajo) </td><td>Enrollador de hoja, gusano totrix del naranjo, gorgojo de la raíz</td><td>Rocielo hasta 1 día antes de la cosecha.</td></tr></table>			Plantas	Plagas	Uso antes de la cosecha	Frijoles de vainas comestibles jugosos, frijoles sin cáscara jugosos y frijoles sin cáscara secos incluyendo: - Ayocote - Habichuela verde - Alubia - Frijol espárrago - Frijol largo chino - Moth Bean (Vigna aconitifolia) - Frijol serpiente - Frijol machete - Frijol de soya (semilla inmadura) - Haba espada - Haba de Lima - Haba - Frijol de carita - Carilla - Lupino - Lupino dulce - Lupino blanco - Lupino blanco dulce - Haba - Frijol colorado - Judía - Frijol pinto - Frijol tépari - Frijol adzuki - Catjang (Vigna unguiculata cylindrica) - Frijol Castilla - Alubia Crowder - Moth Bean (Vigna aconitifolia) - Judía mungo o frijol chino - Rice Bean (Vigna umbellata) - Frijol negro o lenteja negra - Garbanzo - Guar - Frijol de Egipto	hojas, saltahoja, minador de las hojas (adulto), escarabajo del frijol mexicano, gorgojo del guisante, gorgojo de las hojas del guisante, polilla de las plantas, saltahoja de la papa, escarabajo del grano de maíz, gusano del grano del maíz (adulto), cercopídeo, Three-Cornered Alfalfa Hopper (Spissistilus festinus), gusano de la yema del tabaco, oruga de las leguminosas, gusano tejedor y oruga de la polilla tigre o polilla lanuda		FRUTAS Y NUECES			Grupo de las bayas incluyendo: - Mora - Mora roja - Frambuesa roja y negra - Arándano azul (arbusto alto y arbusto bajo)	Enrollador de hoja, gusano totrix del naranjo, gorgojo de la raíz	Rocielo hasta 1 día antes de la cosecha.	2.312"		
Plantas	Plagas	Uso antes de la cosecha																																	
Vegetales de bulbos (especies de Allium) incluyendo: - Ajo; ajo elefante - Chalota verde o echalote verde - Cebolleta o cebollita - Puerro o ajo porro - Cebolla verde o cebollino, cebolla de bulbo seco - Cebolleta o cebolla larga - Chalotas de bulbo seco y verde - Cebolla de verdeo, cebollino o cebolleta	Gusano cogollero, gusano cortador, minador de hojas (adultos), gusano de la cebolla (adultos), chinche hedionda, áfido y trip de la cebolla	Rocielo hasta 7 días antes de la cosecha.																																	
Girasol	Gusano cortador, gusano blanco y cascarudo o saltaperico	Rocielo hasta 30 días antes de la cosecha.																																	
Maíz dulce	Chinche, gusano de la raíz del maíz (adulto), mosca del maíz (Euxesta stigmatias), gusano cortador, alticino, saltahoja, escarabajo japonés (adulto), escarabajo nitidulido (adultos), chinche de las plantas opaca, gusano cogollero, bórer del maíz, gusano cogollero del maíz, saltamonte y áfido	Rocielo hasta 3 días antes de la cosecha.																																	
Legumbres — jugosas y secas Guisantes de vainas comestibles jugosos, guisantes sin cáscara jugosos y guisantes sin cáscara secos (variedades de Pisum) incluyendo: - Guisante enano - Guisante de vaina comestible - Guisante mollar - Tirabeque dulce - Gandul - Petipuas - Arveja - Chicharo verde - Lenteja	Gusano cortador, oruga de la dama pintada, gusano peludo, Silverspotted Skipper (Epargyreus clarus), oruga de la alfalfa, gusano cogollero, gusano cogollero del sur, escarabajo de la hoja del frijol de franjas amarillas, escarabajo de la papa de Colorado, bórer del maíz europeo, bórer del maíz sureño, escarabajo de la raíz del maíz (adulto), Curculio del frijol, escarabajo del pepino, alticino, gusano verde, escarabajo de tierra, gusano de la col importado, escarabajo japonés, devorador de	Rocielo hasta 1 día antes de la cosecha para los guisantes o frijoles jugosos sin cáscara o en vainas y agregue 21 días antes de la cosecha para los guisantes o frijoles sin cáscara secos																																	
Plantas	Plagas	Uso antes de la cosecha																																	
Frijoles de vainas comestibles jugosos, frijoles sin cáscara jugosos y frijoles sin cáscara secos incluyendo: - Ayocote - Habichuela verde - Alubia - Frijol espárrago - Frijol largo chino - Moth Bean (Vigna aconitifolia) - Frijol serpiente - Frijol machete - Frijol de soya (semilla inmadura) - Haba espada - Haba de Lima - Haba - Frijol de carita - Carilla - Lupino - Lupino dulce - Lupino blanco - Lupino blanco dulce - Haba - Frijol colorado - Judía - Frijol pinto - Frijol tépari - Frijol adzuki - Catjang (Vigna unguiculata cylindrica) - Frijol Castilla - Alubia Crowder - Moth Bean (Vigna aconitifolia) - Judía mungo o frijol chino - Rice Bean (Vigna umbellata) - Frijol negro o lenteja negra - Garbanzo - Guar - Frijol de Egipto	hojas, saltahoja, minador de las hojas (adulto), escarabajo del frijol mexicano, gorgojo del guisante, gorgojo de las hojas del guisante, polilla de las plantas, saltahoja de la papa, escarabajo del grano de maíz, gusano del grano del maíz (adulto), cercopídeo, Three-Cornered Alfalfa Hopper (Spissistilus festinus), gusano de la yema del tabaco, oruga de las leguminosas, gusano tejedor y oruga de la polilla tigre o polilla lanuda																																		
FRUTAS Y NUECES																																			
Grupo de las bayas incluyendo: - Mora - Mora roja - Frambuesa roja y negra - Arándano azul (arbusto alto y arbusto bajo)	Enrollador de hoja, gusano totrix del naranjo, gorgojo de la raíz	Rocielo hasta 1 día antes de la cosecha.																																	


The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

INSIDE TEXT (Odd Pages)



The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)

BOOKLET INSIDE

INSIDE TEXT (Even Pages)

1.812"

INSIDE TEXT (Odd Pages)

1.812"

The total number of pages must be divisible by four (4).

The maximum number of pages is forty (40)