

BAC-PACK™

Beneficial Microbial Blend

INGREDIENTS:

CONTAINS NON-PLANT FOOD INGREDIENTS

Natural, beneficial, rhizobacteria*.....0.6%
* $\geq 3.00 \times 10^8$ CFUs per ml. Includes *Bacillus* and *Pseudomonas* spp.
Also contains natural fermentates, and inert and carrier ingredients.....99.4%
Microorganisms in Bac-Pack are not genetically modified.

Target Crops:

Turfgrass
Purpose Statement: This mix of microbiological organisms is intended to produce beneficial physical, chemical, biochemical, and biological changes in the soil. The microbial components transform organic matter into nutrients and humus to reduce the need for supplementary inputs.

Bac-Pack is a trademark of Soil Technologies Corp.
Copyright 2019



**NET CONTENT:
1 GALLON**

NOTICE: Manufacturer and distributor disclaim all crop liabilities for product users when product is used in a manner inconsistent with labeling.

Manufactured By:



SOILTECH
Teaming With Biology

Soil Technologies Corp.

2103 185th Street, Fairfield, IA 52556, U.S.A.

TEL: +1-641-472-3963

FAX: +1-641-472-6189

www.soiltechcorp.com

KEEP OUT OF REACH OF CHILDREN

Storage: Bac-Pack should be stored in a cool, dark place. Refrigeration will extend the shelf life of the product.

Disposal: Do not contaminate water, food or feed by storage or disposal.

Disposal of Container: Triple rinse, discard in trash or recycle with plastics.



BAC-PACK™

Beneficial Microbial Blend

SHAKE WELL PRIOR TO USE

Application Site: Plant root zone.

Dosage: 1-2 pints per acre (1.2-2.4 lt/ha) at 10-14 day intervals during the growing season.

Application Methods: Bac-Pack is to be mixed with water or liquid fertilizer and spray applied or injected through an irrigation system. The live micro organisms in Bac-Pack should not be mixed with chlorinated water. A dechlorinator may be used. The product should be watered in immediately after application.

Application Timing: Applications should be made when soil temperatures are between 50-90°F (10-32°C). Excessive exposure to ultraviolet light (sunlight) can interfere with the activity of Bac-Pack. Allow 48 hours between Bac-Pack applications and the use of herbicides, fungicides and pesticides.

Initial Dosage For Stressed Turf: To promote greater physical, biochemical, and biological change in the soil, Bac-Pack can initially be applied at higher rates, up to 8.0 pints per acre (9.6 lt/ha). Apply as a drench application. For subsequent applications, revert to the usual application rate.

Expires:

