

GROUP 11 P1 FUNGICIDE

PULL HERE TO OPEN ►



syngenta

Fungicide

Broad-spectrum, systemic fungicide for the prevention and control of turf diseases on golf courses, sod farms, institutional, commercial and industrial buildings, and collegiate and professional athletic fields.

1 pound
Net Weight



Active Ingredients:

Azoxystrobin: methyl (E)-2-[2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate*	50.00%
Acibenzolar-S-methyl: Benzo(1,2,3)thiadiazole-7-carbothioic acid-S-methyl ester*	1.18%

Other Ingredients: 48.82%

Total: 100.00%

Heritage Action contains 0.50 lb of azoxystrobin and 0.0118 lb of acibenzolar-S-methyl per pound of product.

Heritage Action is a water-dispersible granule (WG) formulation.

*IUPAC

KEEP OUT OF REACH OF CHILDREN. CAUTION

See additional precautionary statements and directions for use inside booklet.

Reformulation is prohibited. See individual container labels for repackaging limitations.

EPA Reg No. 100-1550 EPA Est. 67545-AZ-1

Product of United Kingdom

Formulated in USA

SCP 1550A-L1A 0315

4049601

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15–20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15–20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear protective eyewear, long sleeved shirt and long pants, socks and shoes. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Protective eyewear
- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber.
- Shoes plus socks

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

continued...

PRECAUTIONARY STATEMENTS *(continued)*

Environmental Hazards

This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Do not apply directly to water except as specified on this label.

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to run-off of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via run-off for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features, such as ponds, streams and springs, will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from run-off water and sediment. Run-off of this product also will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify state and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

Physical or Chemical Hazards

Do not use with or store near any oxidizing agents. Hazardous chemical reactions may occur.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF

THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USES

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

continued...

AGRICULTURAL USE REQUIREMENTS (*continued*)

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

NON-AGRICULTURAL USES

Broad-spectrum, systemic fungicide for the prevention and control of turf diseases on golf courses, sod farms, institutional, commercial and industrial buildings, and collegiate and professional athletic fields.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area being treated must be vacated by unprotected persons.

Do not treat areas while unprotected humans or domestic animals are present in the treatment areas. Because certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information.

Do not allow entry into treatment area until area that was treated with Heritage Action is dry.

PRODUCT INFORMATION

Heritage Action couples a broad spectrum, systemic fungicide with a systemic compound used for control and suppression of disease through induction of host plant resistance. Heritage (azoxystrobin) is a broad-spectrum, systemic fungicide that prevents and controls pathogens causing foliar, stem, and root diseases, including leaf and stem blights, leaf spots, patch diseases, mildew, molds and rusts of turfgrass. Action (acibenzolar) exhibits an additional unique mode of action, beyond Heritage alone, which mimics the natural systemic activated resistance (SAR) response found in most plant species. This SAR response does not involve direct activity against the target pathogens.

Heritage Action fungicide may be used to control these diseases on golf courses, sod farms, institutional, public, commercial and industrial buildings, and collegiate and professional athletic fields. Heritage Action may be applied as a foliar spray in alternating spray programs or in tank mixes with other registered turf-protection products.

USE PRECAUTIONS AND RESTRICTIONS

Adjuvants: When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program is recommended.

DO NOT apply more than 10 lb product/acre/year (3.7 oz product/1,000 sq ft per year).

DO NOT apply to residential lawns.

DO NOT apply by air.

DO NOT graze or feed clippings from treated turf areas to animals.

SPRAY DRIFT MANAGEMENT

DO NOT spray when conditions favor drift beyond area intended for application.

DO NOT apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Conditions that contribute to drift include thermal inversion, wind speed

and direction, sprayer nozzle/ pressure combinations, and spray droplet size. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

PHYTOTOXICITY

Heritage Action is extremely phytotoxic to certain apple and crabapple varieties. Extreme care must be used to prevent injury to apple and crabapple trees. Even trace amounts can cause unacceptable phytotoxicity. **DO NOT** spray Heritage Action where spray drift may reach apple or crabapple trees.

Heritage Action has demonstrated some phytotoxic effects when mixed with products that are formulated as EC's. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.

INTEGRATED PEST (DISEASE) MANAGEMENT (IPM)

Sound turf management resulting in healthy, vigorous turf is the foundation of a good IPM program. Cultural practices such as proper choice of turf variety, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management should be integrated with the use of fungicides to increase turf vigor and reduce the susceptibility to disease. Immunoassay detection kits and extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

RESISTANCE MANAGEMENT

Heritage Action exhibits two modes of action. Plant pathogens are known to develop tolerance to host plant resistance and to products used repeatedly for control. The use of Heritage Action should conform to sensitivity management strategies established for

crops and use areas due to its two modes of activity Heritage Action should be applied in a tank mix or alternation program with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed. Since Heritage Action contains a strobilurin fungicide, avoid alternation with other strobilurins. Do not apply more than two sequential Heritage Action applications for Gray leaf spot and *Pythium* spp. control. For all other diseases when Gray leaf spot and *Pythium* spp. are not present, do not apply more than three sequential applications of Heritage Action.

MIXING INSTRUCTIONS

Thoroughly clean spray equipment before using this product to prevent possible crop injury or nozzle clogging problems from spray tank contamination. To prepare spray solution, partially fill the spray tank with clean water and begin agitation. Add the specified amount of Heritage Action to the tank allowing adequate time for good dispersion and then add an adjuvant if recommended. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray solution stand overnight in the spray tank. Do not prepare more spray solution than is needed for the immediate spray operation. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

Heritage Action is compatible with many commonly used fungicides, liquid fertilizers, herbicides insecticides and biological control products. If tank mixes are desired, observe all directions, precautions, and limitations on labeling of all products used. Consult compatibility charts or your local or state agricultural or turf authorities for compatibility information. Do not combine Heritage Action in the spray tank with any product unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective and non-phytotoxic under your conditions of use. If physical compatibility is unknown, the following testing procedure should be followed: pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand for at least 20 minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

Products should be added to the spray tank in the following order: Heritage Action, other WG or dry flowable formulations, wettable powders and flowable (aqueous suspensions) products. Finish filling the tank to the desired volume to obtain the proper spray concentration. Maintain agitation throughout the spraying operation. Do not allow spray mixture to stand overnight or for prolonged periods. Make up only the amount of spray required for immediate use. Sprayers should be thoroughly cleaned immediately after application.

APPLICATION INSTRUCTIONS

Heritage Action should be applied prior to disease development. Heritage Action mimics the SAR response in plants. Maximum disease control is normally obtained 4 days after a Heritage Action application. Apply Heritage Action at rates and timings given on this label. Use the shortest specified application interval and/or the highest specified rate under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions conducive to disease exist. Repeat applications at specified intervals for as long as required. Mix Heritage Action with the required amount of water and apply as a dilute spray application in 2-4 gallons of water per 1,000 square feet (87-174 gallons per acre). Proper adjustments and calibration of spraying equipment is essential for best disease control. For spot treatments, use 0.2 oz Heritage Action per 1 to 2 gallons of water.

For use with soil injection applications:

Heritage Action may be applied through a liquid fungicide injector for the control of ectrotrophic root diseases such as summer patch and take-all patch. Use Heritage Action only in liquid injection equipment specifically designated for pesticide use.

Apply Heritage Action at 0.2 to 0.4 oz per 1,000 sq ft. Spray carrier volume should fall within 30-150 gallons of water per 1,000 sq ft. Injection hole spacing of 1 inch by 1 inch is recommended for optimum control. Injection depth should be no greater than 2 inches. One inch depth is recommended for optimum results. Application timing should follow disease control strategies used for normal broadcast spray programs.

For use in the establishment of turfgrass from seed or in overseeding of dormant turfgrass:

Heritage Action may be used for control of certain turfgrass diseases associated with turfgrass establishment from seed. Heritage Action may also be used during overseeding of dormant turfgrass.

Heritage Action may be safely applied before or after seeding or at seedling germination and emergence to ryegrass, bentgrass, bluegrass, and fescue turfgrass types. Optimum application timing is during seeding.

TABLE 1: Use Rates and Timings for Listed Diseases

Target Diseases ¹	Use Rate (oz product per 1,000 sq ft)	Application Interval (days)	Remarks
Anthracnose (<i>Colletotrichum graminicola</i>)	0.2-0.4	14-28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Bermudagrass Decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	0.4	28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Brown Patch (<i>Rhizoctonia solani</i>)	0.2-0.4	14-28	Apply when conditions are favorable for disease development.
Brown Ring Patch (<i>Waitea circinata</i> var. <i>circinata</i>)	0.2-0.4	14-28	Apply when conditions are favorable for disease development.

Target Diseases ¹	Use Rate (oz product per 1,000 sq ft)	Application Interval (days)	Remarks
Yellow Patch (<i>Rhizoctonia cerealis</i>)	0.2-0.4	14-28	Make one or two applications in fall or when conditions are favorable for disease development.
Fairy Ring (<i>Lycoperdo</i> , <i>Arachnion</i> , <i>Bovista</i> , <i>Vascellum</i> , and <i>Agrocybe</i> spp.)	0.4	28	Apply preventatively or as soon as possible after fairy ring symptoms develop. Apply only in 4 gallons water per 1,000 square feet (174 gallons /acre). Add the recommended rate of a wetting agent to the final spray. Fairy ring symptoms may take 2 to 3 weeks to disappear following curative applications, and reapplication may be required in some cases. Severely damaged or thin turf may require reseeding.
Fusarium Patch (<i>Microdochium nivale</i>)	0.2-0.4	14-28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.2-0.4	14-28	Begin applications before disease is present and continue applications while conditions are favorable for disease development.

Target Diseases ¹	Use Rate (oz product per 1,000 sq ft)	Application Interval (days)	Remarks
Gray Snow Mold Typhula blight (<i>Typhula incarnata</i> , <i>T. ishikariensis</i>)	0.4	10–28	Make two applications of 0.4 oz spaced 10–28 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide, such as Daconil®, may enhance control under severe disease pressure.
Leaf Rust Stem Rust Stripe Rust (<i>Puccinia</i> spp.)	0.2–0.4	14–28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Leaf spot (<i>Bipolaris</i> spp.)	0.2–0.4	14–21	Apply when conditions are favorable for disease development.
Melting Out (<i>Drechslera poae</i>)	0.2–0.4	14–21	Apply when conditions are favorable for disease development.
Necrotic Ring Spot (<i>Leptosphaeria korrae</i>)	0.2–0.4	14–28	Apply when conditions are favorable for disease development.
Pink Patch (<i>Limonomyces roseipellis</i>)	0.2–0.4	14–28	Apply when conditions are favorable for disease development.

Target Diseases ¹	Use Rate (oz product per 1,000 sq ft)	Application Interval (days)	Remarks
Pink Snow Mold (<i>Microdochium nivale</i>)	0.4	10–28	Make two applications of 0.4 oz spaced 10–28 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide, such as Daconil, may enhance control under severe disease pressure.
Powdery Mildew (<i>Erysiphe graminis</i>)	0.2–0.4	14–28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Pythium Blight Pythium Root Rot (<i>Pythium</i> spp.)	0.2–0.4	10–14	Use preventatively. Begin applications before disease is present. During periods of prolonged favorable conditions, treat on the 10 day application interval. For use on newly seeded as well as established turf.
Pythium Root Dysfunction (<i>Pythium volutum</i>)	0.4	21–28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.

continued...

Target Diseases ¹	Use Rate (oz product per 1,000 sq ft)	Application Interval (days)	Remarks
Red Thread (<i>Laetisaria fuciformis</i>)	0.2–0.4	14–28	Apply when conditions are favorable for disease development.
Rhizoctonia Large Patch/Zoysia Patch (<i>Rhizoctonia solani</i>)	0.2–0.4	14–28	Make one or two applications in fall or when conditions are favorable for disease development. Spring applications may also be required in some locations or when disease pressure is high.
Leaf and Sheath Spot (<i>Rhizoctonia zeae</i>)	0.2–0.4	14–28	Apply when disease conditions are favorable for disease development.
Southern Blight (<i>Sclerotium rolfsii</i>)	0.2–0.4	14–28	Apply when conditions are favorable for disease development.
Spring Dead Spot (<i>Ophiospharella korrae</i> , <i>O. herpotricha</i> , and <i>O. narmari</i>)	0.2–0.4	14–28	Apply 1 or 2 applications approximately one month prior to bermudagrass dormancy. 1/8–1/4 inch of irrigation directly after application is recommended. Reapply 14–28 days later.
Summer Patch (<i>Magnaporthe poae</i>)	0.2–0.4	14–28	Apply when conditions are favorable for disease development.

Target Diseases ¹	Use Rate (oz product per 1,000 sq ft)	Application Interval (days)	Remarks
Take-all patch (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>)	0.2–0.4	28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development. Make two applications, 28 days apart in the spring and two applications 28 days apart in the fall.
Dollar Spot Suppression (<i>Sclerotinia homoeocarpa</i>)	0.2–0.4	14–28	Heritage Action will suppress dollar spot. For effective control always mix Heritage Action with Daconil, Secure® or another Dollar Spot control fungicide
Bacterial Wilt Suppression (<i>Acidovorax</i> spp.) (<i>Xanthomonas</i> spp.)	0.2–0.4	14–28	For suppression of bacterial wilt on bent and Poa greens apply Heritage Action in a seasonal program with Daconil Action.

¹Do not apply more than two sequential applications of Heritage Action for control of Gray leaf spot and *Pythium* spp. For all other diseases when Gray leaf spot and *Pythium* spp. are not present, do not apply more than three sequential applications of Heritage Action.

TABLE 2: Heritage Action Rate Conversion Chart

Ounces Product Per 1,000 Sq Ft	Ounces AI Per 1,000 Sq Ft	Ounces Product Per Acre	Pounds Product Per Acre
0.20	0.10	8.7	0.5
0.30	0.15	13.1	0.8
0.40	0.20	17.4	1.1

TABLE 3: Amount of Heritage Action to Mix 100 Gallons

Heritage Action Use Rate	Spray Volume (gallons/1,000 sq ft)		
	2 gallons	3 gallons	4 gallons
0.2 oz	10 oz	6.7 oz	5 oz
0.4 oz	20 oz	13.3 oz	10 oz

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, sweep and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Handling [less than or equal to 50 pounds]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [bags]

Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

Heritage®, Daconil®, Secure®, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company.



©2015 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1550A-L1A 0315
4049601

GROUP 11 P1 FUNGICIDE

 **Heritage Action**

 syngenta

Fungicide

Broad-spectrum, systemic fungicide for the prevention and control of turf diseases on golf courses, sod farms, institutional, commercial and industrial buildings, and collegiate and professional athletic fields.

1 pound

Net Weight



GROUP 11 P1 FUNGICIDE

 **Heritage Action**

Broad-spectrum, systemic fungicide for the prevention and control of turf diseases on golf courses, sod farms, institutional, commercial and industrial buildings, and collegiate and professional athletic fields.

Active Ingredients:	
Azoxystrobin: methyl (E)-2-[2-[[6-(2-cyanophenoxy)pyrimidin-4-yl]oxy]phenyl]-3-methoxyacrylate*	50.00%
Acibenzolar-S-methyl: Benzo[1,2,3]thiadiazole-7-carbothioic acid-S-methyl ester*	1.18%
Other Ingredients:	48.82%
Total:	100.00%

Heritage Action contains 0.50 lb of azoxystrobin and 0.0118 lb of acibenzolar-S-methyl per pound of product.

Heritage Action is a water-dispersible granule (WG) formulation.
*IUPAC

Reformulation is prohibited. See individual container labels for repackaging limitations.

See additional precautionary statements and directions for use inside booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg No. 100-1550
EPA Est. 67545-AZ-1

KEEP OUT OF REACH OF CHILDREN. CAUTION

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

HOT LINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), call **1-800-888-8372**.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear protective eyewear, long-sleeved shirt and long pants, socks and shoes. Remove and wash contaminated clothing before reuse.

Environmental Hazards: This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Do not apply directly to water except as specified on this label.

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This product may impact surface water quality due to run-off of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via run-off for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is

applied and surface water features, such as ponds, streams and springs, will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from run-off water and sediment. Run-off of this product also will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Notify state and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

Physical or Chemical Hazards: Do not use with or store near any oxidizing agents. Hazardous chemical reactions may occur.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. **Pesticide Storage:** Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, sweep and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Handling: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities. **CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

Heritage® and the SYNGENTA Logo are trademarks of a Syngenta Group Company. ©2015 Syngenta

Manufactured for: Syngenta Crop Protection, LLC
P.O. Box 18300 Greensboro, North Carolina 27419-8300

SCP 1550A-11A 0315 4049601

 syngenta®



(011) 0 07 02941 56376 0