



Restriction: Not for sale, sale into, distribution and/or use in Nassau, Suffolk, Kings and Queens Counties in the state of New York.

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	0.2%
Other Ingredients	99.8%
Total	100.0%

10-00-00

Total Nitrogen (N)	10.00%
10.000% Urea Nitrogen	

DERIVED FROM: Urea.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE BACK OF LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS
FOR USE

FIRST AID	
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 8:00 AM-12:00 PM PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300."	
Note To Physician: No specific antidote is available. Treat the patient symptomatically.	

Mears Fertilizer, Inc. F1958
629 North Industrial Road
El Dorado, KS 67042 NET WEIGHT: 50 Lbs. (22.68 Kgs.)

NET WEIGHT: 50 Lbs. (22.68 Kgs.)

USER SAFETY RECOMMENDATIONS	
Users Should	* Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. * Remove clothing immediately if pesticides get inside. The wash thoroughly and put on clean clothing. * Remove PPE Immediately after handling this product. Wash the outsides of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.


- * **DO NOT** apply this product near fish pools, ponds, streams or lakes.
- * **DO NOT** use more than 200 pounds of this product (0.4 lbs a.i.) per acre per year or 4.5 pounds of this product (0.09 lbs a.i.) per 1,000 square feet/year.
- * **DO NOT** apply to vegetable gardens.
- * **DO NOT** apply this product to areas that are waterlogged or saturated with water. This product may not work effectively in these types of conditions.
- * **DO NOT** harvest or consume fruit or nuts from trees that have been treated within 1 year (365 days).
- * Keep children and pets off treated area until it has dried.
- * **DO NOT** use food utensils such as teaspoons or measuring cups for food purposes after use with this product.
- * **DO NOT** apply this product, by any application method, to linden, basswood or other *Tilia* species in the State of Oregon.
- * Not for sale, sale into, distribution and/or use in Nassau, Suffolk, Kings and Queens Counties in the state of New York.

* Coveralls * Chemical-resistant gloves made of any waterproof material * Shoes plus socks

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, greenhouses and sod farms

APPLICATION RATES FOR USE ON SOD FARMS and TURFGRASSES			
CROP SITE	PEST		DOSAGE
Sod Farms and Turfgrasses	<u>Larvae of:</u> Annual bluegrass weevil European Crane Fly Asiatic garden beetle Northern masked chafer Billbugs Oriental beetle Black turfgrass atenius <i>Phyllophaga</i> spp. European chafer Southern masked chafer		125 to 200 pounds per acre or 3.0 to 4.5 pounds per 1,000 square feet
	<u>Larvae of:</u> Asiatic garden beetle Northern masked chafer Black turfgrass atenius Oriental beetle European chafer <i>Phyllophaga</i> spp. Japanese beetle Southern masked chafer		150 pounds per acre or 3.5 pounds per 1,000 square feet
	<u>Larvae of:</u> Annual bluegrass weevil Billbugs		150 to 200 pounds per acre or 3.5 to 4.5 pounds per 1,000 square feet
	For optimum control, make application prior to egg hatch of the target pest. READ "APPLICATION EQUIPMENT" section of this label.		
	Mole crickets		200 pounds per acre or 4.5 pounds per 1,000 square feet
	For control of mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, accompany application by a remedial insecticide. Consult your local State Agricultural Experiment Station or State Extension Turf Specialists for more specific information regarding timing of application.		
	RESTRICTIONS: Irrigation or rainfall is necessary within 24 hours after application to move the active ingredient through the thatch. On golf courses, irrigate treated areas following application.		
	* DO NOT apply more than 200 pounds (0.4 pound of active ingredient) per acre per year.		
	* DO NOT mow turf or lawn area until after irrigation or rainfall has occurred so that uniformity of application is not affected.		
	* DO NOT graze treated areas or use clipping from treated areas for feed or storage		
* DO NOT apply to areas which are water logged or saturated, which will not allow penetration into root zone of the plant.			

Sample Label



SPREADER SETTING FOR THIS PRODUCT:

Use the following chart as a guide to establish an accurate rate of this product. Use a spreader setting that provides an application rate of 3.5 pounds per 1,000 square feet or 150 pounds per acre. Application rates will vary dependent upon speed, equipment wear and condition of spreader. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

APPLICATION RATES FOR USE ON ORNAMENTALS AROUND RESIDENTIAL, COMMERCIAL & INDUSTRIAL AREAS		
SITE	PEST	DOSAGE
Landscape Plantings:	White grub larvae	
Including: Trees, Shrubs, Evergreens, Flowers, Foliage Plants, Ground Covers	Such As: Japanese beetle larvae, Chafers, Phyllophaga spp. Asiatic garden beetle, Oriental beetle	3.5 to 4.5 pounds per 1,000 square feet
Apply as a broadcast treatment. Prior to application, remove black plastic or other weed barrier material that will not permit water permeability. Mechanical incorporation, irrigation or rainfall is necessary within 24 hours after application to move the active ingredient into the soil.		
Flower Beds	Adelgids Aphids Armored scales (suppression) Japanese beetle (adults) Lacebugs Leaf beetles (including Elm, Viburnum, and Lily)	4 ounces per 12 square feet (3ft. x 4 ft. bed area)
Roses	Leafhoppers (including Glassy-winged sharpshooter) Leafminers (including Birch and Boxwood) Mealybugs Pine tip moth (larvae) Psyllids	2 ounces per plant
Shrubs	Root weevil larvae (including Black vine) Sawflies (larvae) Soft Scales (including Lobate lac) Thrips Whiteflies	4 ounces per foot of plant height
For new plants, apply to soil and mix prior to planting. For established plants, use before pest damage occurs or immediately seen. Repeat applications every 8 weeks during the growing season, not to exceed max rate per year. Apply the required amount of granules evenly on top of soil; mix thoroughly into the top soil layer, being careful not to damage the upper roots, then water in thoroughly. For flower beds, sprinkle granules evenly over bed. RESTRICTIONS DO NOT apply near fish pools, ponds, lakes or streams. DO NOT allow runoff or puddling or irrigation water following application. DO NOT apply this product to areas that are water-logged or saturated with water. This product may not work effectively in these types of conditions. DO NOT harvest or consume fruit or nuts from trees that have been treated within 1 year (365 days). Keep children and pets off treated area until it has dried. DO NOT apply to vegetable gardens. DO NOT use utensils such as teaspoons or measuring cups for food handling after they have been used to measure this product. DO NOT apply more than 200 pounds (0.4 pound of active ingredient) per acre per year or 4.5 pounds/1000 square feet/year. DO NOT apply this product, by any application method, to linden, basswood or other Tilia species in the State of Oregon. Not for sale, sale into, distribution and/or use in Nassau, Suffolk, Kings and Queens counties in the state of New York.		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Carefully handle the container to prevent spills. If material is spilled, carefully contain any material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides below.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill, by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Refer to Precautionary Statements on label for hazards associated with the handling of this material. Contact CHEMTREC at 800-424-9300 for decontamination procedures or any other assistance that may be necessary.

IMPORTANT READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions disclaimer of warranties and limitations of liability.

WARRANTY STATEMENT

LIMITATION OF WARRANTY AND LIABILITY

Read the entire direction for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following

CONDITIONS, DISCLAIMER OF WARRANTIES, and LIMITATIONS OF LIABILITY.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Mears Fertilizer, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Mears Fertilizer, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Mears Fertilizer, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Mears Fertilizer, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Mears Fertilizer, Inc. election, the replacement of product.

In the state of Florida, it is recommended that the user of this product follow the Green Industries and Golf Course Best Management Practices found at: http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/10BMP_Book_Final.pdf and <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>.

If this product is being sold in a State requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available on the internet at: <http://aapfco.org/metals.htm>

In Virginia, this lawn/turf fertilizer contains phosphorous and is intended only for nonagricultural use on (1) turf during its first growing season, (2) on turf areas being repaired or renovated, and (3) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not intended for the routine maintenance of turf.

This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the specified application rates established by the State; or 3) Reestablishing or repairing a turf area.