

Manzate[®] Pro-Stick T&O[™]

FUNGICIDE

Small, uniform particles within the formulation break up faster to create a homogenous mixture that remains in suspension longer than larger particle sized formulations.

Manzate[®] Max T&O

FUNGICIDE

Now in a new easy-to-use flowable formulation with the reliable disease control growers expect.

For Superior Control of Key Diseases in Turfgrass, Sod Farms, and Ornamentals, Manzate T&O Fungicide provides multi-site, broad-spectrum disease control.



- Extended Residual Activity
- Multi-Site Action Allows Better Management of Fungicide Resistance
- Broad-Spectrum Fungicidal Coverage
- Economical Control Provides Better Returns

Turfgrass & Sodfarms Diseases Controlled		
Algae	Gray Leaf Spot	Red Thread
Brown Patch	Leaf Rust	Rhizoctonia Solani
Copper Spot	Leaf Spot	Slime Molds
Dollar Spot	Pink Snow Mold	Stem Rust
Fusarium Blight	Pythium Blight	Stripe Rust

Formulation	Use Rates*	Application Timing
Manzate Pro-Stick T&O	4 oz. -8 oz. in 3-5 gal of water per 1,000 sq. ft.	Typically begin application when disease threatens, or at the first sign of disease. Repeat application as per labeled interval rates as conditions persist.
Manzate Max T&O T&O	5 fl. oz. -14 fl. oz. per 1,000 sq ft.	
*Follow label instructions for disease specific use rates and application timing.		

Resistance Management Tool: Managing fungicide resistance is an important decision when choosing your disease control products. The multi-site action of Manzate T&O Fungicide prevents pathogenic organisms from adapting and mutating, meaning even after decades of use, fungal strains have not developed resistance to Manzate T&O Formulations.

	Formulations	Package Sizes
Manzate Pro-Stick T&O:	75 DF	8 x 6 pound
Manzate Max T&O T&O:	4 SC	2 x 2.5 gallon





630 Freedom Business Center, Suite 402, King of Prussia, PA 19406
1.800.438.6071 • www.upl-ltd.com

Manzate[®] Pro-Stick T&O

FUNGICIDE

Manzate[®] Max T&O

FUNGICIDE



FLOWERS, FOLIAGE PLANTS AND ORNAMENTALS Diseases Controlled

Alternaria	Curvularia	Gloeosporium	Melampsoridium	Ramularia
Asochyta	Cylindrosporium	Glomerella	Microsphaera alni	Rhizoctonia
Bipolaris	Diaporthe	Gnomonia	Monilinia	Rhytisma
Botryosphaeria	Didymascella	Guignardia	Monochaetia	Sclerotium
Botrytis	Diplocarpon	Gymnosporangium	Mycosphaerella	Sirococcus
Cephalosporium	Drechslera	Helminthosporium	Ovulinia	Sphaeropsis
Cercospora	Elsinoe	Kabatiella	Peronospora	Stemphylium
Ciborinia	Entomosporium	Keithia	Pestalotia	Stigmata
Cladosporium	Erwinia	Leptosphaeria	Phaeocryptopus	Taphrina
Coccomyces	Exobasidium	Lophodermium	Phoma	Uromyces
Colletotrichum	Fabraea	Marssonina	Phomopsis	Venturia
Coryneum	Fusarium	Melampsora	Phyllosticta	Whetzelinia
	Fusicladium		Puccinia	

Manzate T&O Fungicide is recommended for use on certain flower, foliage, and ornamental plants. See label for full list of plants. Please note the use of Manzate T&O has been found to be acceptable in specific genera and species of the plants listed on the product label; however, phytotoxicity may occur. Due to the large number of species and varieties of ornamentals and nursery plants, each one is not tested for sensitivity to Manzate T&O. The user should determine if Manzate T&O can be used safely prior to commercial use. Manzate T&O may also be used in aerial applications, as per label specifications.

Formulation	Use Rates*	Application Timing
Manzate Pro-Stick T&O	1-2 lbs. per acre or 100 gals. water	Typically begin application when disease threatens, or at the first sign of disease. Repeat application as per labeled interval rates as conditions persist.
Manzate Max T&O T&O	0.8-1.6 quarts per acre or 100 gals. water	

*Follow label instructions for disease specific use rates and application timing.



630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
1.800.438.6071 • www.upl-ltd.com