



# Stonewall® 0.37% Plus Fertilizer

PRODIAMINE	GROUP	3	HERBICIDE
------------	-------	---	-----------

For selective preemergence control of grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns and landscape ornamentals (including established perennials and wildflower plantings).

**ACTIVE INGREDIENT:**  
 Prodiamine: (N<sup>3</sup>, N<sup>3</sup>-Di-*n*-propyl-2,4-dinitro-6-(trifluoromethyl)-*m*-phenylenediamine).....0.37%  
**OTHER INGREDIENTS:**.....99.63%  
 Total: .....100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If swallowed:	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have a person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
If on skin or clothing:	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled:	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
Emergency Phone Numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 222-1222 Poison Control Center

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

**ENVIRONMENTAL HAZARDS**

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment wash water.

**NON-TARGET ORGANISM ADVISORY STATEMENT:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turf grasses (excluding golf course putting greens) and lawns;
- Landscape ornamentals;
- Established perennials and wildflower plantings.

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when it is activated by at least 1/2 inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

**Not for use on plants being grown for (1) Sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on ornamental plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.**

- DO NOT graze or feed livestock forage cut from areas treated with this product.
- DO NOT apply aerially.
- DO NOT apply to golf course putting greens.
- DO NOT apply this product through any type of irrigation system.
- DO NOT use on sod farms.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

**RESISTANCE MANAGEMENT**

For resistance management, this product is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local

extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.

- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

**SPRAY DRIFT ADVISORIES  
THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.  
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.**

**IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

- Controlling Droplet Size – Ground Boom**
- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
  - Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
  - Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

**BOOM HEIGHT – Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce. **SHIELDED SPRAYERS** Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

**BOOMLESS GROUND APPLICATIONS**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift. **HANDHELD TECHNOLOGY APPLICATIONS** Take precautions to minimize spray drift.

**TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation. **TEMPERATURE INVERSIONS** Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

**WIND**

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spr drift.

**INTEGRATED WEED PEST MANAGEMENT**

Integrate this product into an overall weed pest management strategy whenever the use of an herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Consult local agricultural and weed authorities for additional IPM strategies established for your area.

**WEEDS CONTROLLED**

When used in accordance with this label, this product will provide control of the following weeds:

Barnyardgrass	Goosegrass <sup>1</sup>	Panicum (Texas, Fall, Browntop)	Spurge, Prostrate
Bluegrass, Annual ( <i>Poa annua</i> )	Henbit	Pigweed	Witchgrass
Carpetweed	Itchgrass	Purslane, Common	Woodsorrel, Yellow (from seed)
Chickweed, Common	Johnsongrass (from seed)	Pusley, Florida	
Chickweed, Mouse ear (from seed)	Junglerice	Rescuegrass <sup>2</sup>	
Crabgrass (large, smooth)	Knotweed	Shepherd's Purse <sup>2</sup>	
Crowfootgrass	Kochia	Signalgrass, Broadleaf	
Cupgrass, Woolly	Lambsquarter, Common	Speedwell, Persian	
Foxtails, Annual	Lovegrass	Sprangletop	

<sup>1</sup> In many areas a single application of 0.65 lb. to 1.5 lb. active ingredient per acre (equal to 176 lbs. to 405 lbs./A of this product) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 0.65 lb. to 1.0 lb. a.i. per acre (equal to 176 lbs. to 270 lbs./A of this product) followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Annual Rates** section. Do not exceed the maximum rate for turf grass species listed in the **Maximum Annual Rates** section above.  
<sup>2</sup> Applications for this weed should be made in late summer, fall or winter prior to germination.  
<sup>3</sup> Suppression only. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

## 0-0-7

**GUARANTEED ANALYSIS**

SOLUBLE POTASH (K2O).....7.00%


DERIVED FROM: Muriate of Potash.

**APPLICATION DIRECTIONS**

Apply uniformly with suitable, calibrated application equipment.

**ESTABLISHED TURF**

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.



**USE RESTRICTIONS**

- The following restrictions apply to the use of this product in turf grasses and lawns:
- DO NOT** apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
  - DO NOT** cut (harvest) treated sod before 120 days after application.
  - DO NOT** apply to newly set sod until the following year.
  - DO NOT** apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.
  - DO NOT** allow people or pets to enter the treated area until the spray has dried.

**USE PRECAUTIONS:**

- The following precautions apply to the use of this product in turf grasses and lawns:
- Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
- Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as diking may result in reduced weed control.

**RATES OF APPLICATION**

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

**MAXIMUM ANNUAL RATES**

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate/Calendar Year of turf fertilizer by turf grass species(1)		
Turf Species	Lbs. Product/A	Lbs AI/Acre
Creeping Bentgrass	176	0.65
Creeping Red Fescue	203	0.75
Buffalograss	270	1.0
Kentucky Bluegrass		
Perennial Ryegrass		
Bermudagrass <sup>2</sup>	405	1.5
Bahiagrass		
Centipedegrass		
Seashore Paspalum		
St. Augustinegrass		
Tall Fescue (including turf type)		
Zoysia		

<sup>1</sup>These are the maximum rates per calendar year by species limitations.  
<sup>2</sup> May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 0.5 lb. a.i./acre (equal to 135 lbs./A of this product). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded.

Suppression only of Foxtail, Goosegrass, and Rescuegrass due to reduced product rates used in sprigging situations.

- Do not apply more than 1.5 lbs a.i. per calendar year per acre (equal to 405 lbs./A of this product).
- Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the **Maximum Annual Rates** table.

**WHEN TO APPLY AFTER OVERSEEDING TURF**

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

**WHEN TO OVERSEED AFTER APPLICATION**

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

**Overseeding Rate and Treatment Interval Chart**

Lbs. Product/Acre	Lbs a.I/A	Months Before Overseeding		
		North	Transition	South
135	0.50	4	4	4
176	0.65	5	4	4
203	0.75	6	5	5
216	0.80	-	6	6
270	1.00	-	7	7
308	1.14	-	-	9
351	1.30	-	-	10
405	1.50	-	-	12

**LANDSCAPE ORNAMENTALS**

Including Established Perennials and Wildflower Plantings

This product may be applied for residual preemergence weed control in ornamentals.

**Use Rates and When to Apply**

Apply in fall and/or spring, 0.65 to 1.5 lbs. active ingredient per acre, which is equal to 4.0 to 9.3 lbs. of this product per 1,000 sq. ft. of ornamental plantings; therefore 20 lbs of this product will treat 2,150 to 5,000 sq.ft. Use the higher rates of application for longer control periods. Sequential applications may be made as long as the total amount of product applied does not exceed the maximum annual application rate of 1.5 lbs. a.i. per acre.

**Established Landscape Ornamentals**

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate it into the soil.

**Newly Transplanted Landscape Ornamentals**

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

**Tolerant Ornamental Species**

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when it is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific Name	Common Name (Tested Varieties)	Scientific Name	Common Name (Tested Varieties)
<i>Abies</i> spp.	Fir	<i>Ligustrum amurense</i>	Amur Privet
<i>Abelia grandiflora</i>	Abelia; Sherwood	<i>Ligustrum japonicum</i>	Japanese Privet
<i>Acer palmatum</i>	Japanese Maple	<i>Ligustrum lucidum</i>	Glossy Privet (Wax-Leaf)
<i>Acer platanoides</i>	Norway Maple	<i>Liriope muscari</i>	Liriope (Big Blue, Lily Turf, Silver Mound, Evergreen Giant)
<i>Achillea</i> sp.	Yarrow (King Edward)	<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>Agapanthus africanus</i>	Lily-of-the-Nile; African Lily	<i>Lonicera tatarica</i>	Tatarian Honeysuckle
<i>Arctotheca calendula</i>	Cape Weed	<i>Malephora luteola</i>	Ice Plant
<i>Aucuba japonica</i>	Japanese Aucuba	<i>Malus</i> spp.*	Crabapple*
<i>Berberis gladywnensis</i>	Barberry	<i>Nandina domestica</i>	Heavenly Bamboo
<i>Berberis julianae</i>	Wintergreen Barberry	<i>Nerium Oleander</i>	Oleander
<i>Berberis thunbergii</i>	Japanese Barberry	<i>Olea europaea</i> *	Olive*
<i>Berberis verticillata</i>	Warty Barberry	<i>Ophiopogon japonicus</i> **	Mondo Grass**
<i>Buxus microphylla</i>	Japanese Boxwood	<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Callistemon viminalis</i>	Weeping Bottlebrush	<i>Oxydendron luteum</i>	Sourwood
<i>Calluna vulgaris</i>	Scotch Heather	<i>Persea Americana</i> *	Avocado*
<i>Carpobrotus edulis</i>	Hottentot Fig; Ice Plant	<i>Photinia fraseri</i>	Fraser's Photinia; Redtip
<i>Cassia artemisoides</i>	Feathery Cassia	<i>Picea</i> spp.**	Spruce species** (Colorado Blue, Norway, etc.)
<i>Ceanothus rigidus</i>	Wild Lilac	<i>Pieris japonica</i>	Lily-of-the-Valley Shrub
<i>Chamaecyparis pisifera</i>	False Cypress	<i>Pinus brutia</i>	Calabrian Pine
<i>Cleyera japonica</i>	Cleyera	<i>Pinus canariensis</i>	Canary Island Pine
<i>Coreopsis</i> spp.	Coreopsis; calliopsis (Early Sunrise, Moonbeam)	<i>Pinus elliotii</i>	Slash Pine
<i>Cornus florida</i>	Flowering Dogwood	<i>Pinus halepensis</i>	Aleppo Pine
<i>Cornus stolonifera</i>	American Dogwood	<i>Pinus nigra</i>	Austrian Black Pine
<i>Cortaderia selloana</i>	Pampas Grass	<i>Pinus radiata</i>	Monterey Pine
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster	<i>Pinus palustris</i>	Longleaf Pine
<i>Cotoneaster buxifolius</i>	Cotoneaster	<i>Pinus strobus</i>	Eastern White Pine
<i>Cotoneaster dammeri</i>	Bearberry Cotoneaster	<i>Pinus sylvestris</i>	Scotch Pine
<i>Cotoneaster microphyllus</i>	Rockspray Cotoneaster	<i>Pinus taeda</i>	Loblolly Pine
<i>Craeaegus</i> spp.	Hawthorne	<i>Pinus thunbergiana</i>	Japanese Black Pine
<i>Cupressus sempervirens</i>	Italian Cypress	<i>Pistacia</i> spp.**	Pistachio**
<i>Dodonea alba</i>	White Trailing Ice Plant	<i>Ptilosporum rhombifolium</i>	Queensland Pittosporum
<i>Dodonaea viscosa</i>	Hop Bush	<i>Ptilosporum tobira</i>	Japanese Pittosporum

(012921) Rev 9/8/22 VT

**Net Contents: 50 lb (22.7 kg) SGN 240**

**EPA REG. NO. 60063-40-10404**

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 69759-OR-004 (BAN), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 90561-NC-001 (N), 57131-WI-1-EC, 92626-TN-1, 49991-TX-2, 93243-CT-1, 93243-TX-1.

First letter of lot code indicates manufacturing site.

**#702155**

Distributed by: **LESCO, Inc.**  
 1385 East 36th Street  
 Cleveland, OH 44114-4114

**Tolerant Ornamental Species, continued:**

Scientific Name	Common Name (Tested Varieties)	Scientific Name	Common Name (Tested Varieties)
<i>Elaeagnus pungens</i>	Silverberry	<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Euonymus fortunei</i>	Wintercreeper	<i>Prunus laurocerasus</i>	English Laurel
<i>Euonymus japonica</i>	Japanese spindle tree, Evergreen Euonymus	<i>Prunus spp.*</i>	Almond, Apricot, Nectarine, Peach, Plum, and Prune*
<i>Euonymus kiautschovica</i>	Spreading Euonymus	<i>Pseudotsuga menziesii**</i>	Douglas Fir**
<i>Fatsia japonica</i>	Japanese Aralia	<i>Pyracantha coccinea</i>	Firethorn, scarlet
<i>Forsythia intermedia</i>	Border Forsythia	<i>Pyracantha fortuneana</i>	Firethorn
<i>Forsythia vindissima</i>	Greenstem Forsythia	<i>Pyracantha koidzumii</i>	Firethorn
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine	<i>Pyrus spp.</i>	Bradford Pear
<i>Ilex crenata</i>	Japanese Holly	<i>Quercus rubra</i>	Red Oak
<i>Ilex opaca</i>	American Holly	<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Jasminum nudiflorum</i>	Winter Jasmine	Rhododendron (including Azalea)	Rhododendron & Azalea (Coral Bells, Delaware valley white, Flame creeper, George L. Tabor, PJM, Roseum Elegans, Formosa, Hinocrimson, Wakaebisu, Gumpo)
<i>Juglans sp.*</i>	Walnut*	<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Juniperus chinensis</i>	Chinese Juniper (Nick's Compact)	<i>Rosmarinus officinalis*</i>	Rosemary*
<i>Juniperus conferta</i>	Shore Juniper (Blue Pacific)	<i>Rudbeckia, spp.</i>	Black-eyed Susan
<i>Juniperus horizontalis</i>	Creeping Juniper	<i>Sedum album</i>	Stonewall
<i>Justicia brandegeana</i>	Shrimp Plant	<i>Syzygium paniculatum</i>	Japanese Boxcherry
<i>Lagerstroemia indica</i>	Crape Myrtle	<i>Taxus cuspidata</i>	Japanese Yew
<i>Taxus media</i>	Yew	<i>Viburnum tinus</i>	Laurustinus
<i>Thuja occidentalis</i>	American Arborvitae	<i>Viburnum trilobum</i>	Cranberry Bush
<i>Trachelospermum asiaticum</i>	Star Jasmine	<i>Viburnum wrightii</i>	Leatherleaf Viburnum
<i>Tsuga canadensis</i>	Canada Hemlock	<i>Vinca minor</i>	Common Periwinkle
<i>Viburnum japonicum</i>	Japanese Viburnum	<i>Vitis spp.**</i>	Grape**
<i>Viburnum odoratissimum</i>	Sweet Viburnum	<i>Weigela florida</i>	Old Fashioned Weigela
<i>Viburnum plicatum</i>	Japanese Snowball	<i>Yucca aloifolia</i>	Spanish Bayonet
<i>Viburnum rigidum</i>	Canary Island Viburnum	<i>Yucca filamentosa</i>	Yucca, Adam's Needle

\* Do not use on food producing trees, vines or plants.  
\*\*Not for use on container grown plants.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.  
**STORAGE:** Store this product in its original container in a dry, cool, secured area.  
**PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authori-

**CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc.. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors. LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, INC. BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES. LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

If this product is being sold in a state requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available at the following website: <http://www.aapfco.org/metals.html>. LESCO is a registered trademark and Stonewall and the sweeping design are trademarks of LESCO Inc. SCOTT'S is a registered trademark of The SCOTT Company. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC.

(012921) Rev 9/8/22 VT

**Net Contents: 50 lb (22.7 kg) SGN**

**EPA REG. NO. 60063-40-10404**

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 69759-OR-004 (BAN), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 90561-NC-001 (N), 57131-WI-1-EC, 92626-TN-1, 49991-TX-2, 93243-CT-1, 93243-TX-1.

First letter of lot code indicates manufacturing site.

**#702155**

Distributed by: **LESCO, Inc.**  
 1385 East 36th Street  
 Cleveland, OH 44114-4114

**SPREADER SETTINGS**

Spreader settings vary by make and model of spreader. It is recommended that individual spreaders are calibrated for the specific product that is to be applied. A walking speed of 3 miles per hour is recommended.

Spreader Model	Application Rates <sup>1</sup> a.i. lbs./A (Product lbs./1000 sq ft)			
	0.5 (3.1)	0.75 (4.7)	1.0 (6.2)	1.5 (9.3)
LESCO® Calibration Gauge	#13	#16	#18	#24
PermaGreen	13	16	18	24
Cyclone® or Spyker®	3½	4½	4¾	5¾
Vicon (LESCO® Pendulum)	20	30	33	36
Lely®	3¼	4¼	4¾	5¾

<sup>1</sup> Application Rates for example only - do not exceed the maximum annual application rate of 1.5 lb a.i./acre. (9.3 lb product/1,000 sq. ft.)