

NITRO-PHOS

WEED & FEED 15-5-10

ACTIVE INGREDIENTS:

- Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid.....0.29%
- Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....0.17%
- Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid).....0.07%

OTHER INGREDIENTS:99.47%
 TOTAL:100.00%

THIS PRODUCT CONTAINS:

- 4.80 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.24%
- 2.80 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.14%
- 1.20 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.06%

Isomer Specific by AOAC Methods
 TRIMEC[®] is a registered trademark of PBI/GORDON CORP.

EPA Reg. No. 2217-603-54343
 EPA Est. No. 32802-MO-1(B)

First letter of code on bag shows establishment where manufactured.

STOP! READ THE ENTIRE LABEL FIRST.
 OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

FIRST AID

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call toll free, 1-877-800-5556, for emergency medical treatment information.	

**COVERS UP TO 15,625 SQ. FT.
 NET WT. 50 LBS. /22.73 KG.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

USE RESTRICTIONS:

- Do not exceed specified dosages for any area. Do not apply to newly seeded lawns. During application, close spreader openings when turning.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

USE PRECAUTIONS:

- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.

INSTRUCTIONS:

This product controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, herb, spurge and other listed weeds commonly found in home lawns. In addition, fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days. Do not use on dichondra. Do not overapply. Under conditions of adequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration of St. Augustine grass. Do not use this product on 'Floratam' St. Augustine grass.

1. Mow lawn to nonnal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non target plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, zoysiagrass, St. Augustine grass, Bahiagrass and centipedegrass: Apply 3.2 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application. The maximum broadcast application rate is 4.0 pounds of product per 1,000 square feet per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 square feet.

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage. Do not exceed the application rate for any area.

Typically a lawn utilizes 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When applied as directed, this product supplies 0.48 - 0.6 pounds of nitrogen per 1,000 square feet with each application. Use this product in conjunction with an additional fertilization program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your local agricultural extension agent for the proper amount of nitrogen to be applied in your area.

SPREADER SETTINGS

Central	7 1/2 to 8	Ortho	6	Scotts #35	6 to 7
Cyclone	4 to 5 1/2	Precision	6	Scotts PF1	6
Greenfield	7 to 8	Scotts C-1	G to H	Sears	7 to 8

THE ABOVE SPREADER SETTINGS ARE APPROXIMATE. CHECK YOUR SPREADER ON 250 SQ. FT. AND ADJUST TO APPLY AT THE RATE OF 1 LB. PER 250 SQ. FT.

WEEDS

Annual yellow sweetclover	Carolina geranium	Creeping Jenny	Horsenettle	Poison Oak	Sowthistle	Wild carrot
Aster	Chickweed, common	Curlydock	Lambsquarters	Prostrate knotweed	Spotted spurge	Wild garlic
Black eyed Susan	Chickweed, mouseear	Dandelion	Morningglory	Prostrate pigweed	Stinging nettle	Wild geranium
Black medic	Clover, crimson	Dichondra	Mustard	Prostrate spurge	Thisle	Wild onion
Broadleaf plantain	Clover, sweet	Dollarweed	Nettle	Prostrate vervain	Virginia creeper	Wild radish
Bull thistle	Clover, white	Florida pusley	Oxalis	Purslane, common	Russian thistle	Wild strawberry
Burning nettle	Cocklebur	Ground Ivy	Plantain	Russian thistle	Wayleaf bullthistle	Wild vetch
Buttonweed	Common mullein	Herbit	Poison Ivy	Sheep sorrel	Wild aster	Yarrow

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Distributed By:
 Nitro-Phos Co., Inc.
 P.O. Box 1356
 Houston, TX 77251

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)15%
1.96% Ammoniacal Nitrogen	
13.04% Urea Nitrogen	0.01%
AVAILABLE PHOSPHATE (P2O5)10%
SOLUBLE POTASH (K2O)10%

Chlorine: not more than7.60%

Secondary and/or Micro Plant Nutrient Guarantee:

BORON (B)0.02%
COPPER (Cu)0.05%
Water Soluble Copper (Cu)	
IRON (Fe)0.10%
0.01% Water Soluble Iron (Fe)	
MANGANESE (Mn)0.05%
0.01% Water Soluble Manganese (Mn)	
MOLYBDENUM (Mo)0.0005%
ZINC (Zn)0.05%
0.01% Water Soluble Zinc (Zn)	

Derived From: Ammonium Phosphate, Copper Oxide, Copper Sulfate, Iron Oxide, Iron Sulfate, Manganese Oxide, Manganese Sulfate, Sodium Borate, Sodium Molybdate, Zinc Oxide, Zinc Sulfate, Muriate of Potash, Urea.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapco.org/metals.htm>

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt, long pants, shoes plus socks and chemical-resistant gloves (made of any waterproof material). User should wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.