



Bayleton[®]
FLO

Product

information sheet

// Active Ingredients: Triadimefon (43%); equivalent to 4.15 lb triadimefon per gallon

// FRAC Code: 3 (DMI fungicides)

// Systemicity: Acropetal penetrant

// Formulation: Suspension concentrate

// Signal Word: Caution

Best Uses on Warm-Season Turf

// Dollar spot on seashore paspallum

Best Uses on Cool-Season Turf

// Dollar spot, fairy ring

Key Strengths

// Spring disease management programs, broad-spectrum disease control

Bayleton[®] FLO has a 30-year history of reliable disease control and continues to be a top choice for fairy ring and dollar spot management on turfgrass. Bayleton FLO can be used for simultaneous control of dollar spot and fairy ring in early-season programs and also provides broad-spectrum control of other important diseases.

Attributes of Bayleton FLO

// Systemic disease control of foliar, root and soil diseases

// Preventive and curative activity

// Long residual for dollar spot prevention

Solutions for Tough Diseases

// Proven fairy ring control

// Strong dollar spot control

// Early spring prevention DMI application

// Curative during the growing season

How to Use Bayleton FLO

Apply Bayleton FLO as part of an integrated fairy ring and dollar spot management program when average soil temperatures are 55-60°F at a 2-4-inch depth for 5 days or more. Use Bayleton FLO in seasonal spray programs for the control of other diseases. Avoid using Bayleton FLO under high temperature conditions.

Key Diseases Controlled ¹	Rate (per 1,000 sq ft) ¹	Application Interval
Dollar Spot ••	0.25 fl oz (preventive) ² 0.5 fl oz (preventive) 1.0 fl oz (curative)	14 days 30 days Once, then follow preventive rates and intervals
Red Thread • Powdery Mildew •• Rusts ••	0.5 fl oz (preventive) 1.0 fl oz (curative)	15 - 30 days Once, then follow preventive rates and intervals
Summer Patch •	1.0 - 1.4 fl oz (preventive)	Apply once in spring, 30 to 60 days before initial symptoms normally appear

How to Use Bayleton FLO (continued)

Key Diseases Controlled ¹	Rate (per 1,000 sq ft) ¹	Application Interval
Take-All Patch ³ •	1.0 - 1.4 fl oz	21 - 28 days, water-in after application
Fairy Ring ••	1.0 - 1.4 fl oz (preventive)	14 - 28 days
Brown Patch •	0.5 fl oz Bayleton FLO + 1.5 - 2.2 oz Prostar® 70 WG (preventive) 0.5 fl oz Bayleton FLO + 3.0 oz Prostar 70 WG (curative)	21 - 28 days/apply at curative rate, then follow preventive recommendations
Gray Leaf Spot ••	0.5 - 1.0 fl oz (preventive)	14 days
Large Patch •	1.0 - 1.4 fl oz	Apply once in early fall, then once in early spring

• = cool-season turf diseases

•• = warm-season turf diseases

¹See the Bayleton FLO label for the complete list of diseases controlled. Refer to the Bayleton FLO label for application information. Do not apply more than 1.9 fl oz/1,000 sq ft/application, do not exceed more than 1.4 fl oz/1,000 sq ft on residential lawns and do not exceed more than 3.8 fl oz/1,000 sq ft/year. Always read and follow label instructions carefully. ²Rate not labeled in California. ³Bayleton FLO is not labeled for take-all patch control in California.

Bayleton FLO Rate Chart

Use Rate (fl oz/1,000 sq ft)	Coverage per Container (1,000 sq ft)	Coverage per Container (acres)	lbs Triadimefon (per acre)	Applications at Rate (per year)
0.50	640	14.7	0.71	4
0.75	427	9.8	1.1	3
1.00	320	7.3	1.4	2
1.25	256	5.9	1.8	2
1.40	228	5.2	2.0	1

Package size: 2.5 gallon container.



Dollar spot affecting a Kentucky bluegrass lawn. Active mycelium is visible. Glenview, IL. (Bayer)

www.EnvironmentalScience.Bayer.US



ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS

Bayer Environmental Science, a Division of Bayer CropScience LP, 5000 CentreGreen Way, Suite 400, Cary, NC 27513. For additional product information, call toll-free 1-800-331-2867. Not all products are registered in all states. Bayer, the Bayer Cross, Bayleton, and Prostar are registered trademarks of Bayer. © 2019 Bayer CropScience LP. ES-818-BAYFLO-188-A-R2