

# LESCO® Barricade® 0.38% Plus Fertilizer

For selective preemergence control of grass and broadleaf weeds in established turfgrasses (excluding golf course putting greens), lawns, athletic fields, parks and sod farms, field grown and landscape ornamentals, established perennials and wildflower plantings, non-crop areas, including landscape beds, ornamentals, and established perennial wildflower plantings on or surrounding municipal grounds and government installations.

**ACTIVE INGREDIENT:**  
Prodimamine: (N3, N3-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine).....0.38%  
**OTHER INGREDIENTS:**..... 99.62%  
Total .....100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See the following sections Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> </ul>

**HOT LINE NUMBER**

For 24 Hour Medical Emergency Assistance ( Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), call CHEM-TREC 1-800-424-9300

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact CHEM-TREC 1-800-424-9300 for emergency medical treatment information.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if absorbed through the skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. [Wear protective eyewear]. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long sleeved shirt and long pants, sock, shoes, and gloves. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**Personal Protective Equipment (PPE)  
Mixers, loaders, applicators, and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any water proof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROL STATEMENTS**

When handlers used closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations:**

- Users should:**
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
  - Remove clothing /PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
  - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment wash waters.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. **Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

**NON AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. **Do not enter or allow others to enter until dusts have settled.**

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State/Tribal agency responsible for pesticide regulation. • For use on plants intended for aesthetic purposes or climatic modification. • Read entire label before using this product.


**ESTABLISHED TURF**

Golf courses (except putting greens), lawns, parks, athletic fields and sod farms.

**How to Apply**

- Do not apply more than 9.0 lb product/1000 sq feet (1.5 lb a.i./A) per calendar year.
- Do not make more than two applications per calendar year.
- Refer to Rates for Grass Species table and Granular Spreader Settings table for spreader settings to deliver the rate recommended for your lawn grass species. Uniform coverage is necessary to provide the most effective level of control.
- Apply LESCO Barricade 0.38% Plus Fertilizer to the turf in the early spring prior to crabgrass emergence. See Crabgrass Germination Dates in the map below for the approximate timing in your area. Always apply before weeds emerge. LESCO Barricade 0.38% Plus Fertilizer will not control emerged crabgrass.
- If you have a known serious weed problem or live in an area with an extended growing season, you may need to make a second application 60-90 days later.
- If you live in Zones D and E (see Crabgrass Germination dates map), you may apply a higher rate (Medium or High Rate) or make two applications for more effective crabgrass control, if recommended for your turfgrass species.
- Do not make a second application to Fine Fescue, Creeping Bentgrass, and newly sprigged and plugged Bermudagrass. Do not make a second application if you applied the High Rate to Buffalograss, Kentucky Bluegrass, or Perennial Ryegrass.
- For best control – After application, water in ½ inch of water within 14 days, either from watering or rain.

**Crabgrass Seed Germination Dates (Optional Graphic)**



**Rates for lawn grass species**

Turf Species	Application Schedule <sup>1</sup>	Application Rate <sup>2</sup>		
Bermudagrass (established)	First Application	Base	Medium	High
Bahiagrass	Second Application (optional)	Base or Medium or High	Base or Medium or High	Base or Medium or High
Centipedegrass				
Kikuyugrass	First Application	Base	Medium	High
Seashore Paspalum				
St. Augustinegrass	Second Application (optional)	Base or Medium	Base or Medium	-
Tall Fescue (including turf-type)	One Application Only	Base or Medium or High	-	-
Zoysiagrass				
Buffalograss	First Application	Base	Medium	High
Kentucky Bluegrass	Second Application (optional)	Base or Medium	Base or Medium	-
Perennial Ryegrass	One Application Only	Base or Medium or High	-	-
Fine Fescue (including Red Fescue)				
Creeping Bentgrass (½ inch and greater height) <sup>3</sup>	One Application Only	Base or Medium	-	-
Bermudagrass newly sprigged and plugged <sup>4</sup>	One Application Only	Base or Medium	-	-

1. A second application may be necessary because of heavy weed pressure or extended growing season.
2. See Granular Spreader Settings table for spreader settings to deliver the rate.
3. To avoid grass injury, do not apply LESCO Barricade 0.38% Plus Fertilizer to creeping bentgrass mowed at less than ½ inch in height.
4. Newly sprigged or plugged Bermudagrass rooting may be temporarily slowed.

<b>0-0-7 GUARANTEED ANALYSIS</b>	
SOLUBLE POTASH (K <sub>2</sub> O).....	7.00%
SULFUR (S) .....	2.20%
2.20% Combined Sulfur (S)	
DERIVED FROM: Sulfate of Potash.	
CHLORINE (Cl) Max.....	2.00%

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.  
**PESTICIDE STORAGE:** Store in a cool, dry place. Keep container closed and away from food, feedstuffs, and domestic water supplies.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling if available or dispose of bag in a sanitary landfill or by other procedures approved by state and local authorities.

**WEEDS CONTROLLED**

Barnyardgrass	Goosegrass <sup>s</sup>	Lovegrass	Speedwell, Persian
Bluegrass, Annual (Poa annua) <sup>1</sup>	Henbit	Panicum (Texas, Fall, Brown Top)	Springletop
Carpetweed	Itchgrass	Pigweed	Spurge, Prostrate
Chickweed, Common, Mousseear (from seed)	Johnsongrass (from seed)	Purslane, Common	Witchgrass
Crabgrass (large, smooth) <sup>2</sup>	Junglelice	Pusley, Florida	Woodsorrel, Yellow (from seed)
Crowfoot grass	Knotweed	Rescuegrass <sup>s</sup>	
Cupgrass, Woolly	Kochia	Shepherd's Purse	
Foxtails, Annual	Lambsquarter, Common	Signalgrass, Broadleaf	

1. In areas where Poa annua is a winter annual, apply LESCO Barricade 0.38% Plus Fertilizer (see rate table) in August or September to established, non-overseeded turf before Poa annua seeds germinate. These timings are approximate. Consult State Extension Service for more specific timing for your area.
2. Fall Applications for Spring Crabgrass Control in Cool-Season Grasses: In those areas where the ground freezes in the winter, LESCO Barricade 0.38% Plus Fertilizer can be applied in the fall after the soil temperature falls below 50F but before the ground freezes. This application will control crabgrass the following spring.
3. In many areas, a single application will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, make a second application 60-90 days later to ornamentals and lawns (if recommended for your lawn grass species).
4. Suppression only.

**USE PRECAUTIONS**

- Do not apply to putting greens or turfgrass areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (Poa Annua) are desirable species.
- Do not cut (harvest) treated sod before 90 days after application. To avoid turfgrass injury, do not apply to newly set sod until the sod has rooted into the second inch of soil and exposed edges have filled in.
- Application of LESCO Barricade 0.38% Plus Fertilizer to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- Seeding:**Do not apply to overseeded areas for 60 days after seeding. Or until after the second mowing, whichever is longer. Do not overseed for 4 months after last applications of this product.

**Granular Spreader Settings<sup>1,2,3</sup>**

Make and Models	[0.38 lb. ai/A] [2.3 lb./1,000 sq. ft] Base Rate (Treats xx sq. ft.)	[0.5 LB. ai/A] [3.0 lb./1,000 sq. ft] Medium Rate (Treats xx sq. ft.)	[0.75 lb. ai/A] [4.5 lb./1,000 sq. ft] High Rate (Treats xx sq. ft.)
LESCO Calibration Gauge	12	13	16
PermaGreen	12	13	16
Cyclone® or Spyrker®	3½	3½	4½
Vicon (LESCO Pendulum)	18	20	30

- 1 Assume 3 mph walking speed.
- 2 It is recommended that you calibrate your spreader before application since manufacturer's settings can vary among the same model of spreader. An extended Spreader Setting listing can be found at www.lesco.com.
- 3 Do not apply more than 1.5 lbs. ai/A per year.

**LANDSCAPE, FIELD GROWN ORNAMENTALS AND NON-CROP AREAS**

**How to Apply**

- Apply LESCO Barricade 0.38 Plus Fertilizer as a broadcast, over-the-top application at 3.9-9.0 lbs. product/1,000 sq. ft. Use higher rates when weed pressure was a severe problem in the past.
- Make a single application in fall and/or spring prior to weed emergence. Another application may be made 60-90 days later if necessary. Do not exceed a total of 9.0 lbs. product/1,000 sq. ft. per year.
- After application, water in ½ inch of water within 14 days, either from watering or rain or, incorporate into the top 1-2 inches of soil within 14 days after application.
- On established ornamentals, watering or rainfall soon after application will wash residues off plant foliage and activate LESCO Barricade 0.38% Plus Fertilizer in the soil.
- On newly-transplanted ornamentals, apply after the soil has settled around the transplants.
- For best weed control, apply to soil free of clods, weeds, and debris such as leaves.
- Prior to application, control existing weeds by hand weeding, cultivation, or by the use of an herbicide for emerged weeds.
- Plants under stress (such as heat, drought, or frost damage) may be sensitive to LESCO Barricade 0.38% Plus Fertilizer.

**TOLERANT ORNAMENTALS**

LESCO Barricade 0.38% Plus Fertilizer will not harm most trees, shrubs, vines and flowers. See the table below: Table 1. Tolerant Ornamentals – All States

Table 1. Tolerant Ornamentals – All States			
Scientific Name	Common Name	Scientific Name	Common Name
<i>Abies spp.</i>	Fir Species; Balsam, etc.**	<i>Maleophora luteola</i>	Ice Plant
<i>Acer palmatum</i>	Japanese Maple	<i>Malus sp.</i>	Crabapple*
<i>A. platanoides</i>	Norway Maple	<i>Nandina domestica</i>	Heavenly Bamboo
<i>Actinidia chinensis</i>	Kiwi* (Ornamental Only)	<i>Olea europaea</i>	Olive* (Ornamental Only)
<i>Agapanthus africanus</i>	Lily-of-the-Nile	<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita	<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Arctotheca calendula</i>	Cape Weed	<i>Oxydendrum arboreum</i>	Sourwood
<i>Aucuba japonica</i> Japanese	Japanese Aucuba	<i>Persea americana</i>	Avocado*
<i>Aucuba Persea americana</i>	Avocado*		
<i>Barberis gladywynensis</i>	Barberry	<i>Photinia fraseri</i>	Fraser's Photinia tip Fraser's Photinia (Redtip)
<i>B. julianae</i>	Wintergreen Barberry	<i>Picea spp.</i>	Colorado Blue, Norway,
<i>B. mentorensis</i>	Mentor Barberry	<i>Pieris japonica</i>	Lily-of-the-Valley Shrub
<i>B. thunbergii</i>	Japanese Barberry	<i>Pinus brutia</i>	Calabrian Pine
<i>B. verruculosa</i>	Warty Barberry	<i>P. canariensis</i>	Canary Island Pine

<i>Callistemon viminalis</i>	Weeping Bottlebrush	<i>P. halepensis</i>	Aleppo Pine
<i>Calluna vulgaris</i>	Scotch Heather	<i>P. nigra</i>	Austrian Black Pine
<i>Cassia artemisoides</i>	Feathery Cassia	<i>P. palustris</i>	Longleaf Pine
<i>Chamaecyparis pisifera</i>	False Cypress	<i>P. sylvestris</i>	Scotch Pine
<i>Cleyera japonica</i>	Cleyera	<i>P. taeda</i>	LOBLOLLY Pine
<i>Cornus florida</i>	Flowering Dogwood	<i>Pistachio sp.</i>	Pistachio*
<i>C. stolonifera</i>	American Dogwood	<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>Cortaderia selloana</i>	Pampas Grass	<i>P. tobira</i>	Japanese Pittosporum
<i>Cotoneaster spiculatus</i>	Cranberry Cotoneaster	<i>Podocarpus macrophylla</i>	Podocarpus sp.
<i>C. buxifolius</i>	Cotoneaster	<i>Prunus laurocerasus</i>	English Laurel
<i>C. dammeri</i>	Bearberry Cotoneaster	<i>Prunus spp.</i>	Almond, Apricot, Nectarine, Peach, Plum, and Prune*
<i>Cretaegus spp.</i>	Hawthorne*	<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>Cupressus sempervirens</i>	Italian Cypress	<i>P. fortuneana; koidzumii</i>	Firethorn spp.
<i>Delasperma alba</i>	White Trailing Ice Plant	<i>Pyrus sp.</i>	Bradford Pear sp.
<i>Dodonea viscosa</i>	Hop Bush	<i>Quercus spp.</i>	Oak spp.
<i>Elaeagnus pungens</i>	Silverberry	<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Euonymus fortunei</i>	Wintercreeper	<i>Rhododendron (including azalea)</i>	'Coral Bells' 'Formosa', 'Hino-Orism', 'IPJM', 'Roseum Elegans'
<i>E. japonica</i>	Evergreen Euonymus	<i>Rosa banksiae</i>	Lady Bank's Rosa
<i>E. klatuschovica</i>	Spreading Euonymus	<i>Rosmarinus officinalis</i>	Rosemary
<i>Fatsia japonica</i>	Japanese Aralia	<i>Santolina virens</i>	Santolina
<i>Forsythia intermedia</i>	Border Forsythia	<i>Sedum album</i>	Stonecrop
<i>F. viridissima</i>	Greenstem Forsythia	<i>Syzygium paniculatum</i>	Japanese Boxcherry
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine	<i>Taxus cuspidata</i>	Japanese Yew
<i>Hedera helix</i>	English Ivy	<i>T. media</i>	Yew, Var.
<i>Hibiscus rosa-sinensis</i>	Chinese Hibiscus	<i>Thuja occidentalis</i>	American Arborvitae
<i>Ilex cornuta</i>	Chinese Holly	<i>Tachelospermum asiaticum</i>	Star Jasmine
<i>I. crenata</i>	Japanese Holly	<i>Tsuga canadensis</i>	Common Hemlock
<i>I. opaca</i>	American Holly	<i>Viburnum japonicum</i>	Japanese Viburnum
<i>I. peryi</i>	Holly	<i>V. odoratissimum</i>	Sweet Viburnum
<i>Jasminum nudiflorum</i>	Winter Jasmine	<i>V. plicatum</i>	Japanese Snowball
<i>Juglans sp.</i>	Walnut*	<i>V. ngidum</i>	Canary Island Viburnum
<i>Juniperus chinensis</i>	Chinese Juniper	<i>V. tinus</i>	Laurustinus
<i>J. conferta</i>	Shore Juniper	<i>V. trilobium</i>	Cranberry Bush
<i>J. horizontalis</i>	Creeping Juniper	<i>V. wrightii</i>	Leatherleaf Viburnum
<i>Justica brandegeana</i>	Shrimp Plant	<i>Vinca minor</i>	Dwarf Periwinkle
<i>Lagerstromia indica</i>	Crape Myrtle	<i>Vitis sp.</i>	Grape*
<i>Ligustrum amurense</i>	Amur Privet	<i>Weigela florida</i>	Old Fashioned Weigela

\*Do not use on food-producing trees, vines, or plants. LESCO Barricade 0.38% Plus Fertilizer controls certain grass and broadleaf weeds before they emerge and provides residual control in established turfgrasses, and ornamentals and other non-crop area. LESCO Barricade 0.38% Plus Fertilizer inhibits shoot and root growth in susceptible weeds. LESCO Barricade 0.38% Plus Fertilizer will not control established weeds. Weed control is most effective when LESCO Barricade 0.38% Plus Fertilizer is activated by at least ½ inch of rainfall or irrigation before weed seeds germinate and within 14 days following application.

**CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc.. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors. LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, Inc., BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES. LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag. Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm> LESCO is a registered trademark and the sweeping design is a trademark of LESCO Technologies, LLC. Barricade is a trademark of a Syngenta Group Company. SCOTTS is a registered trademark of The SCOTT Company. Cyclone and Spyrker are registered trademarks of Spyrker Spreaders, LLC. N:\Regulatory\WPI\Regul Private\MstrLbL\ERIEVIEW ADDRESS\Landscape Style Master Lbls (071107) Rev 3/8/17 VT Net Weight: 50 lb (22.7 kg) SGN 240

EPA REG. NO. 100-1339-10404  
EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H)  
First letter of lot code indicates manufacturing site.

**F1560**  
Distributed by: **LESCO, Inc.**  
1385 East 36th Street  
Cleveland, OH 44114-4114

**#097305 PP**