



# Candor™

SELECTIVE HERBICIDE

## Control Trees and Brush with a Low Volatile Premix.

Candor™ herbicide can be used to control weeds in pastures and rangeland, Conservation Reserve Program (CRP) acres, non-irrigation ditchbanks, fence rows, roadsides and other non-crop areas and industrial sites. The post-emergence activity of Candor targets woody plants and brush – including blackberries and poison ivy – as well as annual and perennial broadleaves, while leaving desirable grasses unharmed.

### TOUGH WEED CONTROL

- Controls more than 100 tough annual and perennial broadleaf weed and woody plants without harming desirable grasses
- Proven active ingredients in a pre-mix formulation

### FLEXIBLE APPLICATION

- May be combined with liquid nitrogen fertilizer
- Can be applied by ground equipment, helicopter or by spot treatment
- No grazing restrictions for most livestock, including cattle, horses, gestating mares, sheep and goats
- Rainfast in 2 hours
- Non-restricted use pesticide

**EPA REG. NO.** 228-565

**ACTIVE INGR.** triclopyr (16.5%),  
2,4-D (34.4%)

**FORMULATION** liquid

**CHEM. FAMILY** pyridinecarboxylic acid

**GROUP NO.** 4

**MODE OF ACT.** synthetic auxin

**SIGNAL WORD** caution

**RESTRICTED** no

**PACKAGE SIZE** 4 x 1 gal, 2 x 2.5 gal



# Candor™

## SELECTIVE HERBICIDE

### KEY USES

Conservation Reserve Program (CRP) land, fallow systems, general farmstead, pastures, rangeland, fence rows, non-irrigation ditchbanks, roadsides, other non-crop areas and industrial sites

### KEY WEEDS CONTROLLED

(See product label for complete list)

Amaranth, spiny  
Blackberry  
Burdock  
Buttercup  
Chickweed, mouseear  
Clover, white sweet  
Clover, bur  
Cocklebur  
Croton, woolly  
Dandelion  
Dock, curly  
Dogbane, hemp  
Galinsoga, hairy  
Goatsbeard  
Henbit  
Horseweed  
Ironweed; tall, western  
Ivy, ground  
Kochia  
Kudzu  
Lambsquarters  
Lespedeza  
Lettuce, wild  
Mustard  
Oxalis  
Pennycress, field  
Pepperweed, field  
Pigweed, redroot  
Plantain  
Poison ivy  
Purslane, annual  
Ragweed  
Radish, wild  
Ragwort, tansy  
Shepherd's-purse  
Sneezeweed, bitter  
Sowthistle, annual  
Spurge  
Sunflower  
Thistle  
Vetch  
Violet, wild

### APPLICATION RECOMMENDATIONS

Candor™ can be applied in-season to foliage or during dormancy. For best results from foliar applications, apply when foliage is actively growing.

Rates range from 1 quart to 2 gallons per acre and vary depending on target weeds and timing of applications.

*See label for complete application rates and recommendations.*

### GRAZING AND HAYING RESTRICTIONS

#### GRAZING AND HARVESTING GREEN FORAGE

- No grazing restrictions for livestock including cattle, horses (including gestating mares), sheep, goats, etc.
- Lactating dairy animals - Do not graze or harvest green forage from treated area for 14 days after treatment.

#### HAYING (HARVESTING OF DRIED FORAGE)

- Livestock: Do not harvest hay for 7 days after treatment.
- Lactating dairy animals: Do not harvest hay until the next growing season.

*See label for complete grazing and haying restrictions.*

### SLAUGHTER RESTRICTIONS

Withdraw livestock from grazing treated grass or consumption of treated hay at least 3 days before slaughter. This restriction applies to grazing during the season of treatment or hay harvested during the season of treatment.

*See label for complete slaughter restrictions.*



More where it matters most.  
[www.nufarm.com/us](http://www.nufarm.com/us)

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit [www.nufarm.com/us](http://www.nufarm.com/us) to download a full label.  
©2013 Nufarm. Important: Always read and follow label instructions. Candor™ is a trademark of Nufarm. 13-VM-0011



Scan this code with your smartphone to visit CDMS and view the complete product label.