



Stonewall® 0.29% Plus Fertilizer

For selective preemergence control of grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns and landscape ornamentals (including established perennials and wildflower plantings). Contains LESCO® Poly Plus® Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding.

ACTIVE INGREDIENT: Prodimamine (N3, N3-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine) 0.29% INERT INGREDIENTS: 99.71% Total: 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION See the following sections: Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.

FIRST AID If swallowed: Call a poison control center or doctor immediately for treatment advice. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. If on skin or clothing: Take off contaminated clothing. If inhaled: Move person to fresh air.

HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec at 1-800-424-9300 for emergency medical treatment advice.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Causes moderate eye irritation. ENVIRONMENTAL HAZARDS This product has low solubility in water.


STORAGE AND DISPOSAL STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed or seed by storage or disposal. PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

DIRECTIONS FOR USE This product is a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in: Established turf grasses (excluding golf course putting greens) and lawns; Landscape ornamentals; Established perennials & wildflower plantings.

USE PRECAUTIONS The following use precautions apply to the use of this product in turf grasses and lawns: Application of this product may thin emerged annual bluegrass and newly over-seeded grasses. Do not apply to overseeded turf for 60 days after seeding or until after the second mowing, which ever is longer.

APPLICATION DIRECTIONS Apply uniformly with suitable, calibrated application equipment. ESTABLISHED TURF This product is a selective pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns.

LANDSCAPE ORNAMENTALS Including Established Perennials and Wildflower Plantings This product may be applied for residual preemergence weed control in ornamentals. Use Rates and When to Apply Apply in fall and/or spring, 0.65 to 1.5 lbs active ingredient per acre, which is equal to 224 to 517 lbs of this product per acre (5.1 to 11.9 lbs per 1,000 sq.ft.) of ornamental plantings; therefore 20 lbs of this product will treat 1,680 to 3,920 sq.ft.



RATES OF APPLICATION This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Table with 3 columns: Turf Species, lb LESCO Stonewall 0.29% Plus Fertilizer / acre, lb AI per Acre. Rows include Bermudagrass, Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue, Zoysia, Buffalograss, Kentucky Bluegrass, Perennial Ryegrass, Creeping Red Fescue, Creeping Bentgrass.

1. These are the maximum rates per calendar year by species limitations. 2. May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 172 lb/A. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded.

19-0-3 GUARANTEED ANALYSIS TOTAL NITROGEN (N) 19.00% SOLUBLE POTASH (K2O) 3.00% DERIVED FROM: Polymer Coated Urea, Urea, Muriate of Potash. CHLORINE (Cl) Max. 2.25% *5.13% Slowly available Urea Nitrogen from Polymer Coated Urea.

SUGGESTED SPREADER SETTINGS* Recommended Rates in lb per 1,000 sq ft. Table with columns for Spreader (LESCO - All Rotary Models, PermaGreen, Cyclone/Spyker, Vicon, Lely) and rows for application rates (#17, #18, #21, #18*, 4%, 4%*, 32, 34, 40, 34*, 4% II, 4% II, 5% II, 7 II).

Established Landscape Ornamentals This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate it into the soil. Newly Transplanted Landscape Ornamentals This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application.

Table with 4 columns: Scientific Name, Common Name, Scientific Name, Common Name. Rows include Abies spp., Abelia grandifolia, Acer palmatum, A. platanoides, Achillea sp., Agapanthus africanus, Arctotheca calendula, Aucuba japonica, Berberis gladiwynensis, Berberis julianae, Berberis thunbergii, Berberis verruculosa, Buxus microphylla, Callistemon viminalis, Calluna vulgaris, Carpoprotus edulis, Cassia artemisoides, Ceanothus rigidus, Chamaecyparis pisifera, Clearya japonica, Coreopsis spp., Comus florida, Cornus stolonifera, Cortaderia selloana, Cotoneaster apiculatus, Cotoneaster buxifolius, Cotoneaster dammeri, Cotoneaster microphyllus, Cretaegus spp., Cupressus sempervirens, Delasperra alba, Dodonea viscosa, Eleaegnis pungens, Euonymus fortunei, Euonymus japonica, Euonymus kiautschovica, Fatsia japonica, Forsythia intermedia, Forsythia vindissima, Gardenia jasminoides, Ilex crenata, Ilex opaca, Jasminium nudiflorum, Juglans sp., uniperus chinensis, Juniperus conferta, Juniperus horizontalis, Justicia brandegeana, Lagerstromia indica, Rhododendron (including azalea).

Table with 4 columns: Scientific Name, Common Name, Scientific Name, Common Name. Rows include Ligustrum amurense, Ligustrum japonicum, Ligustrum lucidum, Liriope muscar, Lonicera japonica, Lonicera tatarica, Maleophora luteola, Malus spp., Nandina domestica, Nerium Oleander, Olea europaea, Ophiopogon japonicus, Osteospermum fruticosum, Persea Americana, Photinia fraseri, Picea spp., Plenis japonica, Pinus brutia, Pinus canariensis, Pinus elliotii, Pinus halepensis, Pinus nigra, Pinus radiata, Pinus palustris, Pinus strobus, Pinus sylvestris, Pinus taeda, Pinus thunbergiana, Pistachio sp., Pittosporum rhombifolium, Pittosporum tobira, Podocarpus macrophyllus, Prunus laurocerasus, Prunus spp., Pseudotsuga menziesii, Pyracantha coccinea, Pyracantha fortuneana, Pyracantha koidzumii, Pyrus spp., Quercus rubra, Raphiolepis indica, Rosa banksiae, Rosmarinus officinalis, Rudbeckia, Sedum album, Taxus cuspidata, Taxus cuspidata, Yucca aloifolia, Yucca aloifolia, Yucca aloifolia, Yucca aloifolia.

Table with 4 columns: Tuja occidentalis, American Arborvitae, Viburnum trilobium, Cranberry Bush; Trachelospermum asiaticum, Star Jasmine, Viburnum wrightii, Leatherleaf Viburnum; Tsuga canadensis, Canada Hemlock, Vinca minor, Common Periwinkle; Viburnum japonicum, Japanese Viburnum, Vitis spp., Grape; Viburnum odoratissimum, Sweet Viburnum, Weigela florida, Old Fashioned Weigela; Viburnum plicatum, Japanese Snowball, Yucca aloifolia, Spanish Bayonet.

WEEDS CONTROLLED When used in accordance with this label, this product will provide control of the following weeds: Barnyardgrass, Bluegrass, Annual (Poa annua), Carpetweed, Chickweed, Common, Chick weed, Mouse ear (from seed), Crabgrass (large, smooth), Crowfoot grass, Cupgrass, Woolly, Foxtails, Annual, Goosegrass, Henbit, Johnsongrass, Junglence, Knotweed, Kochia, Lambsquarter, Common, Lovegrass, Panicum (Texas, Fall, Brown Top), Pigweed, Purslane, Common, Pusley, Florida, Rescuegrass3, Shepherd's Purse2, Signalgrass, Broadleaf, Speedwell, Persian, Sprangletop, Spurge, Prostrate, Witchgrass.

1. In many areas a single application of 224 to 517 lb/a of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 228 to 345 lb/a followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate Table above.

WHEN TO APPLY AFTER OVERSEEDING TURF Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seeding secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

Table with 5 columns: LESCO Stonewall 0.29% Plus Fertilizer Rate (lb/A), Product, Months Before Overseeding (North, Transition, South). Rows include Product 172, 224, 259, 276, 345, 393, 448, 517.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc.. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag. Information regarding the contents and levels of metals in this product is available on the internet at: http://www.aapfco.org/metals.htm.

Net Weight: 50 lb 22.7 kg SGN 240 EPA REG. NO. 60063-41-10404 EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H). #702075 PP F1560 Distributed by: LESCO, Inc. 1385 East 36th Street Cleveland, OH 44114-4114