



LockUp™ Extra 2 with Fertilizer

For postemergence control of annual and perennial broadleaf weeds in established turfgrass, including residential lawns, golf courses, parks, sports fields, sod farms and around commercial buildings and other recreational and commercial turfgrass areas.

Contains LESCO® Poly Plus® Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding.

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid.....0.04%

Penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy[1,2,4] triazol[1,5c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide.....0.01%

dicamba: 3,6-dichloro-o-anisic acid.....0.08%

OTHER INGREDIENTS:98.87%

TOTAL:100.00%

Acid equivalent: 2,4 – D acid – 0.86% - 17.2 lb/ton dicamba – 0.08% - 1.5 lb/ton

Engineering Controls

When handlers use closed systems and enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)].

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

User Safety Recommendations:

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticides gets on skin, wash immediately with soap and water.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

PRODUCT INFORMATION

LESCO LockUp™ Extra 2 With Fertilizer is a stable combination of three herbicides. It provides postemergence control of annual, biennial and perennial broadleaf weeds in established turfgrass, including, residential lawns, golf courses, parks, sports fields, sod farms and around commercial buildings and other recreational and commercial turfgrass areas.

ENVIROMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate. 2,4-D has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted entry interval and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls worn over short-sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Protective eyewear

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Entry Restrictions for Non-WPS Uses: Do not enter or allow others to enter until dusts have settled.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **Pesticide Storage:** Store in a cool dry place in original container. **Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. **Container Handling:** Non-Refillable Container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

PRODUCT USE RESTRICTIONS

- Do not apply LESCO LockUp™ Extra 2 With Fertilizer on dichondra, carpetgrass, St. Augustinegrass, bahiagrass, or to golf course putting greens or tees, or turfgrass containing desirable clovers.
- Do not apply LESCO LockUp™ Extra 2 With Fertilizer directly to, or otherwise permit LESCO LockUp™ Extra 2 With Fertilizer to come into contact with, flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious plant injury may occur.
- Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at application.
- Do not collect treated grass clippings for use as mulch around plants.
- Do not water, mow or disturb treated area for 48 hours after application.
- Use LESCO LockUp™ Extra 2 With Fertilizer on solid (monoculture) stands of perennial ryegrass or tall fescue at a maximum rate of 175 lb per acre.
- Do not apply more than 175 lb of LESCO LockUp™ Extra 2 With Fertilizer per acre per application.
- Do not make more than two applications of LESCO LockUp™ Extra 2 With Fertilizer per acre per year.
- The re-treatment interval is 21 days for applications to turfgrass grown for sod or seed.

PRODUCT USE PRECAUTIONS

- The active ingredient in this product may leave residues in treated grass clippings which could be harmful to some plants and trees. The applicator must notify in writing the appropriate residences/individuals of this precaution when this product is applied by a lawn service/commercial applicator.
- To minimize the potential for turfgrass injury, allow four weeks between applications.
- Some yellowing of these turfgrasses may occur within two weeks of application. Any discoloration is transitory and will dissipate under adequate growing conditions.
- Apply LESCO LockUp™ Extra 2 With Fertilizer to newly seeded or sprigged turfgrass only after the third mowing or after the turfgrass seedlings have developed an adequate secondary root system. Reseeding is not recommended for at least three to four weeks after application.

Treatment on Turfgrass Species Not Listed on the Label for LESCO LockUp™ Extra 2 With Fertilizer
Users who wish to use LESCO LockUp™ Extra 2 With Fertilizer on turfgrass species not recommended on this label may determine the suitability for use by treating a small area at a specified rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the target turfgrass species is tolerant. Applications should not be made to turfgrass under stress. The user assumes responsibility for any plant damage or other liability resulting from use of LockUp™ Extra 2 With Fertilizer on turfgrass species not listed on this label.

LESCO LockUp™ Extra 2 With Fertilizer is recommended for use on the following established turfgrass species:

Common Name	Scientific Name
Annual bluegrass	<i>Poa annua</i>
Bluegrass, Kentucky	<i>Poa pratensis</i>
Fescue, chewing	<i>Festuca rubra var. commutata</i>
Fescue, creeping red	<i>Festuca rubra</i>
Fescue, sheeps	<i>Festuca ovina</i>
Fescue, tall (1)	<i>Festuca arundinaceae</i>
Ryegrass, perennial (1)	<i>Lolium perenne</i>

Established Warm Season Turfgrass

Common Name	Scientific Name
Bermudagrass (common or hybrid)	<i>Cynodon dactylon</i>
Centipedegrass (2)	<i>Eremochloa ophiuroides</i>
Fescue, tall (growing in warm season areas) (1)	<i>Festuca arundinaceae</i>
Kikuyugrass	<i>Pennisetum clandestinum</i>
Zoysiagrass	<i>Zoysia japonica</i>
Zoysiagrass	<i>Zoysia tenuifolia</i>

1. Use LESCO LockUp™ Extra 2 With Fertilizer on these species at a maximum rate of 175 lb (0.02 lb ai penoxsulam) per acre per application and only if some injury can be tolerated. A slight yellowing of turfgrass may occur within two weeks of application. Any discoloration is transient and is generally gone within four weeks of treatment under adequate growing conditions.
2. Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at application.

Weeds Controlled or Suppressed

Generally, application rates at the lower end of the rate range will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands and/or larger weeds), the higher rate within the rate range or repeat applications will be needed. Weeds growing in the absence of competition from other vegetation generally require higher rates or repeat applications to obtain satisfactory control or suppression.

Weeds Controlled or Suppressed		Rate	
		lb/acre	lb/1000 sq ft
bedstraw	medic, black morningglory	150-175	3,4-4
burdock	peppergrass		
chickweed	pigweed		
chicory	plantain		
clover	plantain, broadleaf		
dandelion	plantain, buckhorn		
dock	purslane		
dollarweed	ragweed		
elipta	sheepsorrel		
healall	shepherd's purse		
ivy, ground	speedwell		
knotweed	thistle		
lambsquartars			

Application Directions

Apply LESCO LockUp™ Extra 2 With Fertilizer using a drop or rotary-type spreader designed to apply granules. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. To optimize product performance, mow the turfgrass two to three days before applying LESCO LockUp™ Extra 2 With Fertilizer. If the turfgrass is under moisture stress, a thorough sprinkler irrigation of at least ½ inch may be needed at least one to two days before application. For optimal results, apply LESCO LockUp™ Extra 2 With Fertilizer in the morning when dew is present. If necessary, sprinkler irrigate with less than ¼ inch of water just prior to application to maximize particle adherence to leaf surfaces. Avoid watering of treated area for 24 hours following application. Avoid making applications if rainfall is expected within 24 hours after treatment.

Application Techniques for Applying LESCO LockUp™ Extra 2 With Fertilizer

Application Techniques for Applying LESCO LockUp™ Extra 2 With Fertilizer
 •When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution.
 •A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
 •For treating smaller areas, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.
 •A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully measure the amount of product needed to avoid over application.

Spreader Settings as a Guide for Calibration

Note: These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by LESCO or the spreader manufacturer.

Spreader	Settings for Spreader	
	Rate of LESCO LockUp™ Extra 2 With Fertilizer	
	150 lb/acre	175 lb/acre
LESCO – All Rotary Models	14	15
PermaGreen	14	15
Cyclone®/ Spyker® /Z-Spray	4¼	4½
Lely	4¼	4½

Rates and Timing of Application

Apply 150 to 175 lb of LESCO LockUp™ Extra 2 With Fertilizer per acre (3.4 to 4 lb per 1000 sq ft) per application to established turfgrass. Possible re-treatment in approximately 30 days may be necessary if turfgrass is heavily infested with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed 350 lb of LESCO LockUp™ Extra 2 With Fertilizer per acre (8 lb per 1000 sq ft.) per year because damage to turfgrass may occur. Do not make more than two applications per season. **Note:** When applying to warm season turfgrass under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended to reduce application rates by 50% during hot, dry conditions to avoid possible dehydration or browning of turfgrass. Best weed control results are obtained when LESCO LockUp™ Extra 2 With Fertilizer is applied to small, actively growing weeds in the spring or early fall. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control. Broadleaf weed species germinate at different times.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.
Warranty Disclaimer
 Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.
Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies
 To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
 2. Replacement of amount of product used.
- To the extent permitted by law, Seller shall not be liable for losses or damages resulting from handling or use of this product, unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapico.org/metals.htm>. Poly Plus is comprised of Polymer Coated Urea. LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC. Lely is a registered trademark of C Van Der Lely N.V. Vicon is a registered trademark of Ransomes-Cushman-Ryan. Lockup is a registered Trademark of Dow Agro Sciences.

Rev. 6/20/17 VT (012916)

Net Weight: 50 lb (22.7 kg) SGN 240

EPA REG. NO. 62719-590-10404

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 61842-CA-1 (T), 37351-CO-1 (E), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H).

First letter of lot code indicates manufacturing site.

F1560
 Distributed by: **LESCO, Inc.**
 1385 East 36th Street
 Cleveland, OH 44114-4114

#702422 PP

18-0-4

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)18.00%

18.00% Urea Nitrogen*

SOLUBLE POTASH (K₂O).....4.00%

IRON (Fe)5.00%

0.05% Water Soluble Iron (Fe)

DERIVED FROM: Polymer Coated Urea, Urea, Muriate of Potash, Iron Sucrate.

CHLORINE (Cl) Max.....3.00%

*13.50% Slowly available Urea Nitrogen from Polymer Coated Urea.